GENERAL ELECTRIC

Group Technical Resources Operation

INSULATION SYSTEMS LABORATORY OPERATION
100 WOODLAWN AVENUE
PITTSFIELD, MASS. 01201
A/C 413, Tel. 494-4385

SUBJECT:
ARTICLE IN SKIN AND
ALLERGY NEWS - SEPTEMBER, 1973

September 25, 1973

Mr. Paul G. Benignus
Monsanto Chemical Company
800 No. Lindbergh Blvd.
St. Louis, Missouri 63166

Dear Paul:

Enclosed is the page from the above publication with an article by a Dr. Bickers, with the title "Polychlorinated Biphenyls Induce Enzyme Production".

Our medical director does not object to the main content of the article except for the second paragraph which states that polychlorinated biphenyls are carcinogens. Inasmuch as your trademark name, Aroclor, is also cited, we believe that you may want to have your company protest this particular statement.

Very truly yours,

E. L. Raab, Manager
DIELECTRIC SYSTEMS LABORATORY
Building 11, Room 315

ELR/ps

Enclosure

cc: Dr. F. G. McKerr
E. L. Dobbins

DSW 018888

STLCOPCB4005432

Polychlorinated Biphenyls Induce Enzyme Production

International Medical News Service

NEW YORK — Topically applied polychlorinated biphenyls induce production of the enzyme aryl hydrocarbon hydroxylase in both skin and liver, Dr. D. R. Bickers reported at the annual meeting of the Society for Investigative Dermatology.

Polychlorinated biphenyls (PCB) are carcinogens which are widely distributed as environmental pollutants.

They are found in human tissues and are potent inducers of the hepatic mixed function oxidase system that metabolizes drugs and polycyclic hydrocarbons, said Dr. Bickers, of New York University School of Medicine and The Rockefeller University, New York.

There was a twelve-fold increase in aryl hydrocarbon hydroxylase (AHH) in the treated nuchal skin of rats and a ten-fold increase in their liver AHH levels. Untreated dorsal skin of the same animals



Dr. Bickers

showed a much lower level of AHH induction (about three-fold), he reported.

The polychlorinated biphenyl AROCLOR 1254 was applied to shaved nuchal skin of rats on three successive days. After the animals were killed, skin from both the treated and untreated area and liver tissue were analyzed.

Cytochrome P-450 was found in the treated skin of these animals.

The P-450 is the terminal oxidase of the hepatic mixed function oxidase system that metabolizes drugs and polycyclic hydrocarbons. It has an absorption peak at 450 nm on the carbon monoxide difference spectrum.

Dr. Bickers' associates in this study were Drs. L. C. Harber and A. Kappas and A. P. Alvares, Ph.D.

DSW 018889

STLCOPCB4005433