

C. A case-
strie disor-
to solvents
(1976) 14-

Current
ers: la
ational
Publishers
237-248.
neurology
n 1969.
berger PA
duced Par-
(1978) 222-

ity of 2,4-
2, 4, 5-tri-
ort on their
dogs. Arch
-67.
n JR. Peri-
sure to ad-
ssoc (1959)

pidemiologic
term effects
s. Scand j
63-169.
nderson R
ers exposed
quantitative
cular func-
(1978) 976-

medical and
s indicating
sulfide ab-
(1941) 859-

- of car-
in cate-
armacol 35

d estimation
j epidemiol

ural history
d j 2 (1972)

professionale

significant
mpling pro-
retrospective
s sizes. Am

onismus mit
rovaskulären
er. Scheitel-
n Gewerbe-
42-552.

Relationship between clinical and electrophysiological findings and indicators of heavy exposure to 2,3,7,8-tetrachlorodibenzo-dioxin

02-03-021

by Graziella Filippini, MD, Bianca Bordo, MD, Paolo Crenna, MD,
Nicoletta Massetto, MD, Massimo Musicco, MD, Renato Boeri, MD

FILIPPINI G, BORDO B, CRENN A P, MASSETTO N, MUSICCO M, BOERI R. Relationship between clinical and electrophysiological findings and indicators of heavy exposure to 2,3,7,8-tetrachlorodibenzo-dioxin. *Scand j work environ health* 7 (1981) 257-262. In this study the prevalence rate of peripheral neuropathy in a population living in an area polluted with 2,3,7,8-tetrachlorodibenzo-dioxin (dioxin-TCDD) was determined. Of the 723 subjects invited to the first screening in 1977, 470 (65 %) attended. At the second screening in 1978, of the 710 invited subjects, 319 (45 %) attended. Prevalence rate ratios for peripheral neuropathy and the associated 95% confidence limits were calculated for subgroups determined by the presence of (i) predisposing factors to neuropathy (alcoholism, diabetes, occupational exposure to neurotoxic agents, etc) or (ii) conditions thought to result from exposure to dioxin-TCDD such as chloracne or abnormal serum hepatic enzyme levels. The prevalence rate of peripheral neuropathy among those subjects with predisposing factors and among those with chloracne or abnormal serum hepatic enzyme levels was nearly three times greater than among those without these manifestations. The results derived from this study may be useful qualitative pointers for identifying subjects at risk in the neurological follow-up.

Key terms: chloracne, dioxin, environmental pollutants, liver damage, peripheral neuropathy, raised serum hepatic enzyme levels.

Subjects with occupational or accidental exposure to 2,3,7,8-tetrachlorodibenzo-dioxin (dioxin-TCDD) have been found to develop a wide variety of lesions and symptoms. Neurological findings may include polyneuropathies, lower extremity weakness, or sensorial impairment (sight, hearing, smell, taste), all of which have been recorded for dioxin-exposed subjects in whom other long-term effects such as chloracne, raised levels of cholesterol, triglyceride, gamma-glutamyl transpeptidase (GGT), glutamic oxalacetic transaminase (GOT), and glutamic pyruvic transaminase (GPT) were manifest (1, 3, 6, 7, 10, 11, 12, 13, 14, 17, 18). An association

of chloracne with increased GGT, GOT, and GPT activities and abnormal lipid levels has also been reported (19, personal communication from M Moses). The rate of reported congenital anomalies in men fathering at least one child after service in Viet Nam was nearly twice as high among those with recurrent acne, skin rash with blisters, and skin color changes consistent with conditions thought to result from exposure to dioxin-contaminated substances as among those without them (15). For men with these "indicators of exposure" greater frequencies have also been reported for gastrointestinal disturbances, swelling or numbness of joints, and sleep and psychological disturbances (15).

These previous findings prompted us to compare the prevalence rate (PR) of peripheral neuropathy in exposed subjects

Istituto Neurologico "C. Besta," Milan, Italy.

Reprint requests to: Dr G Filippini, Istituto Neurologico "C. Besta," Via Celoria 11, I-20133 Milan, Italy.

with chloracne or increased γ -GT, GOT, and GPT activities with that found in exposed subjects without these manifestations in a population living in a dioxin-polluted area in the Seveso region of Italy.

Subjects and methods

In July 1976, an accident at a trichlorophenol-producing factory in Meda, Italy, resulted in the contamination of a large, densely populated area. Seven hundred and thirty-three residents were evacuated 15 to 20 d after the accident from the most polluted zone (an area of 108 ha with an average dioxin-TCDD level of $240 \mu\text{g}/\text{m}^2$, a maximum level of $5,477 \mu\text{g}/\text{m}^2$, and an unevaluable minimum level) (4). These residents were invited by letter and personal invitation during visits at home to an initial neurological screening in March 1977 and to a second screening in April 1978. At the first screening five subjects had died and five had moved from the area. Of the 723 inhabitants invited to participate in the screening in Seveso, 470 (65 %) attended. During the period March 1977 to April 1978, five died and nine moved from the area. Of the 709 inhabitants invited to the second screening, 319 (45 %) attended, 277 of whom had been examined at the first screening and 42 of whom were examined for the first time. Of the latter 42, 11 children under the age of 5 refused to participate in the electrophysiological examination and were thus excluded from the prevalence calculation. No acute polyneuropathy was found during the study.

Each subject was examined according to a standardized protocol requiring a medical history questionnaire, a clinical examination, and an electrophysiological investigation. The neurological symptoms specifically identified were paresthesia, hypesthesia, pain, and hyposthenia, and the clinical signs were superficial and deep sensory impairment, muscular weakness, and tendon hypo- or areflexia. The electrophysiological study included maximum and minimum motor conduction velocity along the ulnar (axilla-wrist) and peroneal (capitulum fibulae-ankle) nerves and needle electromyography on the first interosseous muscle. However, electromyography was abandoned during the course

of the study because many subjects rejected the use of the needle. Muscle potentials were recorded by a surface electrode (Disa 13 K 60) from the minimum abductor and extensor digitorum brevis muscles at a surface temperature above 27°C . Minimum conduction velocity was determined by the desynchronization method of Cosi et al (5). Motor conduction latencies were corrected to a standard distance of 6 cm. A Disa 1500 electromyograph (Disa Elektronik, Skovlunde, Denmark) was used.

A nonexposed population living in an unpolluted area was used as a source of "normal" reference values of the electrophysiological parameters. A total of 1,000 subjects (91.6 % of the randomized sample) were representative of all ages, sex, and occupation. Seventy-five of these subjects were alcoholic, occupationally exposed to neurotoxic agents, or had diseases predisposing to peripheral neuropathy and were thus excluded from the calculation of the reference values. The "normal" reference values obtained are given by age group in table 1. Conduction velocity and distal latency were considered to be in the pathological range when the former was lower than the mean -2 SD and the latter was higher than the mean $+2$ SD of the reference population of the same age group. These two parameters were normally distributed. Amplitude and duration of muscle action potential were considered abnormal when the former was below the lower range limit and the latter was above the upper range limit of the reference population of the same age group.

For an electrophysiological diagnosis of peripheral neuropathy two or more abnormal parameters, among which conduction velocity was obligatory, were considered necessary, regardless of the involved nerve. All Seveso subjects were investigated for the following factors predisposing to neuropathy: alcoholism (more than 1 l of wine consumed daily), personal history of diabetes or fasting plasma glucose $\geq 100 \text{ g}/100 \text{ ml}$, cancer, nutritional deficiency, infectious and inflammatory diseases, osteoarticular diseases, use of neurotoxic drugs, possible occupational exposure to neurotoxic agents (chemical workers, shoemakers, plastic workers, painters, cabinet makers, printers, tanners, ceramic work-

ers). According to the International Classification of Diseases (1), chloracne was diagnosed when γ -GT was above 27°C . Minimum conduction velocity was determined by the desynchronization method of Cosi et al (5). Motor conduction latencies were corrected to a standard distance of 6 cm. A Disa 1500 electromyograph (Disa Elektronik, Skovlunde, Denmark) was used.

The statistical analysis was performed within the experimental design of the relationship between peripheral neuropathy and indicators of dioxin being tested. The prevalence rate ratio (P) was calculated for subjects with chloracne and subjects without them. Ninety-five percent confidence limits (95 %) were calculated according to the Woolf (20).

Table 1. Motor conduction values (surface electrodes) in

Nerve	Number of referents	Age (n)	Me
Ulnaris (axilla-wrist)	172	6-29	57
	118	30-59	58
	15	≥ 60	59
Peroneal (capitulum fibulae-ankle)	168	6-29	57
	115	30-59	58
	12	≥ 60	41

Table 2. Prevalence rate groups of Seveso residents

Actual cases of peripheral neuropathy
Total number of subjects
PR (%)
PRR
95 % confidence limits

• Evaluated versus group

subjects re-
Muscle action
a superficial
from the digit
for digitorum
temperature
tion velocity
achronization
Motor distal
to a standard
a 1500 electro-
tik, Skovlunde,

on living in an
as a source for
s of the electro-
A total of 380
ed sample) sub-
all ages, parti-
these subjects
nally exposed to
diseases predis-
opathy and were
calculation of the
normal" reference
n by age groups
velocity and distal
d to be in the
the former was
2 SD and the lat-
mean + 2 SD of
of the same age
eters were nor-
ode and duration
were considered
mer was below the
he latter was above
f the reference pop-
e group.

logical diagnosis of
two or more ab-
ong which conduc-
igatory, were con-
ardless of the in-
eveso subjects were
ollowing factors pre-
ny: alcoholism (more
med daily), personal
fasting plasma glu-
er, nutritional defi-
inflammatory diseases,
s, use of neurotoxic
ational exposure to
emical workers, shoe-
ers, painters, cabinet
ners, ceramic work-

ers — according to the International Stan-
dard Classification of Occupations (9)).
Chloracne was diagnosed by a dermatol-
ogist.

Increased enzyme activities were re-
corded when GPT was above 50 U/l (nor-
mal range: up to 36 U/l, as determined ac-
cording to Szasz (16)), GOT was above 50
U/l [normal range: up to 26 U/l, as deter-
mined by the "optimized method" (2)], and
GPT was above 50 U/l [normal range: up
to 32 U/l, as determined by the "optimized
method" (2)].

The statistical analysis of the data was
performed within the exposed population,
the relationship between peripheral neu-
ropathy and indicators of heavy exposure
to dioxin being tested. The PR and the
prevalence rate ratio (PRR) of peripheral
neuropathy were calculated for the Seveso
subjects with chloracne or raised serum
hepatic enzyme levels versus Seveso sub-
jects without them. Ninety-five percent
confidence limits (95 % CL) were estimat-
ed according to the method described by
Wolf (20).

Results

Relationship between the occurrence of peripheral neuropathy and indicators of heavy exposure to dioxin

By 31 June 1977, 22 cases with symptoms
or clinical and electrophysiological signs
of peripheral neuropathy and 20 cases with
electrophysiological abnormalities had been
found among the 470 subjects examined,
for a total PR of 8.9 % (95 % CL, 6.2—
11.6). Among the 308 subjects who under-
went a complete examination, 26 cases
with symptoms or clinical and electro-
physiological signs of peripheral neuro-
pathy and 16 cases with electrophysiologi-
cal abnormalities were found by 31 July
1978, for a total PR of 13.6 % (95 % CL,
9.5—17.8).

Table 2 reports the PR and the PRR of
peripheral neuropathy for subjects with
predisposing factors (group A) and for
subjects with chloracne or raised serum
hepatic enzyme levels (group B) versus
subjects without the symptoms of either

Table 1. Motor conduction velocity, distal latency, amplitude, and duration of muscle action poten-
tial (surface electrodes) in the 305 referents.

Nerve	Number of referents	Age (a)	Maximum conduction velocity (m/s)		Minimum conduction velocity (m/s)		Distal latency (ms)		Amplitude (mV)		Total duration (ms)
			Mean $\pm$ SD	Median	Mean $\pm$ SD	Median	Mean $\pm$ SD	Median	Range	Range	
Ulnaris (antia-	172	6—29	60.2 $\pm$ 5.3	60	55.5 $\pm$ 6.1	55	2.5 $\pm$ 0.8	2.5	4—15	11—13	
radialis)	118	30—59	58.2 $\pm$ 4.4	58	54.3 $\pm$ 4.3	55	2.5 $\pm$ 0.5	2.5	4—15	11—15	
	15	$\geq$ 60	57.8 $\pm$ 5.0	57	53.4 $\pm$ 4.4	52	2.6 $\pm$ 0.5	2.6	4—15	11—15	
Peroneal	166	6—29	52.7 $\pm$ 5.0	52	48.9 $\pm$ 4.9	49	3.0 $\pm$ 0.5	3.1	2—11	10—17	
capitulum	115	30—59	50.9 $\pm$ 4.1	51	47.9 $\pm$ 3.7	47	3.3 $\pm$ 0.8	3.2	2—11	10—17	
(super-	12	$\geq$ 60	49.8 $\pm$ 3.7	50	47.3 $\pm$ 4.1	47	3.9 $\pm$ 0.6	3.4	2—11	10—17	
ficie)											

Table 2. Prevalence rate (PR) and prevalence rate ratios (PRRs) for peripheral neuropathy in sub-
groups of Seveso residents in 1978.

	A With predisposing factors	B With indicators of exposure	C Without A or B
Actual cases of peripheral neuropathy	17	12	13
Total number of subjects	85	55	168
PR (%)	20	22	7.7
PRR ^a	2.6	2.8	1
95 % confidence limits	1.2—5.6	1.2—6.5	---

^a Evaluated versus group C.

Table 3 shows the PRR of peripheral neuropathy for subjects with increased γ -GT, GOT, and GPT activities and for subjects with chloracne versus subjects without any of these manifestations. The PRR was significantly different from one for the subjects with abnormal hepatic enzyme levels and was greater than one for the subjects with chloracne, even though the lower 95 % CL was equal to one.

Clinical symptoms and signs (paresthesia, hyposthenia, weakness, hypo- or areflexia

Discussion

It is evident from the preceding results that a very high prevalence of peripheral

	Abnormal hepatic enzyme levels ^a		Chloracne ^b	
	With	Without	With	Without
Actual cases of peripheral neuropathy	6	19	6	19
Total subjects	18	205	24	199
Prevalence rate ratio	3.6		2.6	
95 % confidence limits	1.3—10.2		1.0—7.2	

Table 4. Relationship between clinical symptoms and signs and peripheral neuropathy in subgroups of Seveso residents.^a

Subjects	A With predisposing factors		B With indicators of exposure		C Without A or B	
	Symptoms and signs	No symptoms or signs	Symptoms and signs	No symptoms or signs	Symptoms and signs	No symptoms or signs
Neuropathic	11 (65) [30]	6 (35) [12]	6 (50) [50]	6 (50) [14]	9 (69) [60]	4 (31) [5]
Normal	26 (38) [70]	42 (62) [85]	6 (14) [50]	37 (55) [86]	6 (4) [40]	149 (50) [187]
	$\chi^2 = 3.85$	$p < 0.05$	$\chi^2 = 7.15$	$p < 0.01$	$\chi^2 = 63.01$	$p < 0.001$

^b There was a significantly higher prevalence of neuropathy for subjects with symptoms and signs versus subjects with no symptoms and signs in each subgroup.

It must be stressed that the neuropathy was not severe. The clinical picture was considered the most similar to that seen with which to study the effects of dioxin-T on peripheral nerves.

At the follow-up nonattendants was analysis was carried out to determine whether the group examined in 1978 was as regards the disease of the presence of signs which could indicate affected subjects to be because of the predisposing towards. The frequency of pathology-related factors was higher among those subjects in respect to those not examined. More subjects with normal serum hepatic enzymes were reexamined in the hypothesis that they were positively susceptible to the disease. Therefore confirmed, the existing neuropathy subjects exposed to

were
peripheral
of the
subjects
ix with
with
hout
eral
up with
dioxin.
for six
acne or
els, the
the last
chlor-

g results
eripheral

oracne or

racne *

Without

19

199

7.2

subgroups

A or B

to symptoms
or signs

4 (31)

[3]

149 (96)

[97]

p < 0.001

signs of the
indicate the
is and signs
ms and signs

neuropathy was observed among subjects who showed indicators of heavy exposure to dioxin (12 out of 55, 22 %). This prevalence was almost three times greater than that of subjects who did not show these indicators. The high frequency of neuropathy found for subjects with chloracne is interesting because chloracne is a good indicator of exposure to dioxin. Moreover, during the monitoring, chloracne was found only in the young age group (0 to 20 a) (8), for which the frequency of neuropathic signs was very low. In fact, the PRR of neuropathy for the subjects under 20 a of age with chloracne versus those without it reached 4.7 (95 % CL, 1.2-18.1), a value greater than the one (2.6) observed in the chloracne subjects of all ages. In contrast, since abnormal hepatic enzyme levels were more common in the adult age group, their association with peripheral neuropathy may not be so unusual.

It must be stressed that no acute polyneuropathy was found during the study, and the polyneuropathic cases found were not severe. The electrophysiological diagnosis of peripheral neuropathy was considered the most appropriate procedure with which to study the possible chronic effects of dioxin-TCDD exposure on the peripheral nerves of a general population.

At the follow-up examination the rate of nonattendants was high, about 56 %. An analysis was carried out to determine whether the group of 277 subjects reexamined in 1978 was positively self-selected as regards the disease under study because of the presence of symptoms and clinical signs which could have encouraged the affected subjects to come to the clinic or because of the presence of factors predisposing towards peripheral neuropathy. The frequency of these possible neuropathy-related factors was not significantly higher among those reexamined with respect to those not reexamined. Moreover, more subjects with chloracne and abnormal serum hepatic enzyme levels were not reexamined in the second screening. The hypothesis that subjects who returned were positively self-selected with respect to the disease under study was not therefore confirmed by this analysis. The consistent neuropathy and chloracne among subjects exposed to dioxin may represent

an important qualitative finding for the identification of subjects at risk.

A neurological follow-up of this population is in progress. The principal objectives are to increase the examination attendance to at least 95 % and to improve the information on the "exposure indicators."

Acknowledgment

This work was supported by a grant from Regione Lombardia.

References

1. Bauer H, Schulz KH, Spiegelberg U. Berufliche Vergiftungen bei der Herstellung von Chlorophenol-Verbindungen. Arch Gewerbepath Gewerbehyg 10 (1961) 530-555.
2. Bergmeyer HU, Büttner H, Hillmann G, Kreutz FH, Lang H, Laue D, Riek W, Schmidt E, Schmidt FW, Stamm D, Szasz G. Recommendations of the German Society for Clinical Chemistry: Standardisation of methods for the estimation of enzyme activity in biological fluids. Z Klin Chem Klin Biochem 8 (1970) 650-660.
3. Bleiberg J, Wallen M, Brodtkin R, Applebaum H. Industrially acquired porphyria. Arch dermatol 99 (1964) 793-797.
4. Bollettino Ufficiale della Regione Lombardia. Supplemento straordinario al N.23: Provvedimenti per Seveso. Milano 14 Luglio 1977, pp 3052-3054.
5. Cosi V, Pinelli P, Torre E. Range of conduction velocity of motor nerve fibres in a given muscle: Method of desynchronisation and method of delay of the partial antidromic block. Electroencephalogr clin neurophysiol 21 (1966) 515.
6. Firestone D. The TCDD problem: A review. In: Rameil C, ed. Chlorinated phenoxy acids and their dioxin. Royal Swedish Academy of Sciences, Stockholm 1978, pp 39-52. (Ecot bull no NFR 27)
7. Goldmann PL. Schwerste akute Chloracne, eine Massenintoxikation durch 2,3,6,7-tetrachlorodibenzodioxin. Hautarzt 24 (1973) 149-152.
8. IARC Working Group on the Evaluation of the Carcinogenic Risk of Chemicals to Man. Some fumigants, the herbicides 2,4-D and 2,4,5-T, chlorinated dibenzodioxins and miscellaneous industrial chemicals. International Agency for Research on Cancer, Lyon 1977, pp 1-102. (IARC monogr 15).
9. International Labour Office. International standard classification of occupations. Geneva, 1968.
10. Jirásek L, Kalensky J, Kubec R, Pazdnerova J, Lukáš E. Acute chloroma, porphyria cutanea tarda and other manifestations of general intoxication during the manufac-

- Use of herbicides. *Cesk dermatol* 39 (1974) 145-157.
11. Jarešek L, Kalensky J, Kubec K, Paderova J, Lukáš E. Chloracne, porphyria cutanea tarda, and moderate intoxication by herbicides. *Hantarz* 27 (1976) 328-333.
 12. Paderova-Vejthlikova J, Němcová M, Páková J, Jirásek L, Lukáš E. The development and prognosis of chronic intoxication by tetrachlorodibenzo-p-dioxin in men. *Arch environ health* 36 (1981) 5-11.
 13. Poland AP, Smith D, Metter G, Possick P. A health survey of workers in a 2,4-D and 2,4,5-T plant, with special attention to chloracne, porphyria cutanea tarda, and psychologic parameters. *Arch environ health* 22 (1971) 316-327.
 14. Schulz KH. Zur klinik und etiologie der chloracne. *Arbeitsmed Sozialmed Arbeits-hyg* 3 (1968) 25-29.
 15. Stellman S, Stellman J. Health work ex among 535 Vietnam veterans exposed to toxic herbicides. *Am J* 112 (1980) 444 (abstract).
 16. Szasz G. Kinetic photometric method for serum γ -GT. *Clin chem* 15 (1969) 11-14.
 17. Thiess AM, Goldmann P. Follow-up über das Trichlorophenol-Dioxin-Geschehen in der BASF AG vom 1. November 1953. Presented at the 15th chem Kongress, Haifa, September 1977.
 18. Ton Tkhut Tung. Patologie humaine animal de la dioxine. *Rev med (Paris)* (1977) 653-657.
 19. Walker AE, Martin JV. Lipid profile in dioxin-exposed workers. *Lancet* 1 (1977) 446-447.
 20. Woolf B. On estimating the relation between blood group and disease. *Ann genet* 19 (1955) 251-253.

Received for publication: 27 May 1981

Key to

High level pressure is a hearing. Less the increasing caused by under the fact that regarded as ha Propeller no amounts to level in relation to the energy by frequency range of noise cause of noise frequent finding audiograms, within this field. The investigation was to from the noise rock drills and under water

- ¹ Norwegian Institute of Occupational Health, Oslo, Norway.
- ² Electronics Research Institute, Norwegian Institute of Technology, Trondheim.

Reprint requests to: Underwater, Norway.