

FILE NAME: Brakes (BRK)

DATE: 1968

DOC#: BRK030

DOCUMENT DESCRIPTION: Unpublished Report for Johns-Manville - The Pulmonary Response to Brake-Drum Dust

Q4a

1 7 4 2 1 7 4 2 1 7 4 5 2 1 7 4 4 2 1 7 4 3 2 1 7 4 2 2 1 7 4 1 2 1 DOCUMENT NUMBER															
DATE		INDEXED BY				EDITED BY				PAGE		OF		DOCUMENT NO.	
DOCUMENT TITLE												3071		LOCATION	
Gross, Paul:															
KEY CODE		KEY WORD OR CHARACTERISTIC								SCOPE NOTES					
1		The Pulmonary Response to Brake-Drum								Not to Circulate!					
72		Dust: A Preliminary Investigation.													
913		Ind. Hy. Foundation - Report for J-M.								Scientific Committee					
43		July 1968								only					
40		UNPUBLISHED)													
53															
80		Summary: "Brake-drum dust has evoked a pulmonary response in rats and hamsters which suggest that this dust is biologically 'inert.' Brake-drum dust is capable of producing ferruginous bodies in the lungs of hamsters that are indistinguishable from asbestos bodies."													

E83280 KEYDEX © 1962 - MCBEE FORM KE-1130

INDEX/TRACING CARD



3097
(412) 621-1100

INDUSTRIAL HYGIENE FOUNDATION OF AMERICA, INC.

4400 FIFTH AVENUE

PITTSBURGH, PA. 15213

CANADIAN ASBESTOS INFORMATION CENTRE
1130 SHERBROOKE ST. W. - STE. 410
MONTREAL (QUEBEC), CANADA H3A 2M8

Report on

THE PULMONARY RESPONSE

TO

BRAKE-DRUM DUST:


A PRELIMINARY INVESTIGATION

for

JOHNS-MANVILLE CORPORATION

by

Paul Gross, M.D.
Director of Research Laboratory
Industrial Hygiene Foundation


Robert T. P. deTreville, M.D.
President

1968

Three of the hamsters were killed 7 1/2 months after the injection and lung washings were obtained from these animals using a modified LaBelle and Brieger technique. The sediments from these washings were smeared upon slides which were allowed to dry and were examined microscopically unstained.

All remaining hamsters and rats were killed 12 months after the intratracheal dosage.

All lungs were distended with formalin under a pressure of 10 to 12 cm of water. Paraffin sections were cut and stained with hematoxylin and eosin. Pertinent fields on these slides were photographed after which the slides were decolorized and impregnated with silver using Gordon and Sweets technique. The same fields were then rephotographed. The coverslips were then removed and the slides subjected to microincineration for one hour at 600° C. After treatment with concentrated hydrochloric acid and washing with water, the same fields were photographed for a third time, but now under dark-field illumination. By printing through a composite negative, made by superimposing the dark-field negative upon either light-field negative, a picture is obtained that relates the mineral dust to the tissues and cells.

Longer-term Reaction

A. Seven and one-half months (hamsters)

The lungs of hamsters killed at this time are nearly normal except for widely scattered foci in which a few alveoli contain clusters of macrophages and the lining cells of their septa are excessively numerous and swollen. An occasional giant cell is also found in the air spaces.

Smears of the sediment from the lung washings show scattered asbestos bodies, many of which are segmented (Fig. 3).

B. Twelve months (rats and hamsters)

The only significant lesions found in the lungs of these animals are accumulations of dust-filled macrophages within widely scattered clusters of alveoli, associated with slightly increased arborescence of alveolar argyrophilic fibers (Figs. 4-6). In some instances, the affected alveoli can be identified as ones that were evaginating off alveolar ducts or respiratory bronchioles. No collagenization of alveolar walls or obliteration of alveolar lumens is demonstrable. Also noteworthy is the fact that in rats in spite of a six-fold greater dosage, the amount of alveolar involvement at one year is not greater than that seen at four days with only one dose of 3.5 mg of dust. A similar discrepancy is found in the lungs of hamsters, which also show a degree of pulmonary involvement at one year with a total dosage of 10.5 mg not greater than that of hamsters four days after a single injection of 3.5 mg.

dust could be ascribed to inadequate dosage. This may be true. However, from a practical point of view, it is not likely that a city dweller is apt to inhale more than an extremely small dose during a 24-hour exposure; and, as seen in these animals, there is no tendency for the lungs to accumulate, compact, and sequester the dust as is seen with certain other dusts.¹⁴ Consequently, removal of the inhaled brake-drum dust from the lungs may be expected to be good.

Nevertheless, it is advisable to repeat this study with a larger number of animals, using also larger dosages and allowing the animals to live out their lives.

Conclusions

Brake-drum dust has evoked a pulmonary response in rats and hamsters which suggests that this dust is biologically "inert."

Brake-drum dust is capable of producing ferruginous bodies in the lungs of hamsters that are indistinguishable from asbestos bodies.

References (continued)

9. Gross, P.; deTreville, R. T. P.; Cralley, L. J.; Davis, J. M. G.:
Pulmonary Ferruginous Bodies, Development in Response to Filamentous
Dusts and a Method of Isolation and Concentration, Arch. Path. 85:539-546,
(May), 1968.
10. Gross, P., et al: The Effect of a Synthetic Ceramic Fiber Dust upon the
Lungs of Rats, Arch. Indust. Health 13:161-166, (February), 1956.
11. Gardner, L. U.: Studies on the Relation of Mineral Dusts to Tuberculosis,
Part II: The Relatively Early Lesions in Experimental Pneumoconiosis
Produced by Carborundum Inhalation and Their Influence on Tuberculosis,
Am. Rev. Tuberc. 7:344, 1923.
12. Gross, P.; Westrick, M. L.; McNerney, J. M.: Experimental Tuberculo-
pneumoconiosis, Arch. Indust. Health 19:320-334, (March), 1959.
13. Gross, P. and Nau, C. A.: Lignite and the Derived Steam-Activated
Carbon: The Pulmonary Response to their Dusts, Arch. Environ.
Health 14:450-460, (March), 1967.
14. Brieger, H. and Gross, P.: On the Theory of Silicosis: III Stishovite,
Arch. Environ. Health 15:751-756, (December), 1967.



Fig. 3 Region of an alveolar duct one year after intratracheal injections of brake-drum dust. The tissue reaction at this time is not significantly different from that of Fig. 1 and still consists only of clustered macrophages. Hematoxylin and eosin. x 150



Fig. 4 The same field as in Fig. 3 but after silver impregnation, showing minimal arborescence of mural argyrophilic fibers. Gordon and Sweets. x 150




Fig. 5 Composite picture of the same field as in Fig. 3 showing that the macrophages still contain abundant mineral dust. x 150



Fig. 6 Smear of lung washings from a hamster injected intratracheally with brake-drum dust seven months previously, showing a typical segmented asbestos body. Eosin only. x1150