

PLAINTIFF'S  
EXHIBIT

USG 274

ARCHIVES OF

# ENVIRONMENTAL HEALTH

PREVENTIVE MEDICINE—OCCUPATIONAL MEDICINE—  
AEROSPACE MEDICINE

---

VOLUME 14

JANUARY THROUGH JUNE, 1967

---

American Medical Association Publication

# Mesothelioma and Asbestos Exposure

Jan Lieben, MD, and Harry Pistawka, MD, Harrisburg, Pa

WITHIN the past five years several articles have called attention to the association between mesotheliomas of the pleura and peritoneum with asbestos exposure.<sup>1-3</sup>

Wagner et al<sup>4</sup> studied 33 cases of mesothelioma. All but one had probable exposure to crocidolite, the blue asbestos.

Asbestos has become a ubiquitous material in our civilization. Cauna et al<sup>5</sup> reported that asbestos bodies were found in the lung smears of 41% of 100 consecutive autopsies at the Presbyterian University Hospital in Pittsburgh. Similar findings were reported by Thompson<sup>6,7</sup> in 500 consecutive lung smears from Capetown, South Africa, and Miami, Fla. Asbestos bodies were found in 30% of the men and 20% of the women. Asbestos has been conclusively associated with increased incidence of lung cancer and other malignancies<sup>8,9</sup> and an increasing number of studies associate its presence with malignant mesothelioma of the pleura and peritoneum.

## Pathology

Mesotheliomas of the pleura and peritoneum are rare neoplasms. Controversy involves histological types, nomenclature, origin, and their very existence. Recently, mesotheliomas as an entity have become generally accepted. Their morphologic variability makes classification difficult and differentiation from other malignancies is often a problem.

Sacconi and Coblenz<sup>10</sup> give the incidence of pleural mesotheliomas as 1.1 cases per 1,000 autopsies. Campbell reported four cases of pleural mesotheliomas in 3,533 consecutive autopsies.<sup>11</sup>

Mesothelioma occurs primarily in adults and is twice as common in males as in females.<sup>10,12</sup>

Pleural mesotheliomas occur equally on the right and left though Weissman feels that pleural mesothelioma occurs on the right more frequently than on the left side.<sup>12</sup>

The criteria for diagnosis of pleural mesothelioma are presence of a firm pleural mass encasing the lung, histological structure compatible with mesothelioma, and absence of a demonstrable primary malignant neoplasm elsewhere. The histologic architecture may be mesenchymal, "epithelial," or mixed.

A complete autopsy is necessary for unequivocal diagnosis, but for the purpose of this study, criteria were made less rigid in that unautopsied cases were accepted if a definite diagnosis of pleural mesothelioma had been made by biopsy, and malignancy elsewhere had been excluded clinically; chest x-ray examination, bronchoscopy, and cytologic studies of sputum or bronchial washing assisted in excluding the presence of bronchial carcinoma.

Primary tumors of the peritoneum are among the rarest of tumors. Generally called mesotheliomas, they form large spreading masses over the peritoneum and histologically appear the same as pleural mesotheliomas. As with pleural mesotheliomas, exclusion of a primary tumor is necessary.

This study is an attempt to learn the extent of the relationship of asbestos exposure with mesothelioma diagnosed in southeastern Pennsylvania.

Submitted for publication June 6, 1968; accepted Sept 13.

From the Division of Occupational Health, Department of Health, Harrisburg, Pa.

Read before the Ninth International Cancer Congress, Tokyo, October 1966.

Reprint requests to PO Box 90, Harrisburg, Pa 17120 (Dr. Lieben).

## Method

One hundred and sixty-two hospitals serving a population of 6½ million people were requested to report all mesotheliomas diagnosed between 1958 and 1963. Replies were received from 152 hospitals. Hospital records of 62 reported cases were reviewed and only mesotheliomas confirmed by operation, biopsy, or autopsy were included in the study. Forty-two cases satisfied our criteria. There were 34 mesotheliomas of the pleura and eight of the peritoneum. Among the pleural mesotheliomas there was a definite preponderance for occurrence on the right side of the chest. Of 34 pleural mesotheliomas only seven occurred on the left side, 25 on the right side, and two were not stated.

Sections of as many of the tumors and lungs as possible were borrowed and reviewed by a single consultant pathologist familiar with mesotheliomas. In the 15 instances in which lung sections were available, they were examined for the presence of asbestos bodies. All slides were reviewed without the benefit of the complete history and autopsy protocol available to the pathologists who made the original diagnosis.

Results of the 33 cases reviewed (Table) by our pathologist agreed with those of the outside pathologists in 17 cases; he rejected seven cases as unacceptable histologically, had serious doubts in nine cases (of these one slide was read as metastasis and one slide as anaplastic malignancy). Of the 15 cases in which lung tissue was available, asbestos bodies were identified in seven.

The discrepancies between the original pathological diagnosis and that of our pathologist are reported to illustrate the difficulties and disagreements which are so frequent in the diagnosis of this tumor. For inclusion in this study the diagnoses of the originally reporting pathologists have been accepted and cases reported by them were investigated.

Efforts were made to obtain a complete employment and residence history for each patient. Inquiry was also made into the employment history of family contacts. Data were obtained from the individual if alive, from family members, or employers.

Of the 42 patients, ten had definite occupational exposure to asbestos during lifetime, three were family contacts of asbestos workers, eight either lived in the immediate neighborhood of asbestos plants or had been employed next to an asbestos plant. Ten patients had a questionable asbestos exposure and, in the other 11 patients, either no asbestos exposure could be elicited on direct questioning or no survivor could be located.

Table 1.—Readings of Consultant Pathologist

| Case No. | Mesothelioma                                        | Asbestos Bodies |
|----------|-----------------------------------------------------|-----------------|
| 1-C      | Yes                                                 | No lung tissue  |
| 3-C      | Not acceptable                                      | Yes             |
| 4-C      | Probably not                                        | No lung tissue  |
| 5-C      | Yes                                                 | Yes             |
| 6-C      | Probably not                                        | Yes             |
| 9-C      | Probably not                                        | No lung tissue  |
| 10-C     | Yes                                                 | Yes             |
| 2-N      | Yes                                                 | No lung tissue  |
| 3-N      | Probably not                                        | No lung tissue  |
| 4-N      | Not acceptable                                      | No lung tissue  |
| 6-N      | Yes                                                 | No lung tissue  |
| 1-F      | Yes                                                 | No lung tissue  |
| 2-F      | Yes                                                 | No lung tissue  |
| 3-F      | Yes                                                 | No lung tissue  |
| 1-Q      | Probably not                                        | No lung tissue  |
| 2-Q      | Yes                                                 | Yes             |
| 4-Q      | Yes                                                 | No              |
| 5-Q      | Probably yes                                        | No              |
| 6-Q      | Not acceptable                                      | No              |
| 7-Q      | Yes                                                 | No lung tissue  |
| 8-Q      | Yes                                                 | No              |
| 9-Q      | Probably not                                        | No              |
| 10-Q     | May be Metast.                                      | No              |
| 1-X      | Yes                                                 | Yes             |
| 2-X      | Yes                                                 | No lung tissue  |
| 3-X      | Not acceptable                                      | No lung tissue  |
| 4-X      | Not acceptable                                      | No lung tissue  |
| 5-X      | Anaplastic malignancy                               | No              |
| 6-X      | Probably not                                        | No              |
| 7-X      | No epithelial cells<br>Mesothelioma must be proven. | No lung tissue  |
| 8-X      | Probably not                                        | No lung tissue  |
| 9-X      | Yes                                                 | No lung tissue  |
| 10-X     | Yes                                                 | Yes             |

**Exposure of Ten Mesothelioma Patients Who Worked With Asbestos.**—Patient 1-C worked 35 years in the textile department of a large asbestos plant.

Patients 2-C, 3-C, and 4-C worked more than 20 years in asbestos insulation plants and patient 5-C worked in an insulation plant for only one year.

Patient 6-C worked 15 years in a plant manufacturing acoustic tile and linoleum.

Patient 7-C was a boiler maker in a railroad yard where, for 25 years, he worked on insulation of steam engines.

Patient 8-C built bakery ovens in which large amounts of asbestos insulation were used.

Patient 9-C was a certified accountant who worked more than 30 years in the office of an asbestos textile plant.

Patient 10-C was a plasterer who came in contact with a large amount of asbestos in his work.

Of these ten men, patients 1-C, 8-C, and 10-C had mesotheliomas of the peritoneum; the other seven men had pleural mesotheliomas.

**Neighborhood Cases.**—Patient 1-N was a 55-year-old woman who never had any occupa-



(from 1925 to 1938, and later in a plastic fabrication plant which utilized fiber glass; he also had been a welder for 25 years. He thinks he has used asbestos but could not recall when and where.

Patient 10-Q, a 65-year-old man, was a core-maker all his life. He worked in two foundries in one of which asbestos cement is used in small quantities but, according to the plant manager, the core-maker did not have any contact with this material. The son stated that his father had used asbestos pipe insulation for pipes in the basement of their home several years prior to the onset of his illness and that he had done some sawing of this pipe insulation material.

All cases in this group had pleural mesotheliomas.

**Mesothelioma Patients Without Asbestos Exposure.**—In the other 11 patients (1-X to 11-X) no occupational, familial or neighborhood asbestos exposure evidence could be elicited. Ten of these patients had mesothelioma of the pleura and one peritoneal mesothelioma.


### Comment

This study presents a good deal of circumstantial evidence and many unanswered questions.

Asbestos is a ubiquitous material used much more frequently than is generally realized. Some uses, such as in the brewery, were not previously known to the writers. There are undoubtedly many others.

If we assume that asbestos may be the causative or trigger agent for malignant mesothelioma in some of the cases described above, we must admit that the minimal dose-effect relationship and duration of a latent period are unknown. The occupational exposure of the insulation workers or asbestos textile workers were certainly many thousand times higher than those of the neighborhood cases or family contacts. Similar findings have recently been reported from England.<sup>1</sup>

The attack rate for mesotheliomas is ex-



Occupational, neighborhood, and family contact patients with mesothelioma associated with one insulation plant.

tremely low. Not all mesotheliomas were reported to us by the hospitals, but, even if we double the incidence, only a minute fraction of the exposed population was affected. There are precedents for this with beryllium disease which acts in a somewhat similar pattern; occupational diseases, familial contacts, neighborhood cases plus a very low attack rate, and a long latent period between exposure and onset of disease.

Another question which arises is why pleural mesotheliomas are so much more frequent than peritoneal mesotheliomas.

The most striking finding of this study was that of the 42 mesotheliomas which came from a geographical area of approximately 30,000 sq mi, six were clustered in and around an insulation plant (Figure). These included two insulation manufacturing workers (Cases 2-0 and 3-0), three neighborhood cases (Cases 2-N, 3-N, and 4-N) and one family contact (Case 2-F)—the daughter of an insulating worker.

#### Summary

Forty-two cases of mesotheliomas reported from 152 hospitals over a five-year period, were studied with regard to exposure to asbestos. Survivors or employers, or both, were questioned regarding the possibility of asbestos exposure. Ten patients actually worked in asbestos plants; eight lived or worked close to an asbestos industry; three patients were family members of asbestos workers. In ten patients a history of assumed exposure to asbestos was obtained after prolonged questioning. In 11 other pa-

tients no history of asbestos exposure could be obtained.

#### References

1. Newhouse, M.L., and Thompson, H.: Mesothelioma of the Pleura and Peritoneum Following Exposure to Asbestos in the London Area. *Brit J Industr Med* 22:261-269, 1965.
2. Hourihane, P.O.: The Pathology of Mesotheliomata and an Analysis of Their Association With Asbestos Exposure. *Thorax* 19:268-277, 1964.
3. Selikoff, I.J.; Churgh, J.; and Hammond, C.: Exposure to Asbestos and Mesothelioma. *New Eng J Med* 272:560-563 (March) 1965.
4. Wagner, J.C.; Sleggs, C.A.; and Marchand, P.: Diffuse Pleural Mesothelioma and Asbestos Exposure in N.W. Cape Province. *Brit J Industr Med* 17:260-271, 1960.
5. Cauna, D.; Totten, R.; and Gross, P.: Asbestos Bodies in Human Lungs at Autopsies. *JAMA* 192:371-373 (May 3) 1963.
6. Thompson, J.G.; Kaschula, R.O.; and MacDonald, R.R.: Asbestos as a Modern Urban Hazard. *S Afr Med J* 37:77-81 (Jan) 1963.
7. Thompson, J.G.: Asbestos and Urban Dweller, read before the Conference on Biological Effects of Asbestos, New York, Oct 19, 1964.
8. Doll, R.: Mortality From Lung Cancer in Asbestos Workers. *Brit J Industr Med* 12:31, 1955.
9. Lieben, J.: Asbestos and Malignancy. *Arch Environ Health* 13:619-621 (Nov) 1966.
10. Sacconi, A., and Coblentz, A.: Endothelioma of the Pleura. With Report of Two Cases. *Amer J Clin Path* 13:186-207, 1943.
11. Campbell, W.: Pleural Mesothelioma. *Amer J Path* 26:473-487, 1950.
12. Weissman, H.: Primary Endothelia of the Pleura. *Dis Chest* 12:562-570, 1946.
13. Lyman, T. (ed.): *Metals Handbook: 1. Properties and Selection of Metals*, ed 8, American Society of Metals, 1961, p 5.
14. *Modern Chemical Processes*, New York: Reinhold Publishing Co., 1952, vol 2, p 167.

#### DISCOVERY OF QUININE

Many of the drug sources known in ancient times were apparently discovered by accident. For example, according to an old South American Indian legend, the curative power of cinchona bark (containing quinine) for malaria was discovered by a sick, feverish Indian lost in the jungle. He quenched his thirst by drinking from a pool of water. From the bitter taste of the water, he recognized that it was tainted with the poison from the quina-quina tree. Although he shared this fictitious belief, he drank deeply so as to bring about a quick death. To his great surprise, he recovered completely. When he returned to his village, he told the story of his cure and thereafter cinchona bark was used as a medicine for the fever prevalent in that region. The Incas later taught the Spanish conquistadors that malaria could be cured with cinchona extract.—Rossman, R.E.: The History and Significance of Serendipity in Medical Discovery. *Trans Coll Physicians Phila* 33:104-120 (Oct) 1965.