GRANTSBURG, WIS.
SENTINEL
W. 2,380

JUN 18 1975

DNR warns against eating fish from certain contaminated water

DNR upon advice of the Wisconsin Division of Health, suggests that people limit their consumption of large trout and salmon from Lake Michigan, and other fish from the Upper Mississippi river, and the Fox river near Portage.

Fish in these waters contain more than the U. S. Food and Drug Administration's acceptable level of PCB's (Polychlorinated Biphenyls).

The FDA has set the public health action level for market fish at 5 parts per million (ppm) of PCB's. Recognizing that some larger fish in these waters contain more than 5 ppm, the Division of Health recommends that, "sports fishermen be advised to refrain from eating more than one meal per week of such fish and that pregnant women and children be cautioned to limit their intake to one average size serving of such fish per week."

Fish from the following waters have been found to exceed these levels:

1. Lake Michigan and Green Bay trout and salmon, including lake trout, coho salmon, chinook salmon, and brown trout, over 24 inches in length.

2. Many fish in the Mississippi river between Prescott and Lower Lake Pepin.

3. Many fish in the Fox river from Portage north to the Columbia county line.

The Wisconsin Department of Agriculture has announced that it will take action and seize any fish offered for commercial sale containing more than 5 ppm of PCB's. Lake trout over 24 inches long caught by commercial fishermen in Lake Michigan incidental to their netting operations for other species will no longer be sold for human consumption. The DNR will provide the Wisconsin Department of Agriculture with samples of these fish for testing and regulatory action.

PCB's have been used in coolant-insulation fluids in some transformers and capacitors, heat transfer agents, machine tool cutting oils, hydraulic fluids, electrical wiring insulation, and epoxy paints. They also occur in sewage treatment plant and some pulp and paper mill effluents. They may enter the air through vaporization or incineration, or enter the water via fallout or runoff.

Like DDT, PCB's are fat soluble with a low solubility in water and therefore are concentrated in the fat of animals, and particularly fish. They also are very stable and tend to persist in the environment.

MONS 031183