EDWARDSVILLE, ILL.
INTELLIGENCER
D. 6,000

APR 29 1977

MONS 001052

Law bans production, distribution by 1979

PCB contamination in air, water, soil—and people

Chicago (AP)

Poly-chlorinated biphenyl (PCB), a manmade chemical used generally in electrical equipment, is found today throughout the United States in the air, water, soil—and in people.

Last year, after tests determined it was a health hazard, Congress passed a law banning all production and distribution of PCBs by 1979.

Approximately 1.4 billion pounds of PCBs have been produced since Monsanto Industrial Chemical Co. of St. Louis began manufacturing the chemical in 1929. At present, government figures estimate that about 750 million pounds are in use in electrical equipment and some 450 million pounds sight in landfills, the air, water, soil or sediments.

Samples of human fat tissues have shown an average of one to three parts per million (ppm) of PCBs in the human population throughout the nation. And a recent study showed that many fish in the Great Lakes contain more than 5 (ppm.) which is the tolerance level for human consumption set by the federal Food and Drug Administration. The FDA is considering lowering the level to 2 ppm.

Karl Bremer, toxic substances coordinator for the U.S. Environmental Protection Agency, said the chemical is "very persistent and slow to break down, if it breaks down at all."

The first indication that PCBs constituted a health hazard occurred when workers manufacturing the substance developed small bumps, sores and lesions on the skin, Bremer said.

In 1968, about 1,000 persons in Japan consumed rice oil that accidentally had been contaminated with about 2,000 parts per million of PCBs. Reactions ranged from sores, acne lesions, blotching and skin pigmentation problems to miscarriages and stillbirths. Bremer said many of the victims still suffer from the poisoning.

Carcinogen

Furthermore, PCBs are a suspected human carcinogen and have produced cancer in rates in studies at the Atlanta Center for Disease Control.

Bremer said that in highly industrialized areas, higher levels of PCBs generally are found in fish, animals and the environment.

He noted, though, that a national survey of PCB content in mothers' milk, conducted by the Colorado State Medical School, showed "it didn't matter much which geographical area the mother came from."

The chemical itself, depending on how much chlorine is present, can be a white, powdery solid or a liquid ranging from a clear, oily substance to a yellow, sticky material.

Its primary use is in electrical equipment, including transformers and capacitors, fluorescent lighting, mercury and sodium vapor street lights, central air-conditioning units and high-efficiency refrigerators.

It also has been used extensively in carbonless copy paper, in waterproofing paint, in hydrolic oils, in resel oils and in plastics.

But in nearly every instance, acceptable substitutes for the uses of PCBs have been found, Bremer said.

A typical problem resulting from use of PCBs is illustrated by what happened when about 30,000 silos in Illinois, Indiana, Michigan and Kentucky were painted with a waterproof paint containing PCBs, Bremer said.

"The PCBs were transferred from the paint to the grain stored in the silos. That was fed to cows and it came out in the milk. Repainting doesn't get rid of the PCBs. We don't know what to do with those silos. The concrete is contaminated and we're having a real disposal problem."

Disposal of PCBs may become even more difficult. In Southern Illinois, citizens of tiny Wilsonville are enraged that the chemical is being hauled from the St. Louis area and deposited in a landfill near their community. They have threatened to blockade the landfill and are fighting the issue in the courts.

The federal ERA is required this month to issue final regulations on disposal and storage requirements for PCBs, Bremer said. Ultimately, PCBs should be completely broken down, but it takes special incinerators that heat to more than 2,000 degrees Fahrenheit to do that.

"The chemical incinerators presently in operation are overloaded," Bremer said. "They're also burning pesticides and other hazardous chemicals. It's a real mess."

PONTIAC, ILL.
LEADER
D. 5,500

APR 29 1977

Seek to halt use of hazardous PCBs by 1979

By CHERYL DEBES
Associated Press Writer

CHICAGO (AP) — Polychlorinated biphenyl (PCB), a manmade chemical used generally in electrical equipment, is found today throughout the United States in the air, water, soil—and in people.

Last year, after tests determined it was a health hazard, Congress passed a law banning all production and distribution of PCBs by 1978.

Approximately 1.4 billion pounds of PCBs have been produced since Monsanto Industrial Chemical Co. of St. Louis began manufacturing the chemical in 1930. At present, government figures estimate that about 750 million pounds are in use in electrical equipment and some 450 million pounds exist in landfills, the air, water, soil or sediments.

Samples of human fat tissues have shown an average of one to three parts per million (ppm) of PCBs in the human population throughout the nation. And a recent study showed that many fish in the Great Lakes contain more than 5 ppm, which is the tolerance level for human consumption set by the federal Food and Drug Administration. The FDA is considering lowering the level to 2 ppm.

Kari Bremer, toxic substances coordinator for the U.S. Environmental Protection Agency, said the chemical is "very persistent and slow to break down, if it breaks down at all."

The first indication that PCBs constituted a health hazard occurred when workers manufacturing the substance developed small bumps, sores and lesions on the skin, Bremer said.

In 1968, about 1,000 persons in Japan consumed rice oil that accidentally had been contaminated with about 2,000 parts per million of PCBs. Reactions ranged from sores, acne lesions, blotching and skin pigmentation problems to miscarriages and stillbirths. Bremer said many of the victims still suffer from the poisoning.

Furthermore, PCBs are a suspected human carcinogen and have produced cancer in rats in studies at the Atlanta Center for Disease Control.

Bremer said that in highly industrialized areas, higher levels of PCBs generally are found in fish, animals and the environment.

He noted, though, that a national survey of PCB content in mothers' milk, conducted by the Colorado State Medical School, showed "it didn't matter much which geographical area the mother came from."

The chemical itself, depending on how much chlorine is present, can be a white, powdery solid or a liquid ranging from a clear, oily substance to a yellow, sticky material.

Its primary use is in electrical equipment, including transformers and capacitors, flourescent lighting, mercury and sodium vapor street lights, central air-conditioning units and high-efficiency refrigerators.

It also has been used extensively in carbonless copy paper, in waterproofing paint, in hydrolic oils, in road oils and in plastics.

But in nearly every instance, acceptable substitutes for the uses of PCBs have been found, Bremer said.

A typical problem resulting from use of PCBs is illustrated by what happened when about 30,000 silos in Illinois, Indiana, Michigan and Kentucky were painted with a waterproof paint containing PCBs, Bremer said.

"The PCBs were transferred from the paint to the grain stored in the silos. That was fed to cows and it came out in the milk. Repainting doesn't get rid of the PCBs. We don't know what to do with those silos. The concrete is contaminated and we're having a real disposal problem."

Disposal of PCBs may become even more difficult. In Southern Illinois, citizens of tiny Wilsonville are enraged that the chemical is being hauled from the St. Louis area and deposited in a landfill near their community. They have threatened to blockade the landfill and are fighting the issue in the courts.

The federal EPA is required this month to issue final regulations on disposal and storage requirements for PCBs, Bremer said. Ultimately, PCBs should be completely broken down, but it takes special incinerators that heat to more than 2,000 degrees Fahrenheit to do that.

"The chemical incinerators presently in operation are overloaded," Bremer said. "They're also burning pesticides and other hazardous chemicals. It's a real mess."

MONS 001053

...OK, OHIO
...ORDAIE
D. 21,000

JUL 23 1977

Threat of toxic PCBs continues

By Newsday

NEW YORK — For a year and a half, 16 female monkeys were fed food contaminated with the chemical PCB (polychlorinated biphenyl) in concentrations similar to those found in fish caught off Long Island and elsewhere in the United States.

When the monkeys mated, seven miscarried. Two could not conceive at all. Uterine growths were recorded. The monkeys developed facial sores and swollen eyelids and lost their hair. Their offspring all were undersized and contained detectable levels of the toxic chemical.

The infants were then transferred. They all developed the same symptoms as their mothers. Within four months, half the infants died. Two years later, even when taken off the breast milk, the other monkeys still exhibit learning and behavioral disabilities.

Last year, the Environmental Protection Agency began testing the breast milk of American women for PCB contamination. The chemical was found in the milk of 75 of the first 80 women tested all over the country. The average levels were about half the lowest concentrations found in the monkeys, and about an eighth of the highest concentrations.

That study, by Dr. J.R. Allen, a pathologist at the University of Wisconsin, is one of the principal reasons why the federal government is preparing to lower drastically its allowable limits of PCBs in fish. In its proposed standards, expected to be enacted this year, the Food and Drug Administration goes even further and cautions women of child-bearing age and children to eat as few PCB-laden fish as possible, even if the fish are just under the limit.

"PCBs are more toxic than we had originally thought," said Elizabeth Campbell, an agency chemist who wrote the new guidelines. "The more information we collect about them, teh worse they seem to be."

Few negative effects from regular exposure to long-term, low-level doses of PCBs have been observed in humans, although consumers have been eating contaminated fish for years. The fatty tissues of workers exposed to PCBs have been found to contain levels dozens of times the proposed limits, but the main detected symptoms have been skin sores that cleared up after the exposure ceased.

Animal studies, however, show reproductive failure, cancer, decreased resistance to disease and other ailments at does not far above what humans could get over a period of many years, since PCBs accumulate in the body, once ingested, and do not break down. Currently, 90 per cent of the people in this country have detectable levels of PCBs in their fatty tissues and the fatty profiles in the breast milk of the average woman has levels that approach or exceed the new safe limits for edible fish.

Recently, studies have found PCBs in the embryonic fluid surrounding developing human fetuses and in the placenta and other tissues, in amounts many times the safe levels. In fact, children are being born with PCBs in their tissues.

Moreover, an official of the National Cancer Institute calculated for Newsday that a person eating the low levels of PCBs present in Long Island fish increases the risk of liver cancer by 6 per cent.

The most widely known and accepted animal studies have found that:

—Complete reproductive failure occurred when minks ate food contaminated with five parts per million of PCBs, which is the current federal safe limit. After 136 days at 3.5 parts per million, all the test minks died. That level is similar to the average in Long Island bass and bluefish. Minks, of course, are smaller than humans, so eating more symptoms at lower doses.

—Chickens, rabbits and birds fed doses of PCBs from 10 to 100 parts per million showed various symptoms including loss of feathers, skin lesions tremors and diminished growth.

Eighty per cent of the exposed rats tested showed cancerous or pre-cancerous growths, compared to less than 1 per cent of an unexposed group. The doses were about 250 times higher than people would get if they ate contaminated fish. An interagency study of PCBs last year said that although such doses (100 parts per million) are high when compared to daily consumption by humans, they are not very high for PCB workers or for people who eat a lot of fish contaminated with the chemical.

All detectable levels of PCBs in water, no matter how small, were found to collect in the tissues of marine organisms. As a result of a part per billion in water, fish offspring became lethargic, stopped feeding, developed lesion and started dying 24 hours after hatching. At one part per billion of PCBs in water, half the exposed shrimp died in 15 days. This concentration is similar to that in the Hudson River from the General Electric Co.'s discharges at plants near Albany. Reproduction rates in all species were reduced at varying concentrations. Resistance to disease was lowered: many species died of viruses that did not infect an unexposed group. Even exposed organisms placed in clean water transferred the chemical to their offspring.

The female rhesus monkeys tested by Allen developed telltale PCB acne after about one month on a continuous diet of the contaminated food.

Of the female monkeys fed food with five parts per million PCBs, reproduction dropped by more than 80 per cent. Of those fed 2.5 parts per million, reproduction dropped by about 50 per cent. Male monkeys, on the other hand, had few symptoms at those doses, Allen said.

"These studies indicate that women who plan to have children had best discontinue eating (contaminated) fish," Allen said. He said that a woman who simply stops eating fish once she gets pregnant may already have consumed and collected PCBs in her tissues. Since the monkeys weighed about 13.5 pounds, a lentil what the average woman weighs. It would take a woman about 10 times as long to accumulate the same doses, he said.

Even the proposed federal two-parts-per million limit on edible fish is not meant for daily consumption, especially by women and children, said the FDA's Campbell. "We figure that four fish meals per month at those levels should be the maximum, especially for women and children," she said.

The caution on the consumption of PCBs by children is based primarily on the fact that since children are smaller than adults, the PCB doses they receive when eating a contaminated fish are much larger per pound of body weight.

An average child nursing at mother's breast for a year would ingest what adults ingest in a decade," said Joseph Highland, a biochemist and PCB expert for the Environmental Defense Fund, a Washington-based organization that prodded the federal government two years ago into taking action to curtail consumption of PCBs.

In the absence of human analysis, the chemical can still be shown to be harmful in theory. PCBs are foreign to the body and are viewed by the system as pollutants. The liver tries to change the substance in some way so that it can be metabolized and excreted as waste. But PCBs are so stable that they cannot be broken down by the body, except slightly, in some instances. The liver works hard to metabolize the substance, and can be damaged in the process.

Laboratory tests indicate that PCBs, even if they do not produce negative effects themselves, can change the liver in a way so that the body is more open to cancer from other toxic substances.

Since the body cannot metabolize PCBs, they essentially cannot be excreted. They concentrate in the lipids, or fatty tissues, which are the body's storage area for most pollutants and excess food energy. The areas include the layer under the skin, the breasts and the abdominal area. They average about 20 per cent of the body's tissues for a person of average proportions.

Negative health effects of PCBs were initially reported in 1936, seven years after the material was first produced commercially. A U.S. Public Health Service report said workers had gotten skin sores from PCB exposures. The sores are believed now to be caused by concentrations of the chemical in the fatty layer under the skin. The ailment became known as "chloracne," or acne from chlorinated hydrocarbons.

Why, then, wasn't the material curtailed?

"The reports were evaluated as an example of misuse and poor handling," said William Papageorge, a biochemist and PCB expert for Monsanto Chemical Co. of St. Louis, the sole manufacturer of PCBs.

But in the mid-1960s the chemical started turning up in human hair, bird eggs and fish. Then, in February, 1968, about 1,200 Japanese consumers ate rice oil contaminated with PCBs. The chemical was used in a closed heating element and had accidentally leaked into cooking oil as it was being prepared in a factory in Yusho, Japan. By March, victims had skin eruptions, swelling of the eyelids and joints, impaired vision, loss of appetite, nausea, vomiting, overall fatigue, darkened skin, abnormal menstruation, impotence, jaundice, fever, hearing difficulties, headaches and spasms.

The offspring born to contaminated parents had darkened skin and PCB concentrations in their tissues. The skin symptoms persisted to a lesser extent after three years. Of the 13 contaminated women who became pregnant, there were two stillbirths and 12 of the 12 babies were smaller than national standards, according to scientific papers in a new book, "PCB Poisoning and Pollution."

Up to Sept. 13, 1972, 22 deaths had occurred among the 1,200 PCB victims. Of those, nine were from cancer, "suggesting a possible excess mortality (from cancer)," according to one of the scientific papers. The paper adds, however, that the "rigorous statistical analysis" has not been done.

Among the 22 dead were a 15-year-old boy and a 36-year-old truck driver. Autopsies discovered internal growths similar to chloracne and various irregularities in lymph nodes, glands or the liver. Neither of those two died from cancer.

The average concentrated of PCBs consumed by the Japanese victims over several weeks was 2,000 to 3,000 parts per million. That is 1,000 to 1,500 times the proposed federal limits for fish consumption.

To reach the minimum levels that caused adverse symptoms at Yusho, a person would have to eat about 100 pounds of contaminated fish. The doses are cumulative, however. Each meal consumed brings one closer to danger, scients say.

The average Long Islander eats about 50 pounds per year — about four times the national average, according to J. L. McHugh, a biologist at the Marine Sciences Research Center of the State University of New York at Stony Brook. A 10-year dose of contaminated fish would equal the minimum Japanese accumulation. Sports fishermen, however, usually consume more than the general population, perhaps as much as 150 pounds annually, according to Robert Rance head of Save Our Strippers, Long Island sport fishing a. sociation. If a fisherman and his family ate only contaminated fish, they could therefore build up enough for the minimum Yusho concentration in less than five years.

Officials are quick to point out that the figures are for the worst case. Not all the fish are contaminated, and there is no way of determining which are, without a laboratory analysis.

In the last five years, the more studies that have been done, the more places PCBs have been found. Recently, the chemical was found in fish 11,000 feet beneath the ocean. The government has been holding hearings in Washington this month on the cleaning possibilities and alternatives. Detailed epidemiological studies on humans are just being started, and in a much-disputed study, University of Pennsylvania researchers linked PCB to cancer in New Jersey workers.

The Environmental Defense Fund's Highland contended that unless drastic action is taken to limit exposures it will be just a matter of time until health problems start showing up in the general population.

But scientists do not yet agree on the best course of action. There is consensus on only two aspects. First there is no easy answer. Second as one federal expert put it, "the animal studies give us very good cause for concern."

MONS 001054