

Castleman File: American Petroleum Institute
w/c = with cover letter or memo If DATE = 0, undated
CD-ROM Document #: API 168 DATE 1964

- ☒ published article from medical journal
- ☐ published article from trade journal
- ☐ published advertisements
- ☐ newspaper article
- ☐ published government report
- ☐ government inspection results
- ☐ unpublished or internal report
- ☐ unpublished presentation from conference
- ☐ letter
- ☐ memorandum
- ☐ industry warning labels
- ☐ industry sales literature
- ☐ industry recommended practices
- ☐ meeting minutes (with attachments)
- ☐ membership list
- ☐ BC notes

TABLE OF CONTENTS

Executive Health

1. What Do We Mean By Stress Disease? 233

J. H. Sterner, M.D.

2. Are Executives Prone to Stress Disease? 238

A. J. Fleming, M.D.

3. Executive Health — How to Maintain It 243

R. F. Schneider, M.D.

- Evaluation of the Worker — Tools and Techniques for the Future 249

R. E. Eckardt, M.D., Ph.D.

- Un Decenio de Mortalidad en una Empresa Petrolera Venezolana 259

Dr. R. Risquez-Iribarren

- What to Say to an Alcoholic 275

- Nickel Carbonyl — Its Detection and Potential for Formation 282

R. S. Brief, F. S. Venable and R. S. Ajemian

- A New Stretcher 294

J. L. J. de Bary, M.D. and A. W. Gardner, M.D.

- Methods for the Prevention of Lead Poisoning in Industry 297

R. A. Kehoe, M.D.

- A Five Year Review of a Mercury Control Program 310

A. H. Diserens

- Checking on Health — Esso Petroleum Company, Ltd. 318

- Abstracts 325

- Newsnotes 328

ABSTRACTS *

The Clinical Significance of Uncomplicated Marked Left Axis Deviation in Men Without Known Diseases. R. S. Eliot; W. A. Millhon and J. Millhon. *American Journal of Cardiology*; 12; 767-771 (Dec. 1963).

One hundred and ninety-five men with marked left axis deviation (-30° to -90°) ranging in age from 23 to 61 years were compared with 100 men with normal axis deviation in the same age distribution at an initial examination and at follow up 3 to 22 months later. Men with other ECG abnormalities, clinical cardiovascular-renal disease or diabetes mellitus known to exist prior to the initial examination were excluded from the study.

113 or 58% of the 195 subjects with LAD had cardiovascular disease or diabetes mellitus on the initial or follow up examination compared to 3% with hypertension and 1% with diabetes in the control group. A more detailed analysis of the abnormal findings revealed:

Angina coronary insufficiency, myocardial infarction or a positive exercise tolerance test occurred or developed in 34% of the study group. None of these diagnoses occurred in those with NAD.

Hypertension alone occurred in 14% of the LAD group compared to 3% of the NAD group. 2% in the LAD group had both hypertension and angina.

Subclinical or latent diabetes mellitus was found in 7% of the study group compared to 1% of the controls.

Cerebral hemorrhage or infarction developed in 2% of the study group. Neither occurred in the controls.

Six patients died of coronary or cerebrovascular disease in the study group during the brief period of follow up.

Body habitus or blood cholesterol levels were not significantly different in the two groups.

The group with LAD had a much stronger family history of cardiovascular disease than did the NAD group.

The authors conclude that the ECG finding of uncomplicated marked LAD should be suspect because of the above associations.

Reprinted with permission from the *Journal of Occupational Medicine*, July 1964.

* These abstracts have been prepared by the Medical Research Division of Esso Research and Engineering Company.

Permanent Threshold Shift
Aging. R. Gallo and J.
June 1964).

With long-term exposure
most hearing level changes
first 15 years, while at 5
linear with exposure time
women. Changes from men

Asbestos Exposure During
Hyg. Assn. J. 25:264 (

Exposures resulting in
naval shipyard insulation
since the cases took a mean
allowable concentration
There is no agreement on
asbestos. Shipboard pipe
employees should wear respirators

Diffuse Mesothelioma in a
Area. W. G. Owen. *J.*

Seventeen cases of diffuse
was sought of exposure
mesothelioma are discussed
described. Exposure to
cases examined, and asbestosis
of lung tissue examined
is a major factor in the

The New Federal Clean
Control Assn. 57th Annual

The Clean Air Act (1970)
assistance and leadership
Federal, State, regional

ABSTRACTS

Left Axis Deviation
W. A. Millhon and
67-771 (Dec. 1963).

left axis deviation
ars were compared
me age distribution
this later. Men with
ial disease or dia-
amination were ex-

ovascular disease or
ation compared to
trol group. A more

ation or a positive
34% of the study
with NAD.

group compared to
both hypertension

in 7% of the study

2% of the study

sease in the study

significantly dif-

history of cardio-

uplicated marked
ns.

edicine, July 1964.

h Division of Esso

Permanent Threshold Shift Changes Produced by Noise Exposure and Aging. R. Gallo and A. Glorig. *Am. Ind. Hyg. Assn. J.* 25:237 (May-June 1964).

With long-term exposures to industrial noise of 90 db octave band SPL, most hearing level changes at 3,000, 4,000, and 6,000 cps occur in the first 15 years, while at 500, 1,000, and 2,000 cps, hearing level change is linear with exposure time. Men have greater hearing level changes than women. Changes from noise and aging may not be differentiable.

Asbestos Exposure During Naval Vessel Overhaul. W. T. Marr. *Am. Ind. Hyg. Assn. J.* 25:264 (May-June 1964).

Exposures resulting in 6 cases of asbestosis among 80 employees of a naval shipyard insulation shop were not picked up in time by X-rays since the cases took a minimum of seven years to develop. The maximum allowable concentration for short duration massive exposures is unknown. There is no agreement on the relative importance of short and long fiber asbestos. Shipboard pipe covering and insulation is hazardous and employees should wear respirators when exposed.

Diffuse Mesothelioma and Exposure to Asbestos Dust in the Merseyside Area. W. G. Owen. *Brit. Med. J.* 2:214 (July 25, 1964).

Seventeen cases of diffuse mesothelioma were investigated and evidence was sought of exposure to asbestos dust. Clinical features of diffuse mesothelioma are discussed and the pathological findings in 12 cases are described. Exposure to asbestos was confirmed in 14 (82%) of the 17 cases examined, and asbestos bodies were found in 7 out of 10 specimens of lung tissue examined. The author concludes that exposure to asbestos is a major factor in the cause of diffuse mesothelioma.

The New Federal Clean Air Act. V. G. MacKenzie. Paper, Air Pollution Control Assn. 57th Ann. Meet., Houston, Texas (June 21-25, 1964) 6 p.

The Clean Air Act (Public Law 88-206) states that Federal financial assistance and leadership is essential for the development of cooperative Federal, State, regional, and local programs to prevent and control air