YOUNGER LABORATORIES
Biochemists . . . Pharmacologists . . . Analysts

5620 TELEGRAPH ROAD
ST. LOUIS 29, MISSOURI
PHONE: TILDEN 6-2540

Certificate of Analysis

July 17th, 1962

C+2 AM

SUBJECT -

Toxicological Investigation Of: Aroclor 2565

Monsanto Sample Number 45
Monsanto Project Number Y-62-43

STUDY CONDUCTED FOR -

Monsanto Chemical Company, St. Louis, Missouri

COPY

EXPERIMENTAL PROCEDURE -

A) Oral L^{+}_{50} (Rats)

The diluted compound was fed by stomach tube to Sprague-Dawley strain albino rats.

After the approximate Minimum Lethal Dose was determined, groups of male and female rats were fed in increasing doses at increments of 0.1 fractional log intervals at four levels designed to blanket the toxicity range thereby supplying data for calculation of the L^{+}_{50} which was done according to a modification of the method of E. J. de Beer.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data, together with the dilution at which the compound was fed, are shown in Table I.

B) Skin Absorption MLD^{+} (Rabbits)

The diluted compound was applied in increasing doses at increments of 0.2 fractional log intervals to the closely clipped, intact skin of New Zealand white female rabbits.

The treated areas were covered with plastic strips and the animals placed in wooden stocks for periods up to twenty-four hours, after which time they were assigned to individual cages.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data, together with the dilution at which the compound was applied, are shown in Table II.

MONS 035085

This confidential report is not to be used in any form of advertising without written permission

SUMMARY -

Aroclor 2565

A) Oral L_{50} (Rats)

The Oral L_{50} for male and female rats was placed at 6310 milligrams per kilogram with lower and upper limits of 5,360 to 7,420 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

B) Skin Absorption MLD (Rabbits)

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 2000 milligrams per kilogram and less than 3160 milligrams per kilogram.

The compound was classed as slightly toxic by skin absorption in female rabbits.

YOUNGER LABORATORIES

BY: FRED M. YOUNGER

The material in this report is to be used in development of the product and may be given to responsible sales contacts, but it is not to be used by them in advertising copy. The source of this material is not to be divulged until it appears in formal publications. No exceptions to the established rule may be made without the approval of the Medical Department in St. Louis. Customers' inquiries regarding matters of toxicity are to be referred as before to the Medical Department in St. Louis for reply.

— Monsanto Chemical Company

T A B L E I

THE ORAL LD_{50} OF 'Aroclor 2565' FOR RATS

Sample Fed As A 20.0% Solution In Corn Oil

Animal No. - Sex	Weight Gm.	Dose Mg./Kg.	Fate
1- Female	195	5,010	Survived
2- Female	210	5,010	Died
3- Female	215	5,010	Survived
4- Male	230	5,010	Survived
5- Male	220	5,010	Survived
6- Male	240	6,310	Survived
7- Female	215	6,310	Died
8- Female	230	6,310	Survived
9- Female	210	6,310	Survived
10- Male	240	6,310	Died
11- Male	225	7,940	Died
12- Male	250	7,940	Survived
13- Female	210	7,940	Died
14- Female	215	7,940	Died
15- Female	225	7,940	Died
16- Male	235	10,000	Died
17- Male	240	10,000	Died
18- Male	230	10,000	Died
19- Female	230	10,000	Died
20- Female	215	10,000	Died

DISCUSSION -

The Oral LD_{50} for male and female rats was placed at 6310 milligrams per kilogram with lower and upper limits of 5360 to 7420 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

Survival time was two to six days with most deaths occurring in three days.

Toxic symptoms included weight loss, increasing weakness, dyspnea, and salivation.

At autopsy there was inflammation of the gastric mucosa and liver congestion macroscopically.

T A B L E II

THE MINIMUM LETHAL DOSE OF 'Aroclor 2565'

BY SKIN ABSORPTION IN RABBITS

Sample Applied As A 20.0% Solution In Corn Oil

Animal No. - Sex	Weight Kg.	Dose Mg./Kg.	Weight Change 5 Days Later Kg.	Fate
1 - Female	2.3	794	0.0	Survived
2 - Female	2.1	1260	0.0	Survived
3 - Female	2.5	2000	- 0.2	Survived
4 - Female	2.3	2000	- 0.1	Survived
5 - Female	2.5	3160	- 0.2	Died - 14 Days
6 - Female	2.4	5010	- 0.3	Died - 8 Days

DISCUSSION -

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 2000 milligrams per kilogram and less than 3160 milligrams per kilogram.

The compound was classed as slightly toxic by skin absorption in female rabbits.

Survival time was eight to fourteen days.

Toxic symptoms included weakness in animals #5 and #6 after several days, reduced appetite, lethargy, and weight loss.

At autopsy there was liver discoloration and pulmonary hyperemia macroscopically.