PROJECT NO: Y-70-43
REPORT FILE

274

YOUNGER LABORATORIES
INCORPORATED
Consulting and Analytical Services
CHEMICAL . . . MICROBIOLOGICAL . . . BIOLOGICAL

128 CLIFF CAVE ROAD
SAINT LOUIS, MO., 63129
PHONE: (314) 846-2540

Certificate of Analysis

May 25th, 1970

SUBJECT - NERAL A-2.00

Toxicological Investigation Of: MCS 1009

Monsanto Sample Number 47

Monsanto Project Number Y-70-43

STUDY CONDUCTED FOR -

Monsanto Company, St. Louis, Missouri

EXPERIMENTAL PROCEDURE -

A) Oral LD_{50} (Rats, Mixed Sex)

The undiluted compound was fed by stomach tube to Sprague-Dawley strain albino male and female rats.

After the approximate Minimum Lethal Dose was determined, groups of male and female rats were fed in increasing doses at increments of 0.1 fractional log intervals at five levels designed to blanket the toxicity range thereby supplying data for calculation of the LD_{50} which was done according to the method of E. J. de Beer.

Observations were made for toxic signs and the viscera of the test animals were examined macroscopically.

The data are shown in Table I.

B) Skin Absorption MLD (Rabbits, Mixed Sex)

The undiluted compound was applied in increasing doses at increments of 0.2 fractional log intervals to the closely clipped, intact skin of New Zealand white male and female rabbits.

The treated areas were covered with plastic strips and the animals held in wooden stocks for periods up to twenty-four hours, after which time they were assigned to individual cages.

Observations were made for toxic signs and the viscera of the test animals were examined macroscopically.

The data are shown in Table II.

MONS 035166

This confidential report is not to be used in any form of advertising without written permission

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 2 (5/25/70) - Y-70-43

EXPERIMENTAL PROCEDURE - (Continued)

C) Skin Irritation (Rabbits, Mixed Sex)

0.5 Milliliter of undiluted sample was applied to the clipped, intact skin of New Zealand white male and female rabbits under a one inch by one inch square patch, two single layers thick. The patches were held in place with adhesive tape. The trunk of each animal was wrapped with plastic strips, to retard evaporation and avoid contamination, for the twenty-four hour exposure period.

Observations were made over a period of seven days for irritation.

The data, scored according to the method of Draize, Woodard and Calvery (Journal of Pharm. and Exp. Therapeutics, Volume 82, December, 1944) are shown in Table III.

D) Eye Irritation (Rabbits, Mixed Sex)

0.1 Milliliter of undiluted sample was placed in the conjunctival sac of the right eye of each of three albino male and female rabbits and observations made over a period of seven days for inflammation.

The eyes were rinsed with warm isotonic saline solution after twenty-four hours.

The data, scored according to the method of Draize, et al, are shown in Table IV.

E) Vapor Inhalation (Male Rats) - Sample Heated To 250° F.

Six mature male rats were placed in a metal chamber of 35-liters capacity and exposed for six hours to a concentrated atmosphere of vapors produced by passing a stream of air through 174.6 grams of the compound contained in a 350-milliliter erlenmeyer flask. The sample was maintained at a temperature of 250° F. by immersing the flask in a heated sand bath.

Vapors from the flask passed into a one liter bottle to remove droplets and then into the chamber.

Air flow through the sample was 4.0 liters per minute as measured by a calibrated rotameter. This was sufficient to violently agitate the liquid. No supplementary air was introduced inasmuch as the above supply was ample for the animals oxygen requirements.

The animals were observed for behavior during exposure and for ten days following exposure. The viscera of the test animals were examined macroscopically.

The data are shown in Table V.

SUMMARY -

MCS 1009

A) Oral LD_{50} (Rats, Mixed Sex)

The Oral LD_{50} for male and female rats was placed at 7700 milligrams per kilogram with lower and upper limits of 7000 to 8470 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 3 (5/25/70) - Y-70-43

SUMMARY - (Continued)

B) Skin Absorption MLD (Rabbits, Mixed Sex)

The Minimum Lethal Dose by Skin Absorption in male and female rabbits was found to be greater than 1260 milligrams per kilogram and less than 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in male and female rabbits.

C) Skin Irritation (Rabbits, Mixed Sex)

The compound was classed as a moderate irritant when applied undiluted to intact skin of male and female rabbits.

The average maximum score was 3.6 out of a possible 8 in twenty-four hours.

D) Eye Irritation (Rabbits, Mixed Sex)

The compound was classed as a slight eye irritant in male and female rabbits.

The average maximum score was 8.6 out of a possible 110 in one hour.

E) Vapor Inhalation (Male Rats) - Sample Heated To 250° F.

All animals survived the six hour exposure as well as the following ten day observation period.

It was concluded that the vapors were practically non-toxic under conditions of the test.

YOUNGER LABORATORIES, INC.

BY: MELVIN D. BIRCH

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 4 (5/25/70) - Y-70-43

TABLE I

THE ORAL LD_{50} OF 'MCS 1009' FOR RATS

Sample Fed Undiluted

Animal No. - Sex	Weight Gm.	Dose Mg./Kg.	Fate
1- Female	200	5,010	Survived
2- Male	240	5,010	Survived
3- Female	215	5,010	Survived
4- Male	220	5,010	Survived
5- Female	225	5,010	Survived
6- Male	230	6,310	Survived
7- Female	250	6,310	Died
8- Male	270	6,310	Died
9- Female	225	6,310	Survived
10- Male	260	6,310	Survived
11- Female	210	7,940	Survived
12- Male	220	7,940	Died
13- Female	230	7,940	Died
14- Male	265	7,940	Survived
15- Female	215	7,940	Died
16- Male	245	10,000	Survived
17- Female	220	10,000	Died
18- Male	265	10,000	Survived
19- Female	195	10,000	Died
20- Male	260	10,000	Died
21- Female	195	12,600	Died
22- Male	215	12,600	Died
23- Female	230	12,600	Died
24- Male	250	12,600	Died
25- Female	210	12,600	Died

DISCUSSION -

The Oral LD_{50} for male and female rats was placed at 7700 milligrams per kilogram with lower and upper limits of 7000 to 8470 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

Survival time was two to eight days with most deaths occurring within four days.

Toxic signs included reduced appetite and activity, increasing weakness, collapse, and death.

At autopsy there was hemorrhagic lungs, slight liver discoloration, and gastro-intestinal inflammation (acute in some cases).

Surviving animals were sacrificed ten days after dosing. Macroscopic examination revealed slight lung congestion in upper dosage levels; viscera of animals at lower dosage levels appeared normal.

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 5 (5/25/70) - Y-70-43

TABLE II

THE MINIMUM LETHAL DOSE OF 'MOS 1009'
BY SKIN ABSORPTION IN RABBITS

Sample Applied Undiluted

Animal No. - Sex	Weight Kg.	Dose Mg./Kg.	Weight Change 5 Days Later Kg.	Fate
1 - Male	2.6	794	- 0.2	Survived
2 - Female	2.2	1260	- 0.1	Survived
3 - Male	1.9	1260	0.0	Survived
4 - Female	1.8	1260	- 0.1	Survived
5 - Male	1.9	2000	- 0.3	Died - 10 Days
6 - Female	1.8	2000	0.0	Survived
7 - Male	1.8	2000	- 0.2	Survived
8 - Female	2.2	3160	-----	Died -- 4 Days
9 - Male	2.1	5010	- 0.3	Died -- 7 Days
10 - Female	2.8	7940	- 0.4	Died -- 5 Days
11 - Male	2.7	7940	-----	Died -- 4 Days

DISCUSSION -

The Minimum Lethal Dose by Skin Absorption in male and female rabbits was found to be greater than 1260 milligrams per kilogram and less than 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in male and female rabbits.

Survival time was four to ten days.

Toxic signs included reduced appetite and activity, lethargy, increasing weakness, slight tremors, collapse, and death.

At autopsy there was lung hyperemia, jaundiced and mottled liver, enlarged gall bladder, and gastreintestinal inflammation.

Surviving animals were sacrificed fourteen days after dosing. Macroscopic examination revealed lung congestion, mottled and discolored liver, enlarged gall bladder, and slight gastreintestinal inflammation.

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 6 (5/25/70) - Y-70-43

TABLE III

SKIN IRRITATION IN RABBITS AFTER APPLICATION OF
'MCS 1009'

Sample (0.5 Milliliter) Applied Undiluted

Animal No. - Sex	Numerical Evaluation At The End Of					
	1 Hour	24 Hours	48 Hours	72 Hours	120 Hours	168 Hours
1 - Female	1	4	3	2	1	0
2 - Male	1	3	3	1	1	0
3 - Female	1	4	4	2	0	0
Average	1.0	3.6	3.3	1.6	0.6	0.0

DISCUSSION -

The compound was classed as a moderate irritant when applied undiluted to intact skin of male and female rabbits.
The average maximum score was 3.6 out of a possible 8 in twenty-four hours.

Observations following application -

1-Hour:	Barely perceptible erythema; no edema
24-Hours:	Well-defined erythema, very slight to slight edema
48-Hours:	Well-defined erythema, very slight to slight edema
72-Hours:	Slight erythema, very slight edema in two instances
120-Hours:	Slight erythema in two instances; no edema
168-Hours:	No erythema or edema; zero readings

Note: Sample had a slight defatting effect on the skin causing the skin to scale off in seven to ten days. There was no injury in depth.

The material in this report is to be used in development of the product and may be given to responsible sales contacts, but it is not to be used by them in advertising copy. The source of this material is not to be divulged until it appears in formal publications. No exceptions to the established rule may be made without the approval of the Medical Department in St. Louis. Customers' inquiries regarding matters of toxicity are to be referred as before to the Medical Department in St. Louis for reply.

— Monsanto Company

MONS 035171

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 7 (5/25/70) - Y-70-43

TABLE IV

EYE IRRITATION IN RABBITS AFTER APPLICATION OF
'MCS 1009'

Sample (0.1 Milliliter) Applied Undiluted

Animal No. - Sex	Numerical Evaluation At The End Of					
	1 Hour	24 Hours	48 Hours	72 Hours	120 Hours	168 Hours
1 - Male	8	2	0	0	0	0
2 - Female	10	2	0	0	0	0
3 - Male	8	2	0	0	0	0
Average	8.6	2.0	0.0	0.0	0.0	0.0

DISCUSSION -

The compound was classed as a slight eye irritant in male and female rabbits. The average maximum score was 8.6 out of a possible 110 in one hour.

Observations following application --

Immediate:	Discomfort was slight
10-Minutes:	Mild erythema and discharge; no edema
1-Hour:	Moderate erythema, moderate to copious discharge; no edema
24-Hours:	Slight erythema; no edema or discharge
48-Hours:	Eyes normal; zero readings
72- 168-Hours:	Eyes normal; zero readings

To: Monsanto Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 8 (5/25/70) - Y-70-43

TABLE V

INHALATION OF 'MCS 1009' VAPORS BY MALE RATS
(SAMPLE HEATED TO 250° F.)

Average Temperature Inside Chamber ..... 74° F.

Average Relative Humidity Inside Chamber ..... 54 %

Amount of Sample --	To start	174.6 Grams	
	Recovered	174.0 Grams	
	Total Vaporized	0.6 Gram	(0.3 %)
	Recovered from condenser	0.0 Gram	(0.0 %)
	Weight of vapors entering chamber	0.6 Gram	(0.3 %)

Animal No. - Sex	Fate	Observations During Exposure
1 - Male	Survived	No immediate effect ...
2 - Male	Survived	Normal respiration ...
3 - Male	Survived	No nasal or ocular discharge ...
4 - Male	Survived	No weakness or drowsiness of consequence.
5 - Male	Survived	
6 - Male	Survived	

DISCUSSION -

All animals survived the six hour exposure as well as the following ten day observation period.

It was concluded that the vapors were practically non-toxic under conditions of the test.

Observations following exposure --
The animals appeared normal when removed from the chamber. The ten day observation period was uneventful. There were no bronchial rales or other evidence of respiratory complications. Weight gain was normal.

Surviving animals were sacrificed ten days following exposure. The viscera appeared normal by macroscopic examination.