Monsanto

FROM (NAME & LOCATION) W. B. Papageorge - B2SK

DATE October 21, 1975

cc R. E. Keller
J. P. Mieure
W. R. Richard

SUBJECT G. E. REQUESTS

REFERENCE Memo JPM to DW, 10/10/75

TO D. Wood

(1) TO R Mauch / QT Lepore
(2) Bill Hammerman
(3) WR File Arochlor Environment

In responding to Items (1) and (2) in Jim Mieure's memo, I propose that we forward a copy of the Fishbein paper (copies are attached) and comment on biphenyl purity as follows:

"Identifiable materials present in our biphenyl contribute to the presence in the polychlorinated biphenyl mixture (the material of primary concern to you) of the following:

| | |
|---|---|
| Chlorinated naphthalene | <3 ppm |
| Chlorinated methyl naphthalene | <6 ppm |
| Chlorinated methyl biphenyl | 100 ppm max. |
| Chlorinated dibenzofurana | <.5 ppm |
| Unidentified | 60 ppm max." |

W B Papageorge
W. B. Papageorge

/pd

MONS 202107

IN 1C REV 1 72

Journal of Chromatography
Chromatographic Reviews
Elsevier Publishing Company, Amsterdam
Printed in The Netherlands

# CHROMATOGRAPHIC AND BIOLOGICAL ASPECTS OF POLYCHLORINATED BIPHENYLS

LAWRENCE FISHBEIN

*National Institute of Environmental Health Sciences, National Institutes of Health, Public Health Service, and Department of Health, Education and Welfare, Research Triangle Park, N.C. 27709 (U.S.A.)*

(Received August 5th, 1971)

CONTENTS

I. Introduction . . . . . 345
II. Ecological aspects . . . . . 347
III. Toxicological aspects . . . . . 351
IV. Column and thin layer chromatography of polychlorinated biphenyls . . . . . 353
V. Gas liquid chromatography . . . . . 359
VI. Gas liquid chromatography-mass spectroscopy . . . . . 400
VII. Analyses of chlorinated naphthalenes and dibenzofurans . . . . . 418
VIII. Summary . . . . . 423
References . . . . . 423

## I. INTRODUCTION

Despite the fact that polychlorinated biphenyls (PCB's) have been available commercially for forty years, it is only within the last five years that they have been recognized to be of environmental concern.

Polychlorinated biphenyls are produced by a comparatively small number of manufacturers, marketed under a number of commercial trade names, *e.g.*, Aroclor, Clophen and Phenoclor, and are represented as "a series of inert, chemically resistant, fire-retarding plasticizers compatible with a wide variety of resins, varnishes, waxes and paints; they vary from mobile, oily liquids to white crystals and hard transparent resins"[1]. The series of Aroclors (Monsanto) are marketed under various numbers and consist of mixtures of chlorinated biphenyls and terphenyls. The first two digits represent the molecular type: 12—chlorinated biphenyls, 25 and 44—blends of chlorinated biphenyls and chlorinated terphenyls (75% biphenyl and 60% biphenyl, respectively), 54—chlorinated terphenyls. The last two digits give the weight percent of chlorine, *e.g.*, Aroclor 1242 is a chlorinated biphenyl containing 42% chlorine.

The viscosity of the PCB's increases in direct proportion to the chlorine content from very fluid liquids to viscous products and solids. In the commercial process for PCB manufacture, biphenyls are chlorinated with anhydrous chlorine with either iron filings or ferric chloride as the catalyst, the byproduct is hydrogen chloride and



MONS 202108

[illegible] is a mixture of several PCB's. In the process of replacing hydrogen atoms [illegible] of chlorine, a large number of substitution combinations arise [illegible].



For example, three monochloropiphenyl isomers are possible, twelve dichlorophenyl isomers, twenty-one trichlorophenyl isomers, and so on. Theoretically, 210 compounds can be prepared by this substitution process. (A typical PCB example would be 2,4,6,2,4-pentachlorophenyl.) Mass spectroscopic studies[2] of Aroclor [illegible] revealed the presence of eleven isomers [illegible], five containing six chlorine atoms, [illegible] containing seven chlorine atoms and one containing eight, while Aroclor 12[illegible] was found to contain eighteen compounds [illegible], one containing three chlorine atoms, [illegible] containing four chlorines, four containing five chlorines, five containing six chlorines, and four containing seven chlorines[3]. The annual production of PCB's in the Western world [illegible] most recently was estimated at 100 million pounds.

The chemical properties that make PCB's desirable industrial chemicals are [illegible] excellent thermal stability, their strong resistance to both acidic and basic hydroxides and action of corrosive chemicals and their *general* inertness. They are [illegible] in water but possess a low finite vapor pressure. Their boiling points range from 278° for Aroclor 1221 to 415° for Aroclor 1268. All are stable to prolonged [illegible]

TABLE 1

USES OF POLYCHLORINATED BIPHENYLS

| Material with which chlorine is combined | Aroclor used with (wt %) | Use |
|---|---|---|
| [illegible] chloride | Aroclor 1248, 1254 and 1260 (7–8%) | Secondary plasticizer to improve flame retardance and chemical resistance |
| [illegible] lacquers | Aroclor 1260 (7%) | Co-plasticizer to enhance resistance |
| [illegible] acetate | Aroclor [illegible], 1232 and 1242 (11%) | Improve quick track and fiber tear properties |
| [illegible] acetate | Aroclor 1254 (11%) | Pressure sensitive adhesive |
| [illegible] resins | Aroclor [illegible] and 1248 (20%) | Increase chemical and oxidation resistance and adhesive quality |
| [illegible] resins | Aroclor 1260 (10–15%) | Effective and economical fire retardant |
| | Aroclor 1260 (10–20%) | Increases strength of fiber glass inforced polyester resins |
| [illegible] | Aroclor [illegible] | Plasticizer |
| [illegible] rubber | Aroclor 1254 (5–10%) | Enhances resistance, flame retardance and improves electrical insulation properties |
| [illegible] | Aroclor 1254 [illegible] | Improves chemical resistance |
| [illegible] | Aroclor [illegible] | Fire retardant |
| | Aroclor [illegible] | Injection moldings |
| [illegible] | Aroclor [illegible] | Plasticizer [illegible] compositions |
| [illegible] | Aroclor [illegible] | Improves water and alkali resistance |
| [illegible] | Aroclor [illegible] | Moisture and flame resistance |



[illegible] at 150° and the lower Aroclors can be distilled at atmospheric pressure without appreciable decomposition.

The largest single use of PCB's is related to their electrical properties, as coolant insulation fluids in transformers and capacitors. Other uses of PCB's include impregnation of cotton and asbestos for braided insulation of electrical wiring, plasticizers of vinyl chloride and polymer freons, as plasticizer in wire and cable coatings and in ballasts for fluorescent fixtures. Because of their thermal stability and fire resistance the PCB's also find application in high pressure hydraulic fluids, heat transfer agents, machine tool cutting oils, specialized lubricants and gasket sealers. Miscellaneous uses include formulations in epoxy paints, resins and chlorinated rubber, printing inks, waxes, synthetic adhesives, textile dyes, protective coatings for wood, metal and concretes, as sealers in water-proofing compounds and putty, and in carbonless reproducing paper.

PCB's have been incorporated into pesticide formulations, especially with such insecticides as DDVP, lindane[4,5], chlordane, aldrin, dieldrin and toxaphene to suppress their vaporization and hence extend their "kill-life" and have also been shown to increase the insecticidal properties of DDT[6].

Table 1 lists a number of uses of the Aroclor polychlorinated biphenyls and the material and percentages with which they are combined.

## II. ECOLOGICAL ASPECTS

The first identification of polychlorinated biphenyls in regard to ecology was by JENSEN[7], who identified PCB's in the bodies of 200 pike taken from different parts of Sweden, in other fish and in eagle feathers collected in 1944. PCB's along with DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)-ethane] are now reported to be the most abundant of the chlorinated aromatic pollutants in the global ecosystem[8]. Extracts of sea eagles, pike and salmon[9] as well as in British[9] and Canadian[10–12] wildlife contained PCB's, and in the former instance it was found that in birds' liver and eggs the PCB residues were greater than the organochlorine pesticide residues. PCB's have also been found in fish, mussels and birds from the River Rhine and The Netherlands coastal areas[3] and in marine animals and wildlife in Sweden, England and the USA[2,8,11,13].

Polychlorinated biphenyls have also been found in human adipose tissue[14], samples of human milk[15] and in foods (margarine, vegetable oils and particularly fish[16]).

Essentially, the same type of residue pattern is becoming apparent for the polychlorinated biphenyls that has been found for the persistent organochlorine insecticides. The PCB's are extremely stable, chemically fat soluble and hence persistent in the environment. The four possible pathways by which PCB's could be dispersed in the environment include: (a) because of their numerous manufacturing applications and general inertness, PCB's could be flushed as wastes into rivers, lakes, etc., to pollute fish and other wildlife; (b) via industrial smoke, in exhaust from aircraft engines, incineration and combustion of PCB-containing products, and leaching of plasticizer from plastic objects in waste disposal areas; (c) via insecticidal formulations to enhance kill-life (described above), and (d) via direct contamination of feeds and foodstuffs through leakage of heat transfer fluids.

Information relative to the biological decomposition of the PCB's is scant and

MONS 202109

TABLE 2

PATHOLOGIC CHANGES INDUCED BY PCB's[a]

| *Treatment* | *Animal* | *Liver* | *Kidney* | *Pericardium and peritoneum* | *Other observable changes* | *Reference* |
|---|---|---|---|---|---|---|
| Single oral dose of 1 g/kg (42% Cl) | Guinea pig<br>Rat<br>Rabbit | Small fat droplets through lobules, slight to moderate central atrophy, focal necrosis noted in a few animals | Essentially normal | No noteworthy changes | Adrenals, spleen and pancreas showed no noteworthy changes | 25 |
| 500 mg/kg daily for 6 days (54% Cl) | Rat | Cells swollen, hyaline granules present, most died within few days | | | | 27 |
| 500 mg/kg daily for up to 8 months (54% Cl) | Rat | Enlarged (33% weight increase), large number of hyaline globules in cytoplasm, several died during experiment | | | | 27 |
| 25, 50 and 100 ppm in diet for 15 days (21–60% Cl Aroclors) | Rat | Increase in weight, effect increasing with increasing chlorine content. Aroclor 1232 10%, 1242 12%, 1254 14%, 1268 24% at 50 ppm | | | | 32 |
| 100 ppm in diet<br>200 ppm in diet<br>400 ppm in diet<br>500 ppm in diet | Chicken | No effect<br>No effect<br>Colored and mottled<br>Diseased | Damaged | Slight<br>Hydropericardium<br>Hydropericardium<br>Hydropericardium, hydropericardium enlarged | | 39 |
| 300 and 400 ppm in diet for 3 weeks (42% Aroclor) | Chicken | No changes noted | Paleness at 300 ppm, extensive hemorrhage, and enlargement at 400 ppm | Increased fluid in pericardial sac at the higher concentration | Reduction of production, enlargement of adrenal and small spleen at low concentrations, at higher concentrations pale cream-colored pancreas, adrenals hemorrhagic | 36 |
| Various doses (34% Cl, Aroclor) | Bengalese finch | No weight changes | Weight was 32.4% of brain weight for controls and 53.5% for those dying from PCB poisoning | Slight weight increase, a few showed liquid in pericardial sac | | 33 |
| 400 ppm in diet for 60 days (60% Cl[a]) | Chicken | Centrolobular necrosis (cpds. 1 and 2); liver weight increased from 2.76 g/100 g to 4.31 g/100 g (cpd. 3); fatty degeneration | Tubular dilatation (cpds. 1 and 2), rare with cpd. 3 | Hydropericardium common with cpds. 1 and 2, rare with cpd. 3. | Increased porphyria, spleen small with reduction of red pulp and atrophy of white pulp (cpds. 1 and 2), spleen decreased from 0.146 g/100 g to 0.130 g/100 g (cpd. 3) | 34 |

[a] Phenoclor DP6 (cpd. 1), Clophen A60 (cpd. 2) and Aroclor 1260 (cpd. 3) were used. Differential effects noted under cpd. numbers. All chickens died on cpds. 1 and 2 within 60 days, only 15% mortality on cpd. 3.





MONS 202110

it was suggested[5] that they are likely more stable than DDT and its metabolites since the PCB's lack the ethane component between the aromatic rings which is the site of action of most of the transformations of DDT. This factor coupled with their physical and chemical characteristics for persistence indicates that these materials are capable of biological magnification up the food chain. (Although the concentration of the PCB's are generally in the p p b[a] range in the environment, their high lipid solubility results in their accumulation in fatty tissues of lower animals and marine life.) Because of the very low aqueous solubility of the PCB's, when they are discharged into a river or lake, they will accumulate on the sediment in relatively high concentration and redissolve very slowly.

## III. TOXICOLOGICAL ASPECTS

Compared to the chlorinated hydrocarbon pesticides, the toxicology of the polychlorinated biphenyls remains rather poorly known. The systemic effects in humans and deaths resulting from exposure to the PCB's and polychlorinated naphthalene was described by GREENBERG *et al.*[18] and the chloroacnegen effects reported by SCHWARTZ and co-workers[19-21], JONES AND ALDEN[22] and VON WIDDEL *et al.*[23]

SAX[24] reported the toxicity of PCB's to man as moderate to high, especially when inhaled as vapors. Animal experiments with rats and guinea-pigs have shown that ingestion of PCB may give rise to liver injury[25,26]. Oral administration of chlorobiphenyls resulted in enlargement of the liver and vacuolar or fatty degeneration of liver cells.

BENNETT *et al.*[27] have previously shown that chlorinated biphenyl resulted in liver changes markedly different from those caused by other well known toxic agents such as carbon tetrachloride and chloroform poisoning and thought that this type of injury was persistent with slow recovery. The effects on mouse and monkey liver of long-term oral administration of chlorobiphenyls (1.5 mg/day or more) have been reported by NISHIZUMI[28].

One PCB has been shown to produce chick hydropericardial edema and growth depression proportional to dietary levels (200 and 400 p p m.) of the toxicant[29,30]. These changes were accompanied by such pathological changes as enlarged hemorrhagic kidney, enlarged adrenal gland, small spleen, cream-colored pancreas, dermatitis, defeathering and transient changes in blood glucose, hemoglobin and hematocrit levels. Hydropericardium and abdominal edema have also been observed in Japanese quail[2] after PCB ingestion. Toxic and teratogenic effects of PCB's in chick embryos have also been reported by MCLAUGHLIN *et al.*[31].

Analogously with the chlorinated hydrocarbon pesticides, the most important effects are long-range sublethal effects. The pathologic changes in various organs are summarized in Table 2, illustrating some interesting differences between mammals and birds. For example, the most striking findings in mammals are alterations to the liver, whereas fluid in the pericardial sac, kidney damage and reduced spleen are found in birds.

The toxicity of eleven Aroclors in terms of oral $LD_{50}$ (rats) and skin MLD (rabbits) is summarized in Table 3.

[a] Throughout this article the American (10[9]) billion is meant.

Polychlorinated biphenyls and triphenyls have been found to be estrogenically active[36]. In a series of PCB's the compounds containing up to 48% chlorine were inactive. On a weight basis PCB compounds such as Aroclor 1221 were shown to have no estradiol degrading potential (in pigeon liver homogenates) about five times that of *p,p'*-DDE or technical grade DDT[8,37]. Other examples of induction of hepatic hydroxylating enzymes by PCB's have been reported in the rat[38] and the American kestrel (*Falco sparverius*)[39].

VILLENEUVE *et al.*[35] studied the effects of PCB administration on microsomal enzyme activity in pregnant rabbits. The no-effect level of Aroclor 1254 for enzyme induction in the pregnant rabbit is between 1.0 and 10 mg/kg body weight when administered for twenty-eight days during gestation. Aroclor 1221 did not induce any enzyme activity in the does, foetus or placenta, so its no-effect level must be considered higher than that for Aroclor 1254. Placental transfer was shown to occur for both Aroclor 1254 and 1221 but does not cause any changes in the biochemical or physiological parameters measured, *e.g.*, total amount of Vitamin A stored per liver, protein levels, aniline hydroxylase enzyme activity, serum cholesterol, no effect in reproductive processes. The drug metabolizing enzymes aniline hydroxylase and aminopyrine *n*-demethylase were both induced by 10 mg/kg Aroclor 1254.

PCB's as inducers of hepatic enzymes (*e.g.*, increasing the rate of circulating estradiol in the liver) may be responsible for aberrations in calcium metabolism in certain species of birds[8] and are generally considered to be more of a potential threat than DDT to our declining bird populations, especially for predatory birds that accumulate fairly high levels of PCB's. ANDERSON *et al.*[11] have suggested evidence that PCB's affected eggshell thickness although to a lesser extent than DDE.

However, VERMEER AND REYNOLDS[40] found no significant correlation between shell thickness and PCB residues for the great blue heron (*Ardea herodias*) and PEAKALL[41] found no significant effect of PCB's in the eggshells of ring doves.

FRIEND AND TRAINER[12] reported on the interaction of PCB with duck hepatitis virus. Ten-day old mallard ducklings fed a polychlorinated biphenyl (Aroclor 1254) at concentrations of 25, 50 and 100 p.p.m. for ten days suffered no apparent clinical

TABLE 4

PERCENT INHIBITION OF HUMAN CELL* GROWTH OBTAINED WITH ENVIRONMENTAL CHEMICALS

| *Chemical* | $ID_{50}$ (p.p.m.)* | |
|---|---|---|
| | *H* | *F* |
| *p,p'* DDT | 55 | 50 |
| *o,p'* DDT | 50 | — |
| Parathion | 44 | 64 |
| PCB (Aroclor 1254) | 64 | 110 |
| Caffeine | 850 | 650 |
| Aspirin | 900 | 1000 |
| Acetylsalicylic acid | 450 | — |

* Human cells: H = HeLa (established cell line); F = normal diploid skin fibroblasts.
† Concentration of additive in p.p.m. that produces 50% inhibition ($ID_{50}$) in comparison with untreated controls.



intoxication. When these birds were challenged five days later with duck hepatitis virus (DHV) they suffered significantly higher mortality than birds which were not exposed to the polychlorinated biphenyl (*e.g.* 14% among virus controls *versus* 35 to 65% among groups receiving PCB plus DHV). This study illustrates one of the potential effects of sublethal concentrations of chemical pollutants and emphasizes the real differences that exist between "sublethal" and "no-effect" concentrations of pollutants.

The effects and interactions of environmental chemicals (aspirin, caffeine, Aroclor 1254, carbaryl, parathion, DDT and several insecticide metabolites) on human cells (HeLa and skin fibroblasts) in tissue culture were described by LITTERST AND LICHTENSTEIN[13]. The dosage causing a 50% inhibition in culture growth, and the interaction between these agents and their effect on protein and nucleic acid synthesis were all determined. Table 4 lists the percent inhibition of human cell growth obtained with selected chemicals and Table 5 shows the effect of aspirin, PCB and caffeine on macromolecule synthesis in human cell cultures after 4½ h of exposure to DDT and parathion. Both cell types responded nearly equally to the presence of an individual chemical. Aspirin and caffeine were eight to twenty times less toxic than the other agents studied while the PCB plasticizer Aroclor 1254 was as toxic as DDT.

## IV. COLUMN AND THIN-LAYER CHROMATOGRAPHY OF PCB's

Because of their similarity in structures and chemical properties, PCB's, if

TABLE 5

EFFECT OF ADDITIVES ON MACROMOLECULE SYNTHESIS IN HUMAN CELL CULTURES* AFTER 4½ h OF EXPOSURE TO INSECTICIDES

| *Additive* | *Insecticide* | | | | | |
|---|---|---|---|---|---|---|
| | *None* | | *DDT (75 p.p.m.)* | | *Parathion (50 p.p.m.)* | |
| | *H* | *F* | *H* | *F* | *H* | *F* |
| Protein | | | | | | |
| None | — | — | 70† | 51† | 85 | 94 |
| Aspirin (25 p.p.m.) | 102 | 94 | 47‡ | 45 | 75 | 105 |
| PCB‖ (10 p.p.m.) | 91 | 108 | 59† | 59 | 82 | 86‡ |
| Caffeine (10 p.p.m.) | 93 | 101 | 72 | 138‡ | 66‡ | 64‡ |
| RNA | | | | | | |
| None | — | — | 70† | 89 | 81 | 67‡ |
| Aspirin (25 p.p.m.) | 91 | 87 | 94 | 103 | 72 | 54 |
| PCB (10 p.p.m.) | 100 | 96 | 72 | 86 | 104‡ | 55 |
| Caffeine (10 p.p.m.) | 100 | 91 | 88‡ | 82 | 91 | 75 |
| DNA | | | | | | |
| None | — | — | 77 | 91 | 91† | 87 |
| Aspirin (25 p.p.m.) | 143 | 86 | 84 | 86 | 81‡ | 91‡ |
| PCB (10 p.p.m.) | 93 | 107 | 94 | 86 | 89 | 103 |
| Caffeine (10 p.p.m.) | 104 | 93 | 81 | 84 | 77 | 85 |

* Human cells: H = HeLa (established cell line); F = normal diploid skin fibroblasts.
† Polychlorinated biphenyl (Aroclor 1254).
‡ Differences between treated cultures (as stated) or additive and control cultures (the data treated with asterisk) were significant at the 5% level.
§ Differences between treated cultures (insecticide plus additive) and insecticide cultures (the data treated with asterisk) were significant at the 5% level.

MONS 202112

present in a sample are carried through the usual pesticide extraction and screening procedures and are frequently mistaken for DDT and analogs in monitoring [illegible]




REYNOLDS[12,13] described an activated Florisil column technique which separated heptachlor, aldrin, DDE and PCB with the first elution (60 ml of n-hexane) from lindane, and heptachlor epoxide, DDD and DDT with the second elution (40 ml of 50% diethyl ether in hexane).

KOLMAN *et al.*[2] eluted a number of apolar compounds including DDE and PCB on activated Florisil columns using n-hexane, then dieldrin and endrin with 10% diethyl ether in hexane.

The separation of PCB's from DDT and its analogs and other common pesticides by column chromatography on silicic acid–Celite was reported by ARMOUR AND BURKE[14]. Aldrin and PCB were eluted with the first fraction (250 ml of petroleum ether) and lindane, heptachlor, heptachlor epoxide, dieldrin, endrin, p,p'-DDE, o,p'-DDT, p,p'-DDT and p,p'-DDD with the second fraction (200 ml of acetonitrile–




[illegible]





hexane–methylene chloride (1:19:80). Determination of PCB's and pesticides could also be made on separate column eluates without cross-interference with recoveries of Aroclors 1254 and 1260 and of several chlorinated pesticides ranging from 76 to 100% and 80 to 100%, respectively. The effect of water content of silicic acid on elution and separation of Aroclor 1260 and DDT and analogs is shown in Fig. 1. The maximum margin of separation of PCB from pesticides with good reproducibility of elution pattern was obtained with silicic acid containing 3% water. A more polar eluent was required for the complete elution of p,p'-DDT and p,p'-TDE, when the water content was 5% less.

A semiquantitative determination of PCB's in tissue samples by TLC was described by MULHERN *et al.*[16]. Cleanup by hexane–acetonitrile partitioning and Florisil column chromatography was performed on samples before oxidative treatment[17] to convert DDE to DCBP (4,4-dichlorobenzophenone), then the PCB components were determined semiquantitatively (with a lower limit of sensitivity of 0.2 μg) by TLC with no prior separation from chlorinated pesticides required. TLC separation was accomplished on silver nitrate incorporated Aluminum Oxide G plates developed with 5% benzene in hexane. The spots were detected using a spray consisting of 5 ml of water and 10 ml of 2-phenoxy-ethanol diluted to 200 ml with acetone and containing one to five drops of 30% hydrogen peroxide, followed by exposure to germicidal UV light. The $R_F$ values of Aroclor mixtures chromatographed by this TLC system

TABLE 6

$R_F$ VALUES FOR AROCLOR MIXTURES AND PESTICIDES ON $AgNO_3$-INCORPORATED ALUMINUM OXIDE G PLATES DEVELOPED WITH 5% BENZENE IN HEXANE

| Compound | $R_F$ value |
|---|---|
| 1242 | 0.91 |
| 1248 | 0.93 |
| 1254 | 0.93 |
| 1260 | 0.93 |
| 1262 | 0.94 |
| DDE | 0.93 |
| DDT | 0.88 |
| DDD | 0.74 |
| Heptachlor epoxide | 0.58 |
| Dieldrin | 0.48 |
| DCBP | 0.30 |

are listed in Table 6. Fig. 2 illustrates a chromatogram of a pelican egg sample containing oxidized samples as well as Aroclor 1254 and pesticide standards. The lower limit of detection of PCB by this TLC procedure is 0.2 μg.

Analytical data of pelican egg and eagle liver samples are shown in Table 7. PCB was determined by TLC and DDE data were obtained by the GLC procedure of REICHEL *et al.*[18]. The PCB/DDE ratio was approximately 1.0 except for the pelican eggs from California.

ARMOUR AND BURKE[14] used precoated aluminum oxide sheets with both n-heptane and acetone–n-heptane (2:98) as solvent systems for the separation of p,p'-DDT and p,p'-DDE from Aroclor 1254 and 1260. However, the above Aroclors and DDE were not resolved.

MONS 202113




Fig. 2. Thin layer chromatogram of a pelican egg sample. $AgNO_3$ incorporated Aluminum Oxid G. 5% benzene in hexane. 1 = 0.6 g of oxidized sample, 2 = 1.0 g of oxidized sample, 3 = 0.6 g of unoxidized sample, 4 = 1.0 g of unoxidized sample, 5 = 1 µg of Aroclor 1254, 6 = 5 µg of Aroclor 1254, and 7 = 5 µg of each of DCBP, dieldrin, heptachlor epoxide, DDD, DDT and DDE (in ascending order).

The separation and identification of DDT analogs in the presence of polychlorinated biphenyls by two-dimensional TLC was described by FEHRINGER AND WESTFALL[10]. Separation resulted from the different migration patterns and migration distances relative to *p,p'*-DDE that mixtures of PCB's exhibit under two sets of conditions. Two-dimensional TLC was employed using MN-Kieselgel G-HR as the absorbent with *n*-heptane alone and *n*-heptane acetone (98:2) as developing solvents and silver nitrate incorporated into the absorbent layer as the chromogenic agent. Fig. 3 illustrates a plate spotted for first dimension development to be performed in a sandwich chamber saturated in mobile solvent of *n*-heptane. Fig. 4 shows a plate spotted for second dimension development to be performed in a partially saturated tank.

TABLE 7

EXAMPLES OF PCB QUANTITATION BY TLC

| *Sample* | *Location* | *PCB (p.p.m.)* | *DDE (p.p.m.)* | *PCB/DDE* |
|---|---|---|---|---|
| Pelican eggs | Calif. | 12.0 | 113 | 0.09 |
| Pelican eggs | Calif. | 5.0 | 70 | 0.[illegible] |
| Pelican eggs | S.C. | 12.0 | 11.6 | 1.04 |
| Pelican eggs | S.C. | 8.0 | 10.5 | 0.76 |
| Pelican eggs | Fla. | 5.0 | 6.6 | 1.21 |
| Pelican eggs | Fla. | 3.0 | 2.6 | 1.15 |
| Eagle liver | Ariz. | 5.0 | 5.5 | 1.01 |

[illegible] Fig. 5 through 12 illustrate the two-dimensional TLC separation of DDT analogs from Aroclors 1254, 1260, 1221, 1232, 1232, 1232, 1242 and 1265 respectively. Fig. 13 depicts the two-dimensional separation of [illegible] from a sample containing PCB's, kelthane, *p,p'*-DDT, *o,p'*-DDT and *p,p'*-DDE. Figs. 14 and 15 depict the two-dimensional separa-









Fig. 3. Plate spotted for first dimension development to be performed in a sandwich chamber. Mobile solvent: *n*-heptane.

Fig. 4. Plate spotted for second dimension development to be performed in a partially saturated tank. Mobile solvent: *n*-heptane-acetone (98:2). Locations of spots after first dimension development are not indicated since spots are not visible at this point.







Fig. 5. Separation of Aroclor 1254 from the DDT analogs.

Fig. 6. Separation of Aroclor 1260 from the DDT analogs.







Fig. 7. Separation of Aroclor 1221 from the DDT analogs.

MONS 202114



Fig. 9. Separation of Aroclor 1242 from the DDT analogs

Fig. 10. Separation of Aroclor 1248 from the DDT analogs



Fig. 11. Separation of Aroclor 1262 from the DDT analogs

Fig. 12. Separation of Aroclor 4465 from the DDT analogs



Fig. 13. Two dimensional separation of a 6% Florisil eluate from a fish sample containing PCB's, heptachlor, *p,p'*-DDT, *o,p'*-DDT and *p,p'*-DDE



Fig. 14. Two dimensional separation of PCB's isolated from a fish sample by the silicic acid column separation of ARMOUR AND BURKE[45]

respectively (both isolated from fish sample by the silicic acid column separation of ARMOUR AND BURKE[45])

Aroclors 1254 and 1260 are the two of greatest interest in pesticide work because their GLC peaks elute in the same general region as the DDT analogs and chromatograph on many TLC systems at approximately the same $R_F$ value as *p,p'*-DDE.

The synthesis and TLC of twenty-three chlorobiphenyls has been described by HUTZINGER *et al.*[51]. All chlorobiphenyls prepared were homogeneous on silica gel thin-



Fig. 15. Two-dimensional separation of kelthane, *p,p'*-DDT, *o,p'*-DDT and *p,p'*-DDE isolated from PCB's in a fish sample by the silicic acid column separation of ARMOUR AND BURKE[45]

layer chromatograms developed in heptane and gave one peak when chromatographed on a 4% SE-30 column.

Table 8 lists the names, structural formulae, starting material sources, method of purification, melting points and $R_F$ values of the individual chlorobiphenyls.

## V. GAS–LIQUID CHROMATOGRAPHY

The discovery of PCB's in the environment was not accomplished until 1966



MONS 202115

| [illegible] | Formula | Start of material (Source) | Method of purification | Physical data m.p. (°C) | [illegible] |
|---|---|---|---|---|---|
| [illegible] |  | [illegible] | TLC, MeOH aq | 31–[illegible] | [illegible] |
| [illegible] |  | 3-Chloroaniline (I) | Dist., MeOH aq | 16–17 | [illegible] |
| [illegible] |  | | MeOH | 77–78 | [illegible] |
| Dichloro-biphenyls 2 |  | 2-Chloroiodobenzene (V) | TLC, MeOH | 59–60 | [illegible] |
| [illegible] |  | 3,3'-Dinitrobiphenyl (II) ↓ 3,3'-Diaminobiphenyl | TLC, EtOH aq | 26–27 | 15 |
| [illegible] |  | — | MeOH | 148–149 | [illegible] |
| [illegible] |  | 2,4-Dichloroaniline (I) | TLC, MeOH aq | 24–25 | [illegible] |
| [illegible] |  | 3,4-Dichloroaniline (III) | TLC, MeOH | 43–46 | |
| [illegible] |  | 2,5-Dichloroaniline (II) | Col., MeOH aq | 33–[illegible] | |
| [illegible] |  | 2,4,6-Dichloroaniline (I) | Col., TLC, EtOH aq | [illegible] | |



[illegible]

| [illegible] | Formula | Start of material (Source) | Method of purification | Physical data m.p. (°C) | [illegible] |
|---|---|---|---|---|---|
| 4,4'- |  | 2,4-Dichloro-2-nitrobiphenyl[illegible] ↓ 2-Amino-4,4'-dichlorobiphenyl | MeOH aq | 57–58 | 51 |
| [illegible]chloro-biphenyls 3,6',4'- |  | 2,2-Dichlorobenzidine[illegible] or 2,4-Dichloroiodobenzene[illegible] | Dist., Col., EtOH aq | 41 | 58 |
| 3,3',4'- |  | 3,4,3',4'-Tetraaminobiphenyl (VI) or 3,3'-Dichlorobenzidine (VIII) | TLC, EtOH aq | 173 | 41 |
| 3,3',5'- |  | 3,5,3',5'-Tetrachlorobenzidine[illegible] | TLC, MeOH, hexane | 164 | 65 |
| 2,6',6'- |  | 2,6-Dichloroaniline (I) ↓ 2,6-Dichloroiodobenzene | Col., EtOH aq | 198 | 40 |
| 3,4,5- |  | 2,3,4,5-Tetrachloronitrobenzene (IV) ↓ 2,3,4,5-Tetrachloroaniline | Col., MeOH | 91 | 48 |
| 3,5,6- |  | 2,3,5,6-Tetrachloronitrobenzene (II) ↓ 2,3,5,6-Tetrachloroaniline | Col., hexane | 76 | 50 |
| [illegible]tachloro-biphenyl 3,4,5,6- |  | Pentachloronitrobenzene (I) ↓ Pentachloroaniline | Col., MeOH | 123 | 51 |
| [illegible]achloro-biphenyls 3,4,6,2',4',6'- |  | 2,6,2',6'-Tetrachlorobenzidine[illegible] or 2,4,6-Trichloroiodobenzene[illegible] | Col., MeOH, EtOH | 112–113 | 60 |

[illegible]

MONS 202116




Fig. 16. Gas chromatograms of (a) a standard pesticide mixture (b) Aroclor 1254 and (c = b) a 50–50 mixture of both. The concentration of Aroclor 1254 is approximately ten times those of the pesticides in (a). Gas chromatograph (Packard 7620) conditions: nitrogen gas flow 80 ml/min; temperatures—oven 200°, inlet 230°, detector 218°; $^{63}$Ni electron capture detector; 5 ft × 1/4 in glass column packed with 5% SE-30 on 80–90 mesh Chromosorb Q.

cause interference with pesticide residue analysis under these or similar operating conditions.

The GLC was carried out using a Varian Model 1200 gas chromatograph fitted with a tritium electron capture detector and a spiral glass column (6 ft × 1/8 in O.D.) packed with 6% QF-1 and 4% SE-30 on acid-washed Chromosorb W. (The number of theoretical plates for DDT = 2227.) The operating conditions were: column temperature 200°, injector temperature 250° and detector base temperature 250°, nitrogen flow rate 20–30 ml/min, Varian Aerograph Model 20 recorder of 1 mV full-scale deflection and a chart speed of 2/3 in./min.

It is of note that the peaks of the commonly found pesticides all have a corresponding PCB peak that would interfere if present in the same extract. The problem of the interferences of PCBs with the analysis of chlorinated pesticides has been largely overcome by prior column chromatography[8,19] or via an initial extraction with hexane as in regular pesticide analyses, partitioning with acetonitrile to remove fats, passage of cleaned-up extract through both Florisil and silica acid columns, then finally analysis by GLC with preferably a chloride specific detector. The pesticides are retained on the silica acid column for later elution and identification.

The use of a column (30 cm × 2.5 cm O.D.) packed with 40 ml (ca. 30 g or 10 cm in height) Florisil (60–80 mesh) stored at 130° until used and eluted with 200 ml of hexane permitted the separation of the PCBs and organochlorine pesticides

MONS 202117




Fig. 1. (a) Interference with organochlorine pesticide residue analysis on a florisil column (60 ft × $\frac{1}{4}$ in. packed with 4% SE-30 and 6% OV-1 on 60–80 mesh Chromosorb W (a) Standard mixture of organochlorine pesticides 1 = 0.05 ng of lindane, 2 = 0.10 ng of heptachlor, 3 = 0.20 ng of aldrin, 4 = 0.14 ng of heptachlor epoxide, 5 = 0.20 ng of DDE, 6 = 0.20 ng of dieldrin, 7 = 0.32 ng of DDD, 8 = 0.50 ng of DDT. (b) 5 μg of Aroclor 1254 (the fourteen major peaks being numbered I to XIV). (c) Combination of the above organochlorine standard pesticide mixture and Aroclor 1254. Injector temp. 250°, column temp., 200°, detector base temp. 250°, nitrogen flow rate 20–30 cc/min.

TABLE 9

PERCENT RECOVERIES OF PCB'S AND PESTICIDES FROM FLORISIL COLUMNS BY ELUTION WITH HEXANE[11]

Recoveries are based on peak height comparisons and each value represents the average of duplicate determinations

| PCB peak No. (ref. C) | Expt. I 100 ml of hexane | Expt. II 200 ml of hexane | Pesticide* peak | Expt. III 100 ml of hexane | Expt. IV, 200 ml of hexane |
|---|---|---|---|---|---|
| 1 | 50.1 | 92.2 | Lindane | None | None |
| 2 | 86.7 | 103.1 | Heptachlor | None | 92.7 |
| 3 | 65.0 | 100.0 | Aldrin | 62.8 | 94.1 |
| 4 | 98.2 | 101.0 | Hept. epox. | None | None |
| 5 | 42.1 | 100.0 | DDE | 20.5 | 97.5 |
| 6 | 44.0 | 98.7 | Dieldrin | None | None |
| 7 | 62.9 | 100.— | DDD | None | None |
| 8 | 96.8 | 105.2 | p,p′ DDT | None | None |
| 9 | 66.4 | 103.8 | | | |
| 10 | 76.6 | 104.8 | | | |
| 11 | 76.9 | 99.0 | | | |
| 12 | 57.— | 100.0 | | | |
| 13 | 100.— | 100.0 | | | |
| 14 | 71.1 | 100.0 | | | |

* Under the experimental conditions used, a total of 20% ether in hexane is required to elute the pesticides although some can appropriately remove them with the exception of DDE, aldrin and heptachlor)[13]. Table 9 shows percent recoveries of PCB's and pesticides from Florisil columns by elution with hexane and indicates almost quantitative removal of the PCB's was effected with 200 ml of hexane while under the same conditions only three of the eight pesticides tried showed evidence of elution. Further experiments indicated the feasibility of column separation of PCB's and pesticides when mixed and when they were present in the extract from an animal tissue (Table 10).

After the Florisil separation of PCB's and pesticides has been effected the two eluates are chromatographed separately on an SE-30/QF-1 column and compared with appropriate standards for quantitation; this is then normally followed by confirmation of the identities of the pesticides.

Despite the separation of the PCB's and their elimination as sources of interference it is still necessary to confirm the identity of the pesticides by additional techniques because of the non-specific nature of the electron capture detector. The sequence recommended by REYNOLDS[14] involves (a) the analysis of the pesticide fraction on a more polar liquid phase column, e.g., polyester such as DEGS or DEGA



TABLE 10

PERCENT RECOVERIES OF PCB'S AND PESTICIDES FROM A MIXTURE AFTER SEPARATION ON FLORISIL (EXPTS. V AND VI)[11]

The peaks are arranged in order of their appearance (increasing retention time) from the GLC column and where a PCB and a pesticide appear in the same line (horizontally) they have similar retention times.

| Eluted with 200 ml of hexane | | | Eluted with 250 ml of 20% ether in hexane | | |
|---|---|---|---|---|---|
| PCB and/or pesticide peak | % Recovery[a] | % Recovery[b] | Pesticide peak | % Recovery[a] | % Recovery[b] |
| | | | Lindane | 93.6 | 98.3 |
| Heptachlor | 92.7 | 98.1 | Heptachlor | None | None |
| CB1 | 104.0 | 101.7 | | | |
| CB2 + Aldr | 105.1 | 96.6 | Aldrin | 7.3 | 4.0 |
| CB3 | 100.0 | 106.0 | | | |
| CB4 | 104.2 | 102.6 | Heptachlor epoxide | 96.4 | 102.2 |
| CB5 + DDE[c] | 97.9 | 102.4 | DDE | None | None |
| CB6 | 101.3 | 100.0 | | | |
| | | | Dieldrin | 100.0 | 100.0 |
| CB7 | 97.8 | 103.1 | | | |
| CB8 | 100.0 | 100.0 | DDD | 102.3 | 95.0 |
| CB9 | 91.6 | 97.0 | | | |
| CB10 | 101.7 | 104.3 | DDT | 99.8 | 95.3 |
| CB11 | 100.0 | 105.5 | | | |
| CB12 | 100.0 | 100.0 | | | |
| CB13 | 100.0 | 100.0 | | | |
| CB14 | 96.4 | 100.0 | | | |

[a] A standard mixture of PCB's and pesticides in pure hexane was placed on the Florisil column; first elution was made with 200 ml of hexane, receiver was changed and the column eluted with 250 ml of 20% ethyl ether in hexane.

[b] Same as in [a] except that the PCB's and pesticides were first mixed with an extract from an animal tissue which was known to be essentially free of pesticides.

[c] Since a single peak was obtained, the recovery was calculated by a comparison of the peak height against that in the combined standard mixture of PCB and pesticide; for all other cases the peak height was compared to that in the standard injected separately.




Fig. 18. Separation of PCB's and organochlorine pesticides on a polar phase column (5% DEGS/2[illegible]). (a) Standard mixture of organochlorine pesticides of the same amounts as in Fig. 17. 1 = heptachlor and aldrin; 2 = lindane; 3 = heptachlor epoxide; 4 = DDE; 5 = dieldrin; 6 = DDT; 7 = DDD. (b) 5 ng of Aroclor 1254. (The fourteen major peaks are numbered I to XIV as in chromatogram b of Fig. 17, assuming that the order of elution is unchanged.) (c) Combination of the organochlorine standard pesticide mixture and Aroclor 1254. Other GLC parameters as in Fig. 17.




Fig. 19. Comparison of Aroclors 1254 and 1260. (a) 5 ng of Aroclor 1254; the fourteen major peaks are numbered I to XIV as in chromatogram (b) of Fig. 17. (b) 5 ng of Aroclor 1260; the seventeen major peaks are numbered I to XVII (the early peaks corresponding to those in Aroclor 1254). GLC parameters as in Fig. 17.



[illegible] (b) derivatization and use of characteristic GLC [illegible] types. For example, reaction with ethanolic potassium hydroxide yields products (via mainly dehydrochlorination) with shorter retention times. This technique is most effective for DDT, DDD and their isomers as well as α- and γ-BHC (the β-isomer is unaffected).

Typical GLC separation of PCB's, pesticides and a mixture of the two groups is shown in Fig. 18.

The two most commonly found PCB's in wildlife samples resemble Aroclor 1254 and 1260. Certain differences are exhibited in their GLC patterns that can be used to distinguish one from the other (Fig. 19). Aroclor 1260, with a higher chlorine content, shows about seventeen major peaks on an SE-30/QF-1 column compared to eleven with Aroclor 1254. Another major difference is the peak height ratio of peaks 10 and 13, e.g., this ratio is approximately 9.3 and 1.0 for Aroclors 1254 and 1260, respectively.

Fig. 20 illustrates GLC chromatograms of a number of Aroclor mixtures determined on SE-30/QF-1 columns. The more highly chlorinated biphenyls (Aroclors 1254 and 1260) are easily detected while the compounds of the lower chlorinated mixtures, e.g., Aroclors 1221, 1232 and 1242 and the higher-molecular-weight mix-




Fig. 20. GLC profiles of the more popular Aroclor mixtures under normal analytical conditions. All peak numbers correspond to those of Aroclor 1254. (a) 5 ng of Aroclor 1221; (b) 5 ng of Aroclor 1232; (c) 5 ng of Aroclor 1242; (d) 5 ng of Aroclor 1254; (e) 5 ng of Aroclor 1260; (f) 5 ng of Aroclor 5460. GLC parameters as in Fig. 17.



MONS 202119




[illegible] the presence of PCB's of the Aroclor 1260 type in a duckling [illegible] of Aroclor 1260 numbered I to XVII as in chromatogram b [illegible] equivalent of 0.35 mg of duck sample (hexane portion of Florisil split). Note the [illegible] on to PCB [illegible] GLC parameters as in Fig. 17.

[illegible] (Aroclor 5460) are less responsive with the usual operating parameters and are thus more [illegible] to go undetected.

Fig. 22 illustrates gas chromatograms of Aroclor 1260 found in a duckling and [illegible] 22 indicates a resin powder extract from a fish hatchery trough indicating the presence of PCB's of the Aroclor 1254 type and illustrates an example of PCB's [illegible] contamination as a direct result of industrial application.

The separation of PCB's from organophosphorus compounds by the Florisil technique has also been affected[12]. GLC utilizing a dual detector (electron capture [illegible] a phosphate or thermionic) connected to a dual pen recorder permits the differentiation of the PCB's and organophosphates as shown in Fig. 23. The PCB's




[illegible] Resin powder extract indicating the presence of PCB's of the Aroclor 1254 type (a) 5 ng of Aroclor 1254 numbered I to XIV as in chromatogram (b) of Fig. 17. (b) The equivalent of [illegible] mg of resin powder extract (hexane portion of Florisil split) numbered I to XIV as in Aroclor 1254. GLC parameters as in Fig. 17.






Fig. 23. Differentiation of Aroclor 1254 and organophosphorus pesticides using dual detectors and effluent split. (a) Organophosphorus standard pesticide mixture: 1 = 1.75 ng of phorate; 2 = 2.0 ng of diazinon; 3 = 3.0 ng Ronnel; 4 = 13.0 ng of malathion; 5 = 5.0 ng of parathion; 6 = 15.0 ng of methyl Trithion; and 7 = 10.0 ng of ethion. (b) Combination of organophosphorus mixture (chromatogram a) and 10 ng of Aroclor 1254 numbered I to XIV as in chromatogram b of Fig. 17 (a compromise of the EC response is necessary to obtain maximum response on the phosphorus detector). P = Phosphorus detector; EC = electron capture detector. Column 4% OV-100/6% OV-210 on 60–80 mesh acid-washed Chromosorb W. Other GLC parameters as in Fig. 17.

similarly to the organochlorine pesticides are not detected on the phosphate detector.

The estimation of PCB's following GLC analysis has been generally achieved via the following techniques: (1) the PCB estimation based on the peak of Phenoclor DP6 having $R_x$ (relative retention time with dieldrin = 1) equal to 1.45 (ref. 2), (2) the PCB's are estimated on the assumption that they have similar electron capture responses to *p,p'*-DDE and a factor is applied to fit the assumed 54% chlorine content of the PCB's[54], (3) estimates of the PCB's are reported as the *sum* of all the PCB components[13], and (4) estimation of PCB's based on an average of two peaks[2].

A GLC study[45] of Aroclors 1221, 1232, 1254, 1260, 1262, 4465, 5442 and 5460 on columns of 10% DC-200 and 15% QF-1/10% DC-200 with electron capture detection showed the Aroclors to be multicomponent mixtures with retention times throughout and beyond the retention time range of common chlorinated pesticides. With a detector sensitivity producing ½ full scale recorder response to 1 ng of heptachlor epoxide, significant responses were obtained with about 10–20 ng of the various Aroclors.

In the above study of ARMOUR AND BURKE[46] the efficacy of silicic acid columns for the separation of Aroclors and chlorinated pesticides added to trout and salmon extracts was also elaborated. Figs. 24 and 25 illustrate gas chromatograms of the mixtures of Aroclors and DDT analogs in trout and salmon, respectively, before and after column separation. Determinations by electron capture GLC showed the salmon



MONS 202120




[illegible] GLC curves of [illegible] extract [illegible] with 2.0 ppm Aroclor 1254, 0.1 ppm [illegible]-DDT, 0.15 ppm [illegible]-DDE and 0.3 ppm [illegible]-DDT. (a) Before separation on silicic acid column containing (b) petroleum ether eluate for [illegible] and (c) benzene eluate for [illegible] separation. (b) petroleum ether eluate from silicic acid column containing (c) [illegible]-DDT, (d) [illegible]-DDE and (e) [illegible]-DDT. 0.8 g sample injected for each curve.

[illegible] GLC curves of Lake Michigan [illegible] extract containing residues of PCB and DDT and [illegible] before and after separation on silicic acid column. (a) Before separation. (b) petroleum ether eluate for silicic acid column containing 2.0 ppm PCB (as Aroclor 1254). (c) benzene eluate for [illegible] containing (1) 0.1 ppm [illegible]-DDE, (2) 0.4 ppm [illegible]-DDT, (3) 0.1 ppm [illegible]-DDT and (4) 0.2 ppm [illegible]-DDE. 2.7 mg of sample injected for curves a, b and c, while 0.8 g of sample injected for curve c, used for [illegible]-DDE quantitation.

to contain 14.6 ppm PCB, 3.2 ppm [illegible]-DDT, 1.3 ppm [illegible]-DDT, 0.4 ppm [illegible]-DDE and 0.4 ppm [illegible]-DDE. The PCB residue was measured in terms of Aroclor 1254 using total area of the GLC response.

Hoffman *et al.*[7] described the detection of PCB's in British wildlife. In birds' livers and eggs they are often in greater quantities than organochlorine pesticide residues. GLC of keste[illegible] liver was achieved utilizing three columns, viz., SE-52, Apiezon L and XE-60. The actual retention times recorded for the peaks on these columns are shown in Table II. Fig. 26 shows a chromatogram of an extract from kestrel 1, indicating the peaks corresponding to [illegible]-DDE (0.43 ng), [illegible]-BHC (0.3 ng), PCB (0.03 ng). The other peaks form a series from PCB's. Fig. 26 also shows the analogous chromatogram of a commercial PCB resin and illustrates the similarity of peaks of high retention times. The peaks on the chromatogram for the kestrel liver extract in Fig. 26 were considered to be equivalent to about 12 ng of polychlorobiphenyl compounds. This is contrasted against the organochlorine pesticide residues which were found to be 0.3 ng of [illegible]-BHC, an isomer of BHC which is [illegible] insoluble in water. Further confirmation of the components of [illegible]



TABLE II

RELATIVE RETENTION TIMES ON THREE DIFFERENT GLC COLUMNS FOR SOME CHLORINATED PESTICIDES[a]

KLE = kestrel liver extract; PCB = polychlorobiphenyl resin

| *Silicone Gas SE-52 column* | | *Apiezon L column* | | *Cyanosilicone GE/XE-60 column* | |
|---|---|---|---|---|---|
| *KLE* | *PCB* | *KLE* | *PCB* | *KLE* | *PCB* |
| 1.42 | 1.42 | 2.31 | 2.30 | 1.24 | 1.21 |
| 1.73 | 1.73 | 2.67 | 2.67 | 1.34 | 1.33 |
| 1.91 | 1.90 | 3.04 | 3.05 | 2.20 | 2.19 |
| 2.82 | 2.81 | 5.06 | 5.06 | 2.62 | 2.60 |
| 3.50 | 3.58 | 5.96 | 5.93 | 2.07 | 2.05 |
| 3.71 | 3.70 | 6.57 | 6.56 | 3.69 | 3.50 |
| 4.68 | 4.68 | | | 4.63 | 4.63 |
| 5.82 | 5.62 | | | 5.11 | 5.11 |
| 7.10 | 7.11 | | | | |

[a] Dieldrin = 1.00

using 1% acetone in hexane as developing solvent. Spots corresponding to the PCB's are found at $R_F$ values between 0.8 and 1.0, whereas the commonly occurring organochlorine pesticides and their metabolites produce spots of lower $R_F$ values.

A similar separation can also be shown with reversed-phase paper chromatography[18], whereby the unknown compounds (PCB's) have very much lower $R_F$ values compared with the chlorinated pesticides and their metabolites.

HOLDEN AND MARSDEN[19] reported that seals and porpoises in Scotland and




Fig. 26. GLC liquid-phase chromatogram of (a) an extract of kestrel liver and (b) a commercial polychlor[illegible]

MONS 202121

[illegible] porpoises and [illegible]

[illegible] GLC on a Varian [illegible] following two glass columns 5 ft × 1.8 in. O.D. One [illegible] DC [illegible] washed DMCS-treated SO [illegible] and the other [illegible] coated with 5% DC-200 [illegible] temperature was 200° [illegible] gas flow 50 ml/min and [illegible] solution for dieldrin [illegible] to 2600 theoretical plates. While [illegible] of the common [illegible] residues was possible on one or the other of these [illegible] PCB residues with p,p'-TDE and p,p'-DDT necessitated [illegible] (alcoholic potassium hydroxide hydrolysis) which [illegible] converts p,p'-DDE to p,p'-DDE and p,p'-TDE to p,p'-MDE. [illegible] PCB's are unaffected. A [illegible] p,p'-DDT and p,p'-TDE from PCB [illegible] confirming the latter.

Unidentified peaks on the chromatograms appeared to be similar to those produced by PCB's and to enable comparison to be made, the ratios ($R_x$) of the retention times of [illegible] peak (relative to dieldrin = 100) were calculated for [illegible] temperature of 200° (these values being temperature dependent). The $R_x$ values of PCB compounds from commercial PCB formulations were also de-

TABLE 12

RELATIVE RETENTION VALUES ($R_x$) OF RESIDUES[a]

| DC-200 column | | | DC-200 [illegible] column | | |
|---|---|---|---|---|---|
| Residue | | Sample | Residue | | Sample |
| PCB | = 68.5 | 68 | PCB | = 68.5 | 70 |
| p,p'-DDE | 100 | 100 | p,p'-DDE | = 84 | 83.5 |
| Dieldrin | | | Dieldrin | 100 | 100 |
| PCB | 121 | 125 | p,p'-TDE | = 121 | 121 |
| p,p'-TDE | = 129 | 128 | PCB | = 123 | 122 |
| PCB | = 148 | 149 | PCB | = 146.5 | — |
| p,p'-DDT | = 170 | 171 | p,p'-DDT | 147 | 147 |
| PCB | = 174 | 173 | PCB | 241 | 241 |
| PCB | 205 | 204 | PCB | 282 | 278 |
| PCB | = 282 | 287 | | | |
| PCB | 330 | 338 | | | |

[a] Dieldrin = 100

[illegible] both types of column (Table 12). At least seven PCB-type peaks were [illegible] including those which [illegible] with p,p'-TDE and p,p'-DDT.

The GLC analysis of [illegible] and heavy metal residues in [illegible] was described by [illegible] et al.[illegible] eggs collected in 1968 from [illegible] in Wisconsin, Maine and Florida all contained residues of DDE, DDD, dieldrin



TABLE 13

DDT[a] AND PCB RESIDUES IN MARINE BIRDS[b] AND IN THE PEREGRINE FALCON[c] (refs. 9)

| Species, locality, date | Total DDT[d] (p.p.m.) | % DDE | PCB (p.p.m.) | DDT/PCB |
|---|---|---|---|---|
| Cassin's auklet[e] | 5.8 | 98 | 0.16 | 36 |
| Ancient murrelet[f] | 0.75 | 90 | 0.15 | 5 |
| Fulmar[g] | 0.41 | 76 | 0.08 | 5 |
| Fulmar[g] | 3.4 | 89 | 0.34 | 10 |
| Red phalarope[h] | 0.78 | 70 | 0.10 | 8 |
| Rhinoceros auklet[i] | 2.7 | 97 | 0.36 | 8 |
| Slender billed shearwater[j] | 32.0 | 92 | 2.1 | 15 |
| Sooty shearwater[k] | 12.3 | 94 | 1.2 | 10 |
| Sooty shearwater[k] | 10.1 | 86 | 0.0 | 12 |
| Peregrine falcon | | | | |
| Breast muscle, second year female, migrant from Arctic | 104 | 99 | 22 | 4.5 |
| Breast muscle, immature California | 13 | 90 | 10.5 | 1.2 |
| Breast muscle, adult female California | 112 | 98 | 100 | 1.0 |

[a] From RISEBROUGH et al.[illegible] and RISEBROUGH et al.[illegible]
[b] Entire bird analyzed except peregrine falcons
[c] From RISEBROUGH et al.[illegible]
[d] Includes p,p'-DDT, p,p'-DDD, p,p'-DDE, p,p'-DDMU, p,p'-DDE, p.p.m., wet weight
[e] *Ptychoramphus aleuticus*, adult female, Farallon Islands, April, 1966
[f] *Synthliboramphus antiquus*, Monterey Bay, Nov. 1, 1966
[g] *Fulmarus glacialis*, Monterey Bay, Nov. 1, 1966
[h] *Phalaropus fulicarius*, Monterey Bay, Nov. 1, 1966
[i] *Cerorhinca monocerata*, Monterey Bay, Nov. 1, 1966
[j] *Puffinus tenuirostris*, Monterey Bay, Dec. 12, 1966
[k] *Puffinus griseus*, Monterey Bay, Nov. 1, 1966

of chlorine per molecule). Many also contained traces of DDT. Fractions were analyzed on OV-17/3% XE-60 according to REICHEL et al.[illegible]. Confirmatory residue analyses were performed on a 12% DEGS on Anakrom SO (100–110 mesh) column with a temperature of 100° and a nitrogen flow rate of 85 ml/min (the retention time of dieldrin was 9.5 min).

The widespread distribution of PCB's and chlorinated hydrocarbons in marine ecosystems of the Pacific Ocean has been reported by RISEBROUGH and co-workers[illegible]. Birds contain higher concentrations of these chemicals than fish, e.g., tissues of the peregrine falcon have contained the highest amounts which have so far been recorded. The PCB's have thus far not been detected in samples of airborne particulates, but their observed distribution in the sea indicates that they are dispersed by wind currents and that their fallout pattern is similar to that of the DDT compounds. Tables 13 and 14 depict the DDT and PCB residues in marine birds and in peregrin falcon and in the eggs of several bird species, respectively. Table 15 shows the DDT and PCB residues in marine fish. Of the commercial PCB preparations, those having the following retention times relative to p,p'-DDE on DC-200 and QF-1 columns have been detected in marine fish and birds.

DC-200: 1.25, 1.48, 1.75, 2.05, 2.11, 2.50, 2.00, 3.11, 3.88, 5.53

MONS 202122

[illegible]

[illegible] DDT [illegible]

[illegible] indicated. Similar values [illegible] have been [illegible] extracts of seals from the North [illegible]

[illegible] AND PCB CONTENTS IN FATS OF SEA-RANGE BIRDS[illegible]

| [illegible] | No. | Total DDTs (μg.) | p,p'-DDT (%) | PCBs (μg.) | DDT/PCB |
|---|---|---|---|---|---|
| [illegible] | | | | | |
| [illegible] | | | | | |
| [illegible] | 17 | 226 | 91 | 213 | 2.9 |
| [illegible] | | | | | |
| [illegible] | | | | | |
| [illegible] | 2 | 1.3 (1.3–1.3) | 99 | 62 | 2.1 |
| [illegible] | | | | | |
| [illegible] | 5 | 1945 (611–3611) | 96 | 559 (364–1019) | 3.5 |
| [illegible] | | | | | |
| [illegible] | 1 | 110 | 93 | 20 | 5.5 |
| [illegible] | 1 | 195 | 91 | 62 | 1.7 |
| [illegible] | | | | | |
| [illegible] | | 1.7 (1.7–1.7) | 97 | 15 | 10 |
| [illegible] | | | | | |
| [illegible] | 1 | 121 | 95 | 115 | 1.6 |
| [illegible] | 1 | 215 | 92 | 1.2 | 2.1 |
| [illegible] | 1 | 2.3 | 57 | 150 | 0.95 |
| [illegible] | | | | | |
| [illegible] | 1 | 52 | 59 | 130 | 1.6 |
| [illegible] | 1 | 569 | 98 | 22 | 36 |
| [illegible] | | | | | |
| [illegible] | 2 | 1260 (1.15–1.35) | 89 | 505 (460–550) | 1.7 (1.3–2.0) |
| [illegible] | 5 | 1110 | 88 | 10.0 (550–1600) | 1.4 |
| [illegible] | | | | | |
| [illegible] | 2 | 1005 (5.5–5.5) | 89 | 181 (131–132) | 5.5 |
| [illegible] | | | | | |
| [illegible] | 2 | 10 (3–17) | 82 | 3.1 (2.2–3.9) | 3.0 |
| [illegible] | | | | | |
| [illegible] | 1 | 2.5 | 87 | 371 | 1.0 |

[illegible]

[illegible] from [illegible] PCB [illegible] [illegible] not calculated from 61–62 [illegible]

[illegible]



[illegible] concentrations of chlorinated hydrocarbons [illegible] both DDT and PCB [illegible] to [illegible] in order of magnitude higher in marine birds than in fish (Tables 13 and 15). As shown in Tables 13 and 14, PCB is widely distributed among marine birds which are [illegible] terminal carnivores of a complex mesh of food chains in the sea.

The chlorine content of extracts of oilier birds and fish was determined with

TABLE 15

DDT[a] AND PCB RESIDUES IN MARINE FISH[a]

| Species, locality, date | No. | Mean wt. (g) | Total DDT (p.p.m.) | p,p'-DDT (%) | PCB (p.p.m.) | DDT/PCB |
|---|---|---|---|---|---|---|
| Northern anchovy | | | | | | |
| Terminal Island | 22 | 11.7 | 14.0 | 93 | 2.0 | 7.1 |
| June 25, 1965 | | (5.5–20) | ±1.9 | | | |
| Shiner perch | | | | | | |
| San Francisco Bay | 14 | 5.5 | 1.0 | 29 | 1.2 | 0.9 |
| October 29, 1965 | | (3–8) | ±0.1 | | | |
| San Francisco Bay | 10 | 28.7 | 1.4 | 35 | 0.4 | 3.5 |
| October 20, 1965 | | (10–28) | ±0.3 | | | |
| San Francisco Bay | 15 | 15.3 | 1.1 | 33 | 1.2 | 0.9 |
| November 4, 1965 | | (8–20) | ±0.1 | | | |
| English sole | | | | | | |
| San Francisco Bay | 18 | 24.3 | 0.55 | 25 | 0.11 | 5 |
| July 29, 1965 | | (5–55) | ±0.07 | | | |
| San Francisco Bay | 33 | 17.3 | 0.55 | 24 | 0.11 | 5 |
| November 4, 1965 | | (7.5–53) | ±0.12 | | | |
| San Francisco lightship | 15 | 253 | 0.19 | 63 | 0.05 | 4 |
| December 1, 1965 | | (175–306) | ±0.02 | | | |
| Monterey | 15 | 195 | 0.76 | 70 | 0.04 | 19 |
| February 15, 1966 | | (89–262) | ±0.16 | | | |
| Jack mackerel | | | | | | |
| Channel Islands | 31 | 81.8 | 0.56 | 57 | 0.02 | 28 |
| November 22, 1965 | | (45–121) | ±0.10 | | | |
| Hake | | | | | | |
| Puget Sound | 22 | 281 | 0.18 | 23 | 0.16 | 1.1 |
| January 29, 1966 | | (185–350) | ±0.05 | | | |
| Channel Islands | 6 | 384 | 1.8 | 65 | 0.12 | 15 |
| February 22, 1966 | | (61–872) | ±1.1 | | | |
| Bluefin tuna | | | | | | |
| Body muscle | 7 | — | 0.36 | 45 | 0.04 | 14 |
| | | | ±0.22 | | | |
| Liver | 9 | — | 0.22 | 45 | 0.02 | 6 |
| | | | ±0.13 | | | |
| Yellowfin tuna | | | | | | |
| Liver | 13 | — | 0.07 | 13 | nil | > 7 |
| | | | ±0.02 | | | |
| Liver | 13 | — | 0.62 | 30 | 0.04 | 15 |
| | | | ±0.19 | | | |
| Skipjack tuna | | | | | | |
| Liver | 3 | — | 0.049[b] | 83 | 0.1 | 0.5 |
| Body muscle | 11 | — | 0.058 | 98 | nil | >30 |
| | | | ±0.014 | | | |
| Liver | 25 | — | 0.056 | 11 | nil | >30 |
| | | | ±0.021 | | | |
| Liver | 12 | — | 0.020 | 21 | nil | >20 |
| | | | ±0.005 | | | |

[a] From [illegible] *et al.*[illegible] concentrations on wet weight [illegible] means at indicated [illegible] 95% [illegible]

MONS 202123

TABLE 17

PEAK HEIGHTS OF ISOMERS AND HOMOLOGUES OF CHLORINATED BIPHENYLS

| Sample | $R_x$ | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | 69 | 81 | 100[a] | 151 | 153[b] | 171 | 200 | 234 | 283 | 333 | 379 |
| Raw sludge A | 36 | 63 | 52 | 104 | 100 | 92 | 52 | 19 | 21 | 10 | 5 |
| Raw sludge B | 39 | 24 | 17 | 31 | 100 | 60 | 29 | 53 | 84 | 29 | 25 |
| Aroclor 1254 | 30 | 72 | 59 | 119 | 100 | 92 | 25 | 12 | 09 | 8 | 4 |
| Aroclor 1260 | 3 | 14 | 4 | 30 | 100 | 60 | 29 | 55 | 97 | 30 | 24 |

[a] This peak includes *p,p'* DDE in sludge.
[b] Peak height = 100 for reference

The peak pattern for sludge A closely resembles a PCB formulation containing about 50% by weight of chlorine (Aroclor 1254) but the pattern for sludge B suggests Aroclor 1260. On the basis of wet weight of raw sludge the estimated concentrations of PCBs using Aroclor 1254 as reference in fifteen samples were in the range <0·1 to 14 p.p.m. In terms of dry matter the values ranged from 1 to 185 p.p.m. It was estimated that an average concentration of PCB of the order of 1 p.p.m. would be equivalent to a discharge in the Clyde estuary of the order of one ton of PCBs per year. PCB compounds found in wildlife, using GLC at about 200°, have been shown to contain mostly six to seven chlorine atoms[2] although compounds with more chlorine atoms may not be sufficiently volatile to be detectable at this temperature.

The chronic toxicity, uptake and retention of Aroclor 1254 in two estuarine fishes, pinfish (*Lagodon rhomboides*) and spot (*Leiostomus xanthurus*) was described by HANSEN *et al.*[51] The results suggested that chronic exposure to Aroclor 1254 increased susceptibility of the test fish to disease and also appeared to be toxic to these fish. This PCB is rapidly stored by pinfish and spot and persists in tissues for approximately three months. Chemical analyses for residues of Aroclor 1254 in pooled samples of fish tissue were performed by GLC following (a) elution and concentration from a Florisil column[63] with 6% ethyl ether in petroleum ether, and (b) extraction of samples of less than 1 g by a modification of the ENOS[64] micromethod. The column eluates were analyzed by electron capture gas chromatographs equipped with DC-200 columns. The multiple-peaked Aroclor 1254 was quantitated by averaging the heights of five major peaks having retention times relative to aldrin of 1·31, 1·55, 2·32, 2·74, and 3·27. Interference from DDT and its metabolites was negligible due to relatively high concentrations of Aroclor 1254 found in the samples.

Spot exposed to 1 p.p.b. Aroclor 1254 for fifty-six days rapidly stored this PCB with maximum levels being attained in fourteen to twenty eight days. Thereafter the relative amount (p.p.m.) was generally constant, whereas the absolute amount (*μg*) continued to increase as the fish grew. The liver concentrated the greatest relative amount of this PCB followed in decreasing order by the gills, whole fish, heart, brain, and muscle. Maximum concentrations in whole spot were 3·7 × tot times that in the test water. In earlier work with DDT[65] it was shown that when pinfish and Atlantic croaker (*Micropogon undulatus*) were exposed to 1 and 10 p.p.b. *p,p'* DDT in water the DDT content increased for two weeks, then remained constant at 10[4] to 3·8 × 10[4] times the exposure concentration. Table 17 lists the test conditions

MONS 202124

TABLE 17

[illegible] CONDITIONS, MORTALITY OF FISHES, AND UPTAKE OF AROCLOR 1254 IN FLOWING WATER BIOASSAYS

| Fish | SL[a] (mm) | Number[b] exposed | Concentration (p.p.b.) | Mortality 25% day | Mortality 50% day | Terminal percent | Days exposed | Aroclor 1254 (p.p.m.)[d] in fish after exposure | Water temperature (°C) | Sample (%) |
|---|---|---|---|---|---|---|---|---|---|---|
| Spot | 25 | 150 | 1 | | | 17 | 33 | 17 | 11–16 | 16–32 |
| Spot | 40 | 150 | 1 | | | 1 | 50 | 27 | 21–32 | 10–33 |
| Spot | 3 | 50 | 5 | 12 | 18 | 51 | 20 | 36 | 11–18 | 16–[illegible] |
| Spot | 4 | 100 | 5 | 21 | 46 | 53 | 26 | 120 | 8–10 | 20–[illegible] |
| Spot | [illegible] | 50 | 5 | 28 | 48 | 62 | 45 | 152 | 28–33 | 21–34 |
| Finfish | 10 | 50 | 5 | 9 | 12 | 66 | 14 | 11 | 16–22 | 20–3[illegible] |
| Finfish | 27 | 54 | 5 | 17 | — | 41 | 35 | 100 | 22–32 | 14–33 |

[a] SL = Sample length.
[b] An equal number of fish were held in control aquaria; control mortality never exceeded 7%.
[c] Mortality of spot exposed to 5 p.p.b. Aroclor was significantly greater in all tests ($p < 0.001$) than that of control fish ($\chi^2$ values were all greater than 10.84).
[d] Milligrams per kilogram.

TABLE 18

LEVELS OF AROCLOR 1254 IN p.p.m., WET WEIGHT, IN SPOT EXPOSED TO 1 p.p.b. OF THIS CHEMICAL (EACH SAMPLE CONSISTED OF TISSUES FROM TEN FISHES)

Total micrograms of Aroclor in parentheses

| Tissue | Number of days exposed | | | | | | | Number of days of flushing | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 0 | 3 | 7 | 14 | 28 | 42 | 56[a] | 14 | 28 | 42 | 56 | 70 | 84[a] |
| Brain | 0.1 | 3.6 | 6.0 | 8.6 | 8.4 | 12.0 | 8.3 | 7.6 | 5.8 | 5.0 | 3.5 | 3.4 | 2.9 |
| | (0.1) | (1.1) | (2.1) | (3.9) | (3.7) | (5.9) | (5.2) | (5.6) | (4.7) | (4.1) | (3.5) | (3.4) | (3.1) |
| Gills | 0.2 | 9.0 | 11.0 | 29.0 | 40.0 | 39.0 | 46.0 | 30.0 | 30.0 | 24.0 | 12.0 | 14.0 | 1[illegible].0 |
| | (0.1) | (8.3) | (13.0) | (41.0) | (64.0) | (82.0) | (74.0) | (75.0) | (72.0) | (65.0) | (35.0) | (46.0) | (37.0) |
| Heart | ND[b] (< 0.1) | 5.5 | 6.8 | 10.0 | 12.0 | 17.0 | 13.0 | 10.0 | 7.8 | 6.1 | 5.3 | 4.7 | 2.5 |
| | | (0.3) | (0.5) | (1.0) | (0.8) | (1.6) | (1.5) | (1.7) | (1.0) | (0.8) | (1.0) | (0.8) | (0.6) |
| Liver | 1.2 | 34.0 | 75.0 | 96.0 | 210.0 | 107.0 | 83.0 | 65.0 | 52.0 | 28.0 | 17.0 | 29.0 | 22.0 |
| | (0.1) | (12.0) | (42.0) | (92.0) | (126.0) | (86.0) | (72.0) | (83.0) | (60.0) | (38.0) | (30.0) | (58.0) | (35.0) |
| Muscle | 0.1 | 1.0 | 1.8 | 3.6 | 7.4 | 7.6 | 6.5 | 6.9 | 4.8 | 3.4 | 3.3 | 2.5 | 2.0 |
| | (0.1)[c] | (1.8) | (4.5) | (17.0) | (24.0) | (36.0) | (45.0) | (65.0) | (45.0) | (34.0) | (44.0) | (40.0) | (32.0) |
| Whole fish | 0.1 | 3.5 | 7.2 | 17.0 | 37.0 | 30.0 | 27.0 | 25.0 | 19.0 | 12.0 | 9.4 | 9.3 | 7.2[d] |
| | (1.5) | (72.0) | (201.0) | (661.0) | (1417.0) | (1553.0) | (1983.0) | (2452.0) | (2041.0) | (1373.0) | (1309.0) | (1486.0) | (78[illegible].0) |

[a] The same tissues analyzed from control spot contained no detectable Aroclor 1254 (< 0.1 to < 1.0 p.p.m. depending on sample weight).
[b] ND = not detected.
[c] Total micrograms in sample; sample approximately 26% of total amount of muscle.
[d] Other tissues analyzed on day 84 were: gall bladder 1.9 (0.9), gonad 0.9 (0.3), gut 2.5 (7.8), rest of muscle 4.5 (133.0), and skin 12.





MONS 202125

[illegible] Aroclor 1254 [illegible] Bioassay and Ta[illegible] [illegible] fish in s[illegible] [illegible] 1254.

[illegible] 1242 on growth [illegible]

[illegible] levels of the PCB [illegible] extracted from a dried [illegible] with acetone and analyz[illegible] [illegible] O.D. glass column packed [illegible] [illegible] XXX [illegible] O.D. glass [illegible] 5% OV-[illegible] 95% [illegible] Chromosorb XXX. The operating [illegible] 215 [illegible] operating 200° detector temper[illegible] [illegible] Identification of Aroclor 1242 was accomplished [illegible] relative retention times of the five major peaks along with relative [illegible] Quantification was accomplished by comparison of the area under the [illegible] with the standard Aroclor. Verification of the peak identi[illegible] [illegible] Aroclor 1242 as the major PCB component of the [illegible] Aroclor 1242 mixture were [illegible] GLC eluters were believed to be possible [illegible] of Aroclor 1242.

*Crassostrea clostrum* were found to absorb and concentrate Aroclor 1242 [illegible] to 100 times above the level in sea water. At 0.1 p.p.m. concentration in sea [illegible] PCB [illegible] levels of RNA and chlorophyll of this [illegible] diatom.

The occurrence of Aroclor 1254 in the water, sediment and biota of Escanaba [illegible] was described by Duke *et al.*[8]. Tissue of fish, crabs, oysters and shrimp [illegible] extracted for 4 h with petroleum ether in a Soxhlet apparatus following initial mixing with anhydrous sodium sulfate. Extracts were concentrated, partitioned with acetonitrile [illegible] acetonitrile then evaporated just to dryness, the residue trans[illegible] [illegible] with petroleum ether and the Aroclor 1254 eluted from [illegible] 6% [illegible] ether in petroleum ether.

Sediment samples were dried at room temperature and extracted for 4 h with [illegible] petroleum ether in a Soxhlet apparatus, extracts evaporated to dry[illegible] and the residues eluted from [illegible] as described above. The extracts [illegible] were identified and measured by electron capture GC using [illegible] of different polarity [illegible] DC-200, QF-1 and mixed DC-200 and QF-1 [illegible] identification. [illegible] few samples, TLC was employed for additional [illegible] and to assess the amount of DDT present. Interference from DDT was [illegible] due to the relatively high residues of Aroclor 1254 found in the samples. Qu[illegible] of this multiple peak compound was approximated by averaging the [illegible] peaks indicating recovery rates above 80%.

[illegible] studies [illegible] that [illegible] were the most sen[illegible] to Aroclor 1254 and were killed when exposed to 5.0 p.p.m. in flowing sea water [illegible] in water from [illegible] Bay even near the mouth of the river [illegible] less than 1 p.p.m. and [illegible] the bay contained a maximum [illegible] The above study [illegible] the need [illegible] both conducting continued [illegible] of estuaries in order to preserve these [illegible] grounds for valuable fish [illegible] tests on the effect of sublethal concen[illegible]



TABLE 19

ORGANOCHLORINE RESIDUES (ppb) FRESH WEIGHT IN BREAST MUSCLE OF ROBINS

| | µg of PCB [illegible] | p,p'-DDE | p,p'-DDT | PCB |
|---|---|---|---|---|
| Birds fed with PCB (Clophen A60) | 60 | 95 | 7 | 375 |
| | 55 | 96 | 25 | 167 |
| | 65 | 75 | 25 | 341 |
| | 55 | 64 | 21 | 164 |
| | 60 | 91 | 22 | 365 |
| | 55 | 56 | 2 | 287 |
| Control birds | — | 72 | 23 | 120 |
| | — | 69 | 27 | 70 |
| | — | 70 | 30 | 90 |
| | — | 66 | 26 | 51 |

tions of PCBs on estuarine organisms in sensitive stages of their life history.

The effect of PCB on the nocturnal activity in caged robins (*Erithacus rubecula* L.) was described by ULFSTRAND *et al.*[70].

The breast muscles of six birds fed a diet containing Clophen A50 were analyzed by GLC utilizing a Varian Aerograph 204 instrument equipped with electron capture detectors and three columns containing SF-96 (4%), QF-1 (8%) and SF-96/QF-1 (3:1) as the stationary phases on 100–120 mesh Gas-Chrom P. Table 19 depicts the organochlorine residues (*p,p'*-DDE, *p,p'*-DDT and PCB) found in breast muscles. The PCB level in the experimentally contaminated birds was four times that found in







Fig. 27. Gas chromatogram of an extract of human milk (heptachlor epoxide added as internal standard). 1 = PCB peak; 2, 3 = heptachlor epoxide; 4 = *p,p'*-DDE disturbed by PCB peaks; 4 and 5, 4 = PCB peak; 8, 5 = *p,p'*-DDT disturbed by PCB peak. (Cf. Fig. 30.)

Fig. 28. Thin-layer chromatograms of stannaros and extracts of pike. 1 = Dieldrin; 2 = heptachlor epoxide; 4 = *p,p'*-DDE; 4 = lindane; 5 = α-BHC; 6 = *p,p'*-DDE; 7 = *p,p'*-DDT; 8 = *p,p'*-DDE; heptachlor; 9 = aldrin; 10 = PCB.

MONS 202126




Fig. 31. Gas chromatograms performed for the analysis of PCB and chlorinated pesticides in a sample of Baltic herring (heptachlor epoxide added as internal standard). For standards *cf.* Figs. 29 and 30. (a) original extract. (b) extract of Area II, Fig. 28 before oxidation. (c) extract of area II, Fig. 28 after oxidation. (d) extract of area I, Fig. 28.

and carbon tetrachloride-treated rats. Hexane extracts of tissues were analyzed by GLC-electron capture analysis on a Varian Aerograph Model 600D gas chromatograph fitted with a coiled 4 ft × 1/4 in. O.D. glass column containing 4% SE-30 and 6% QF-1 on 80–100 mesh Chromosorb W. The nitrogen flow rate was 120 ml/min, with column and injection temperatures of 193° and 225°, respectively.

Residues were found in all tissues analyzed (blood, heart, kidney, brain, liver and fat), with the greatest concentration in the fat. The GLC-EC pattern of the residues was different from the standard mixture administered, indicating that all components were not metabolized at the same rate. Higher residues were found in the carbon tetrachloride-treated rats. Aroclor 1254 residues in the brain, liver, spleen, blood, testes, heart, kidney and fat were reduced by 90, 84, 80, 79, 78, 76, 64 and 33%,




Fig. 32. GLC traces from (a) 24 mg of Aroclor 1254 and (b) the residue found in the liver of a Group 2 rat (7 days).

MONS 202127

[illegible] STANDARD

| | Peak No. | | | | | | Peak No. | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 1 | 2 | 3 | 4 | 5 | 6 | 1 | 2 | 3 | 4 | 5 | 6 |
| | Group 1 | | | | | | Group 2 | | | | | |
| | 6 | 21 | 9 | 20 | 21 | 23 | 3 | 17 | 2 | 11 | 34 | 33 |
| | 3 | 15 | 9 | 7 | 25 | 41 | 1 | 19 | 4 | 11 | 35 | 30 |
| | 2 | 11 | 8 | 30 | 29 | 20 | 0 | 15 | 3 | 18 | 36 | 28 |
| | 5 | 21 | 11 | 21 | 19 | 23 | 0 | 15 | 2 | 14 | 34 | 35 |
| | 7 | 25 | 11 | 25 | 17 | 15 | 1 | 19 | 2 | 20 | 28 | 29 |
| | 8 | 23 | 5 | 22 | 30 | 23 | 2 | 19 | 4 | 12 | 34 | 29 |
| | 4 | 12 | 6 | 21 | 26 | 31 | 1 | 20 | 3 | 12 | 33 | 31 |
| | 1 | 7 | 6 | 21 | 26 | 39 | 1 | 17 | 2 | 8 | 31 | 40 |
| | Group 3 | | | | | | Group 4 | | | | | |
| | 7 | 25 | 14 | 19 | 18 | 17 | — | — | — | — | — | |
| | 9 | 23 | 12 | 19 | 19 | 19 | 1 | 20 | 4 | 12 | 31 | 33 |
| | 4 | 21 | 11 | 21 | 22 | 21 | 0 | 19 | 4 | 29 | 29 | 37 |
| | 8 | 22 | 15 | 15 | 15 | 15 | | 21 | 4 | 8 | 32 | 34 |
| | 10 | 29 | 16 | 21 | 13 | 12 | 1 | 29 | 5 | 20 | 26 | 25 |
| | 6 | 25 | 11 | 20 | 15 | 11 | 1 | 22 | 3 | 11 | 30 | 32 |
| | 5 | 22 | 13 | 21 | 15 | 13 | 2 | 22 | 3 | 13 | 29 | 31 |
| | 5 | 17 | 13 | 18 | 5 | 17 | 1 | 20 | 3 | 10 | 32 | 32 |

* The six major components are present in the following percentages: (1) 12% (2) 15% (3) 18% (4) 12% (5) 15% (6) 10% (6) 13%.

respectively, on twenty days. Aroclor 1254 significantly increased the size of the liver and also the percent lipid of the liver as well as to potentiate the toxicity of carbon tetrachloride in the rat.

Fig. 32 shows the chromatograms from 24 ng of Aroclor 1254 and from the residue found in the liver of a rat. The amount ($C_n$) that each of the six major peaks of the Aroclor contributes to the total residues in the tissues is presented in Table 20.

Fig. 32 and Table 20 show that the components of the Aroclor 1254 mixture with the shorter retention times (peaks 1, 2 and 3, and presumably with the lowest chlorine contents) were metabolized to a greater degree than those with the longer retention times. This observation agrees with that reported in studies with Phenoclor DP6 fed to Japanese quail.[3] Examination of the Aroclor 1254 residues in the tissues of rats fed in Diet 2 (Table 21) shows that the ratios determined on the longest day following the oral dose (before any new liver and fat (Group 1 vs. 2) and the ratios of the liver (liver and fat (Group 3 vs. 4). Table 22 depicts residues in tissues of rats fed, as after being orally administered a single dose of Aroclor 1254 (500 mg/kg).

A comparative toxicologic study with PCBs in chickens with special reference to [illegible] and tissue residues was described by Nosal and Kovacs.[4] Aroclor 1254, PCBs were used in a comparative toxicity test [illegible] chickens, e.g., Phenoclor DP6 (I), Clophen A60 (II) and Aroclor 1260 (III). Using [illegible] and subcutaneous



TABLE 21

CONCENTRATIONS OF THE AROCLOR 1254 RESIDUES IN TISSUES RELATIVE TO THOSE IN BLOOD

| | Group | | | |
|---|---|---|---|---|
| | 1 | 2 | 3 | 4 |
| Blood | 1.00 | 1.00 | 1.00 | 1.00 |
| Testes | 9.51 | 10.21 | 8.62 | 22.48 |
| Heart | 12.33 | 13.88 | 16.21 | 24.68 |
| Spleen | 14.58 | 13.86 | 9.51 | — |
| Kidney | 15.59 | 26.67 | 14.90 | 44.32 |
| Brain | 49.30 | 9.55 | 10.89 | 23.83 |
| Liver | 59.17 | 42.88 | 206.88 | 75.16 |
| Fat | 506.21 | 1601.57 | 233.89 | 4536.20 |

and centrolobular liver necrosis) as parameters, a significant difference in toxicity was found, i.e., I and II showed the highest and III the lowest. Microscopically centrolobular liver necrosis was found in chicks fed compounds I and II. Atrophy of the spleen was found in all test groups and chemical porphyrin was found as a general PCB effect: increased fecal excretion of coproporphyrin and protoporphyrin and fluorescence of tissues occurred in all test groups. This porphyrogenic effect was also observed in Japanese quail and rats treated with 2500 p.p.m. of compound I. GLC analysis of liver and brain of dead chicks gave PCB levels that varied from 120 to 2900 p.p.m. Tissues were extracted with petroleum ether in a Soxhlet extraction apparatus after drying with anhydrous sodium sulfate. Clean-up of the samples was performed by liquid–liquid partition with dimethylformamide and column chromatography using activated Florisil. The PCB preparations in the final extract were measured by GLC using a Varian Aerograph Model 204-1B with electron capture detection and a Pyrex glass column (5 ft. × 1/8 in.) filled with 10% DC-200 on 80–100 mesh Gas-Chrom Q operated at 200° with nitrogen as carrier gas at a flow rate of 50 ml/min. For quantitation one peak, considered to be representative, was selected (the peak with $R_r$ = 1.45 relative to dieldrin). The approximate amount of total PCB

TABLE 22

RESIDUES IN TISSUES OF RATS 45 DAYS AFTER BEING ORALLY ADMINISTERED A SINGLE DOSE OF AROCLOR 1254 (500 MG/KG)

Group 2 of Experiment II killed on day 47

| | Residue | | |
|---|---|---|---|
| | Wet tissue (p.p.m.) | | Relative to blood |
| Blood | 0.182 | ± 0.115* | 1.00 |
| Heart | 2.71 | ± 0.91 | 15.06 |
| Kidney | 1.40 | ± 0.29 | 18.51 |
| Brain | 1.39 | ± 0.41 | 21.25 |
| Liver | 19.01 | ± 1.71 | 50.11 |
| Fat | 197.40 | ± 31.50 | 1405.22 |

* Mean of seven values.

MONS 202128

TABLE 24

EXTRACTION METHOD FOR BIOLOGICAL MATERIALS




produced poisoning among the general population in Western Japan from summer to fall of 1968[6] with a total of 600 patients to date. It was found that contamination of the oil occurred by both the leakage of chlorobiphenyls into the rice bran oil through pin holes in a pipe used for heat exchange and deodorizing in the manufacturing process. It was also demonstrated that some newborn and still-born infants with darkish skin were born from the poisoned mothers.

Kojima *et al.*[7] described the utility of GLC for the analyses of chlorobiphenyls from the skin of a still-born infant who was born from a patient suffering from chlo-

MONS 202129




Fig. 34. Chromatograms of the skin of a still-born infant and control compared with chlorobiphenyls in rice oil ——, Skin of still-born infant — — — skin of infant (control) — · — · — chlorobiphenyls in rice oil




Fig. 33. Gas chromatograms of rice oil —— Rice oil (material) — — rice oil (control) — · — · — Kanechlore 300

two systems. One instrument was a Packard Model 7834 gas chromatograph equipped with a 150 mCi tritium affinity ionization detector operated at 50 V and a 1.83 m × 4.0 mm I.D. glass column packed with 5% SE-30 on 60–80 mesh acid-washed, DMCS-treated Chromosorb W (conditioned for seven days at 250° before use) with a column pressure of 25 p.s.i. of nitrogen (flow rate, 125 ml/min). The injection, oven and detector cell temperatures were 240°, 190°, and 215°, respectively. The second instrument was a Jarrell-Ash Model 28-700 gas chromatograph equipped with a 100 mCi tritium electron affinity ionization detector and operated at 20 V. A 1.22 m × 4.0 mm I.D. glass column contained a 1:1 mixture of 5% QF-1 and 5% DC 200 coated on 80–90 mesh Anakrom AS and conditioned for seven days at 250° before use (column pressure 16 p.s.i., nitrogen flow rate 100 ml/min). The injection port, oven and detector temperatures were 250°, 190° and 210°, respectively. For TLC,




Fig. 35. Chromatograms of the skin of infantile mice and control compared with chlorobiphenyls in rice oil — — Skin of infantile mice (material) — — skin of infantile mice (control) — · — · — chlorobiphenyls in rice oil

MONS 202130



Fig. 36. Chromatograms of extracts of mackerel muscle and content compared with chlorinated hydrocarbon standards. ——— = Skin of mackerel; – – – = skin of mackerel muscle extract; · · · · = chlorinated hydrocarbon standards.

Aluminum Oxide G plates were developed with n-heptane, sprayed with the Mercuric reagent[9] and exposed to UV light for 30 min.

Table 25 summarizes the GLC behavior of eleven Aroclor PCBs and a number of insecticides analyzed with two different columns in separate gas chromatographs. The observed retention times were compared with those obtained after the injection of aldrin, heptachlor, heptachlor epoxide, aldrin, dieldrin, DDT, DDE and TDE. The GLC of nearly all of the plasticizers resulted in chromatograms containing multitude of peaks, in some cases as many as fifteen.

Figs. 37 and 38 illustrate the TLC of plasticizers on Aluminum Oxide G. It

TABLE 25

COMPARISON OF PCBs AND CHLORINATED INSECTICIDES BY GLC

Chromatographic conditions: (1) Packed glass tube gas-liquid column 5% SE-30 on 60–80 mesh Chromosorb W, acid washed, DMCS treated; 6 ft × 4 mm I.D.; 200° (GLC-1). (2) Glass capillary column 300 ft × 0.01 in. I.D. coated with SE-30; 200° (GLC-2).

Abbreviations: BH = benzene hexachloride; HE = heptachlor epoxide; AL = aldrin; DE = dieldrin; DDE = *p,p'*-DDE; DD = DDD; TD = TDE.

| Pattern no. | Aroclor | % weight chlorine | GLC-1 | | GLC-2 | |
|---|---|---|---|---|---|---|
| | | | No. of peaks | Detected as | No. of peaks | Identified with |
| 1 | 1221* | 21 | 4 | None | 5 | DE |
| 2 | 1232 | 32 | 13 | BH, HE, AL, DE, DD, DE, TD, DE | 15 | BH, HE, DE, DD, DE |
| 3 | 1242 | 42 | 14 | BH, HE, AL, DE, DD, DE, TD | 14 | BH, HE, DE, DD, DE |
| 4 | 1248 | 48 | 15 | BH, HE, AL, DE, DD, DE, TD, DE | 15 | BH, HE, DE, DD, DE |
| 5 | 1254 | 54 | 11 | AL, DE, DD, DE, TD, DE | 10 | DE, DD, DE, TD |
| 6 | 1260 | 60 | 13 | TD, DE | 13 | DE, TD, DE |
| 7 | 1016 | 41 | 4 | None | 9 | None |
| 8 | 1254 + 1260 | 57 | 11 | TD | 14 | DE |
| 9 | 1242 + 1254 | 51 | 4 | None | 9 | BH, HE, DE |
| 10 | 1248 + 1260 | 54 | 9 | None | 9 | None |

* 1221 = Chlorinated diphenyl; 1232 = chlorinated diphenyl; 1016 = chlorinated diphenyl.




Fig. 37. Thin layer chromatogram of plasticizers A, B, C, D and E on Aluminum Oxide G. Solvent 1% acetone in n-heptane. STD = Insecticide standards for comparison.

Fig. 38. Thin layer chromatogram of plasticizers F, G, H, I, J and K on Aluminum Oxide G. Solvent 1% acetone in n-heptane. STD = Insecticide standards for comparison.

is likely that these PCBs were not fully separated or resolved in the above system since only one or two spots were found per plasticizer.

Polychlorinated biphenyls have been detected in all samples of rainwater obtained over a 12-month study in Great Britain from seven stations[73]. Hence, aerial fallout is one source of PCB in the sea. Aliphatic chlorinated hydrocarbons identified in surface waters of the North Atlantic are derived in part from barge dumping of waste products of the vinyl chloride industry[89].

PCB contamination of Escambia Bay, Fla., has been traced to an industrial plant on the Escambia river by Duke *et al.*[90]. PCBs have been found in sewage sludge in Sweden[91] and it has been estimated by Harvey[92] that approximately a ton of PCB a year is entering the Clyde estuary in Scotland as a component of raw sewage sludge and that an equivalent amount enters the estuary of the Thames in sewage sludge.

The input of PCBs into California coastal waters from urban sewage outfalls has been described by Seymour *et al.*[93].

Aliquots of concentrated extracts were analyzed by GLC using a Microtek 220 gas chromatograph equipped with a $^{63}$Ni electron capture detector and a 3% QF-1 column on 80–100 mesh, acid-washed, DMCS-treated Chromosorb. The column, injection port and detector temperatures were 190°, 230° and 250°, respectively. Nitrogen was used as carrier and purge gas, flow through the column was 68 ml/min

MONS 202131




Fig. 1. GLC analysis of sewage sludge.

TABLE 25

PCB AND DDT COMPOUNDS IN URBAN SEWAGE OUTFALLS IN CALIFORNIA

| Sampling station (discharge in m³/d)[a] | Sample No. | PCB type | Parts per billion | | Estimated kg/day[b] | |
|---|---|---|---|---|---|---|
| | | | PCB | DDT[c] | PCB | DDT |
| Richmond | 1 | ND | — | 0.010 | — | 0.005 |
| 21.5 | 2 | ND | — | 0.071 | | |
| | 3 | ND | — | 0.068 | | |
| | 4 | ND | — | 0.053 | | |
| El Cerrito | 1 | 1260 | 1.8 | 0.031 | 2.1 | 0.020 |
| 45.5 | 2 | 1260 | 3.2 | 0.036 | | |
| | 3 | 1260 | 3.6 | 0.042 | | |
| | 4 | 1260 | 1.1 | 0.024 | | |
| San Francisco | 1 | 1260 | 3.8 | 0.020 | 0.6 | 0.003 |
| 3.5 | 2 | 1260 | 5.8 | 0.031 | | |
| Oakland | 1 | ND | — | 0.005 | — | 0.006 |
| 19.0 | 2 | ND | — | 0.15 | | |
| Hyperion Waste | 1 | 1254 | 0.19 | 0.043 | 0.1 | 0.05 |
| Water 340 | 2 | 1254 | 0.37 | 0.057 | | |
| Hyperion Sludge | 1 | 1254 | 911 | ND | 1.6 | — |
| 5.0 | 2 | 1254 | 75.5 | ND | | |
| White Point | 1 | Mixture | 72[d] | 68.1 | 100 | 97 |
| 35[e] | 2 | 1254 and | | 98.7 | | |
| | 3 | 1254 | | 92.0 | | |
| | 4 | | | 52.6 | | |
| Terminal Island | 1 | 1254 | 12.8 | 0.098 | 0.35 | 0.002 |
| 9.1 | 2 | 1254 | 5.8 | 0.029 | | |
| Orange County | 1 | 1242 | 0.21 | 0.021 | 0.18 | 0.010 |
| 4.0 | 2 | 1254 | 0.04 | 0.035 | | |
| | 3[e] | 1254 | 0.23 | 0.061 | | |
| San Diego | 1 | ND | — | 0.057 | — | 0.04 |
| 56 | 2 | ND | | 0.033 | | |

[a] Mean ... [b] Based on ... [c] Sum of ... [d] ... [e] ...



and flow through the paper was 25 ml/min. PCB was identified by the presence of profile on chromatograms identical to those of one of the commercial PCB preparations (Aroclor 1254) (Fig. 3a). Other chromatograms were identical except for minor peaks to chromatograms of Aroclor 1242, mixtures of Aroclor 1242 and 1254 or of Aroclor 1260. PCB was quantified by direct comparison with standard preparations of these commercial mixtures. The profile of PCB peaks was unchanged by either saponification or passage through a sulfuric acid-Celite column[83] used for additional clean up of extracts.

Table 26 lists the PCB and DDT compounds in urban sewage outfalls in California. It is noted that daily outputs of PCB in the order of 2 kg are equivalent to a ton a year and because of the low solubility of these compounds in water and their high affinity for lipids, a large fraction of the PCB can be expected to pass into marine food chains. The highest output was recorded from Los Angeles County, one of the most industrialized areas of the state.

The utility of reversed liquid–liquid partition and GLC in the determination of PCBs and chlorinated pesticides in water and sewage sludge was described by ARLING AND JENSEN[84]. In this method the water is passed through a filter (3 g) containing a mixture of *n*-undecane and Carbowax 4000 monostearate on Chromosorb W, and the absorbed pesticides eluted with petroleum ether (10 ml). When detected by means of GLC with electron capture detection, the sensitivity is 10 ng/m³ of lindane with a sample size of 200 l. The recovery of added pesticides was 50–100% (DDT, 80%) and for PCB (Clophen 40 and A50, Bayer), 93–100%. The filter column was 30 cm × 1 cm I.D. with a glass filter disk G 1 (100–120 μ). GLC was carried out using a Varian Aerograph Model 204 chromatograph equipped with an electron capture detector and a 160 cm × 0.20 cm borosilicate glass column containing 4% methylsilicone oil or 8% fluorosilicone oil (QF-1) on HMDS-treated 80–100 mesh Chromosorb W. The detector and injector temperatures were 205° and 220°, respectively, the carrier gas was nitrogen (purified with a 6-in. molecular sieve) at a flow rate of 30 ml/min. The column temperature was chosen to give DDT a retention of 20 min (about 190°).

Table 27 lists the results from recovery experiments using a variety of filters for the separation of a number of chlorinated pesticides and PCB and shows the superiority of 60–80 mesh Chromosorb W, covered with a mixture of Carbowax monostearate and *n*-undecane. Figs. 4a and b show gas chromatograms of extracts from distilled water and natural water, respectively, analyzed on SF-96 columns. Fig. 4c



Fig. 4. (a) Gas chromatogram from the SF-96 separation of an extract from distilled water showing ... (b) ... SF-96 ... extract from natural water.

MONS 202132

TABLE 27

RESULTS FROM RECOVERY EXPERIMENTS

| Filter | Filter (g) | Water (l) | Flow speed (ml/min) | Standard pesticides* (pt/l) | Petroleum ether (ml) | Recoveries BHC | Lindane | Aldrin | Dieldrin | DDE | DDD | DDT | PCB |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Chromosorb W + ca 80% undecane | 1.5 | 6 | 65 | 5* | 10 | trace | 20 | 42 | 80 | 30 | 60 | 30 | |
| 10% Carbowax + | 1.5 | 6 | 65 | 5 | 10 | trace | trace | 28 | b | 17 | 17 | 61 | |
| 10% Carbowax + ca 80% undecane | 1.5 | 6 | 65 | 5 | 10 | 15 | 51 | 13 | 75 | 75 | 77 | 70 | |
| 10% Carbowax + 30% undecane[a] | 2 | 6 | 65 | 5 | 5 | 45 | 30 | 34 | b | 17 | 16 | 15 | |
| 10% Carbowax + 30% undecane[a] | 2 | 6 | 130 | 5 | 5 | 45 | 30 | 44 | b | 33 | 30 | 35 | |
| 10% Carbowax + 30% undecane | 3 | 6 | 65 | 5 | 5 | 68 | 40 | 59 | 97 | 65 | 104 | 53 | |
| 10% Carbowax + 30% undecane | 4 | 6 | 65 | 5 | 5 | 88 | 67 | 56 | 69 | 100 | 100 | 90 | |
| 10% Carbowax + 30% undecane | 2.5 | 18 | 65 | 1.33 | 5 | 64 | 42 | 13 | 62 | 100 | 100 | 68 | |
| 10% Carbowax + 30% undecane | 5 | 44 | 65 | 0.18 PCB[a] | 5 | 156 | 111 | 100 | 100 | 90 | 86 | 93 | |
| 10% Carbowax + 30% undecane | 1 | 5 | 65 | 10 | 5 | | | | | | | | 53 |
| 10% Carbowax + 30% undecane | 3 | 5 | 65 | 10 | 5 | | | | | | | | 93 |

* The standard had the following composition (ml concentration in ng/µl): HHC, 1.0; lindane, 1.0; aldrin, 1.0; DDE, 1.0; dieldrin, 5.0; DDT, 8.0; DDD, 1.0; and PCB (Clophen A50), 10.0.

[a] The extract was treated with sulfuric acid which destroyed the dieldrin.






Fig. 41. Gas chromatogram from an SE-96 column of an extract from sewage sludge.

illustrates a gas chromatogram from an SE-96 column of an extract from sewage sludge.

ZITKO[84] described the solubilization of PCB in water by non-ionic surfactants as well as the determination of concentration of the solubilized PCB by UV spectrophotometry and fluorescence and GLC and the toxicity of PCB to Atlantic salmon (*Salmo salar* parr.).

GC was carried out on a Varian 600D instrument equipped with a 250-mCi tritium electron capture detector, using a 5 ft × 1/8 in. glass column containing 4% SE-30 on 100–120 mesh Chromosorb W operated at 180°. UV and fluorescence spectra were recorded on a Beckman DK-2A and on a Perkin-Elmer MPF-2A instrument, respectively. Retention times and electron capture responses have been reported for mono- and dichlorobiphenyls[85,86]. EMERY AND GASSER[87] have described the GLC separation on polyphenyl thioethers of mono- and some di- and trichlorobiphenyls. As a general rule, chlorine substitution in position 2 shortens and vicinal disubstitution lengthens the retention time. The reported relative electron capture detector responses are 1, 0.2, 2.3, 7.6, 5.8, and 3.5 for 4-, 3-, and 2-mono-, and 4,4'-, 3,3'- and 2,2'-dichlorobiphenyl, respectively[88]. The retention times and electron capture detector responses of a number of PCB's have been reported by ZITKO *et al.*[88]. A Packard A7901 gas chromatograph with a 6 ft. × 4 mm glass column containing 4% SE-30 on 100–120 mesh acid-washed Chromosorb operated at 200° was used. The injector and detector temperatures were 210° and the carrier gas was nitrogen at a flow rate of 60 ml/min. D.C. voltage in the electron capture detector was 95 V and meter sensitivity was $1 \times 10^{-9}$ A. Solutions of chlorinated biphenyls in pesti-



MONS 202133

[illegible] were used in concentrations of 250 ng/ml for injections on co[illegible] and volumes from 2 to 5 µl were used to obtain peak heights of 30–70% of [illegible]. Relative retention times on a 2% SE-30 and electron capture detector [illegible] mixtures of chlorinated biphenyls and Aroclor 1254 and 1260 are shown in Table 2. Under these conditions ortho substitution at positions 2 and 6 shortens and vicinal substitution lengthens the retention time also in tri- and tetrachlorobiphenyls. In some instances these effects overshadow the effect of the number of chlorines in the molecule, which increases the retention time. The detector response increases strongly with increasing number of chlorine atoms in the molecule (e.g., the response of decachlorobiphenyl is 500 times greater than that of 2-chlorobiphenyl). PCB's found in wildlife are generally of the Aroclor 1254 or 1260 type[3,4,7]. In the study of ZITKO *et al.* above[58] these Aroclors gave fourteen and sixteen peaks with relative retention times from 0.48 to 3.28 and from 0.72 to 6.80 respectively. The presented data indicate that the electron capture response of all chlorinated biphenyls with four to nine atoms of chlorine per molecule is not likely to exceed the value of 1.6 relative to *p,p'*-DDE under the described conditions.

DE VOS AND PRINT[59] described the reversed phase partition TLC and GLC of the polychlorinated biphenyls.

Kieselgur plates impregnated with 8% liquid paraffin (b.p. 40–60°) were developed with acetonitrile–acetone–methanol–water (40:18:40:2), detected with

TABLE 26

RETENTION TIME AND ELECTRON CAPTURE DETECTOR RESPONSE OF CHLOROBIPHENYLS

| *Compound* | *Relative retention time (p,p'-DDE = 1.00)* | *Relative response per ng ± standard deviation (p,p'-DDE = 1.00)* |
|---|---|---|
| 4-Chlorobiphenyl | 0.17 | 0.0033 ± 0.00005 |
| 2-Chlorobiphenyl | 0.11 | 0.0030 ± 0.00004 |
| 3-Chlorobiphenyl | 0.14 | 0.0006 ± 0.00001 |
| 4,4'-Dichlorobiphenyl | 0.30 | 0.0135 ± 0.0002 |
| 2,4-Dichlorobiphenyl | 0.25 | 0.0155 ± 0.0001 |
| 2,5-Dichlorobiphenyl | 0.15 | 0.0131 ± 0.0001 |
| 2,3-Dichlorobiphenyl | 0.28 | 0.0455 ± 0.0002 |
| 3,4-Dichlorobiphenyl | 0.49 | 0.0450 ± 0.0004 |
| 2,6-Dichlorobiphenyl | 0.10 | 0.0815 ± 0.0030 |
| 2,4,4'-Trichlorobiphenyl | 0.39 | 0.295 ± 0.010 |
| 2,4,6-Trichlorobiphenyl | 0.24 | 0.270 ± 0.008 |
| 2,4',5-Trichlorobiphenyl | 0.51 | 0.290 ± 0.003 |
| 3,3',4,4'-Tetrachlorobiphenyl | 0.90 | 0.770 ± 0.027 |
| 2,3,6,4'-Tetrachlorobiphenyl | 0.33 | 0.0403 ± 0.00016 |
| 2,3,5,6-Tetrachlorobiphenyl | 0.70 | 0.625 ± 0.045 |
| 2,4,5,3'-Tetrachlorobiphenyl | 0.60 | 0.715 ± 0.010 |
| 2,3,6,3'-Tetrachlorobiphenyl | 0.50 | 0.505 ± 0.010 |
| 2,3,5,6,4'-Pentachlorobiphenyl | 1.00 | 1.30 ± 0.017 |
| 2,4,5,2',5'-Pentachlorobiphenyl | 0.75 | 0.515 ± 0.020 |
| 2,3,4,5,6-Pentachlorobiphenyl | 2.08 | 1.15 ± 0.011 |
| 2,3,4,6,2',5'-Hexachlorobiphenyl | 2.62 | 1.58 ± 0.020 |
| 2,3,5,6,2',5'-Hexachlorobiphenyl | 2.45 | 1.83 ± 0.070 |
| Decachlorobiphenyl | 8.20 | 4.64 ± 0.080 |
| [illegible] | | 0.0440 ± 0.0015 |
| [illegible] | | 1.35 ± 0.010 |






Fig. 42. Thin-layer chromatograms of (a) technical PCB mixture, (b) extract of owl liver and (c) *p,p'*-DDE. Kieselguhr impregnated with liquid paraffin, developed three times with acetonitrile–acetone–methanol–water (40:18:40:2), and visualized with silver nitrate and UV light.

Fig. 43. Gas chromatograms of (a) Phenochlor DP6 (0.6 µg) and (b) a hexane extract of owl liver. Instrumental conditions are given in Table 29.

TABLE 29

RELATIVE RETENTIONS (DIELDRIN = 1.00) OF PCB PEAKS 1–13 IN FIG. 43a

Instrument: Microtek MT 220. Column: glass, 6 ft × 1/4 in. O.D., packed with 3% OV-1 on 80–100 mesh Gas Chrom Q. Operating temperatures: column, 200°; injector, 220°; transfer line (Teflon-coated aluminium), 220°. Detection system: Dohrmann C-250-A microcoulometer with oxidative pyrolysis furnace and T-300-S titration cell. Sensitivity: 200 ohms. Gain: low (200). Recorder sensitivity: 1 mV. Chart speed: 1 cm/min.

| *Peak No.* | *Relative retention* |
|---|---|
| 1 | 0.71 |
| 2 | 0.84 |
| 3 | 1.04 |
| 4 | 1.18 |
| 5 | 1.28 |
| 6 | 1.47 |
| 7 | 1.60 |
| 8 | 1.73 |
| 9 | 4.00 |
| 10 | 4.33 |
| 11 | 4.80 |
| 12 | 5.13 |
| 13 | 5.76 |

MONS 202134

[illegible] with UV (Philips T[illegible] [illegible] (Black spots on a white background usually [illegible] Fig. [illegible] shows the [illegible] chromatograms of a technical PCB [illegible] DP6 containing 60% chlorine, an extract of owl liver, an [illegible] DDE).

Fig. [illegible] illustrates gas chromatograms of both Phenochlor DP6 and a hex[illegible] extract of owl liver. The instrumental conditions employed and the relative retention of the peaks in Fig. 43a are summarized in Table 29.

Peak numbers 5, 6, 8, 9, 10, 11, 12 and 13 in Fig. 43b have the same retention times as the corresponding peaks in Fig. 43a. Previous studies[1] utilizing a combination [illegible] chromatograph–mass spectrometer have shown that in the chromatogram of the technical PCB mixture containing 60% chlorine, peaks 1 and 2 are produced by pentachlorobiphenyls, peaks 3, 4, 5, 6 and 8 by hexachlorobiphenyls, peaks 7, 9, 10, 11 and 12 by heptachlorobiphenyls, and peak 13 by octachlorobiphenyl. Table 30

TABLE 30

RELATIONSHIP BETWEEN SPOTS OBTAINED BY THIN LAYER CHROMATOGRAPHY (FIG. 42a) AND PEAKS OBTAINED BY GAS CHROMATOGRAPHY (FIG. 43a) OF PHENOCHLOR DP6

| *Spot No.* | *Corresponding peak numbers* |
|---|---|
| 1 | 13 |
| 2 | 11 |
| 3 | 6, 9, 12 |
| 4 | 7, 8, 10 |
| 5 | 1, 4, 5 |
| 6 | 1, 6 |
| 7 | 1, 3 |

shows the relationship between spots obtained by TLC (Fig. 42a) and peaks obtained by GLC (Fig. 43a) of Phenochlor DP6. Of the chlorinated pesticides that are frequently detected in wildlife samples only the fungicide hexachlorobenzene (HCB) interferes with one of the PCB spots (No. 3). The lowest detectable amounts obtained with the above TLC system are of the order of 100 ng of single compounds such as DDE and HCB, and a few micrograms for PCB mixtures such as Phenochlor DP6.

The application of carbon skeleton chromatography to the qualitative differentiation between PCB's and DDT was demonstrated by Asai *et al.*[20]. PCB's and biphenyl yield identical carbon-skeleton chromatograms that are strikingly different from that of DDT. Comparisons of the relative retention times of products yielded by the PCB's with known compounds, plus partial identification by their UV spectra, suggested that the products formed at 300° catalyst temperature were cyclohexylbenzene and biphenyl, and at 260° were cyclohexylbenzene and a small amount of bicyclohexyl. The apparatus consisted of a NIE Beroza carbon skeleton determinator (National Instrument Laboratories, Inc., Rockville, Md.) attached to the [illegible] port of an Aerograph Model A-600-B gas chromatograph equipped with a flame ionization detector. For the electron capture gas chromatograms the same [illegible] chromatograph was equipped with a tritium detector. A sodium chloride [illegible] DMCS-treated 80–100 mesh [illegible]






Fig. 44. Carbon-skeleton chromatograms of biphenyl and Aroclor 1260 at two catalyst temperatures.

Chrom Q)[21] was used to change the carbon-skeleton determinator. The temperature of the catalyst bed was maintained at either 260° or 300°. The carbon-skeleton chromatograms were obtained using a 3 ft. × 1/8 in. stainless-steel column packed with 5% DC-200 on DMCS-treated 80–100 mesh Gas-Chrom Q. The column temperature was held at 105° and the hydrogen flow rate was 20 ml/min. The carbon-skeleton chromatograms of Aroclor 1260, biphenyl, bicyclohexyl- and cyclohexylbenzene were obtained on a 3 ft. × 1/8 in. stainless-steel column packed with 80–100 mesh




Fig. 45. Gas chromatograms of Aroclor 1260 and a mixture of p,p'-TDE and p,p'-DDT using the electron capture detector.

MONS 202135




Fig. 47. TICM chromatogram of a standard Aroclor 1254 mixture of polychlorinated biphenyls. Programmed temperature analysis: 2 min at 185°, to 210° at 5°/min, isothermal at 210°. (See text for remaining instrumental parameters and partial peak identification.)




Fig. 48. TICM chromatogram of a standard Aroclor 1260 mixture of polychlorinated biphenyls. Programmed temperature analysis: 2 min at 200°, to 210° at 5°/min, isothermal at 230°. (See text for remaining instrumental parameters and partial peak identification.)




Fig. 49. TICM chromatogram of human adipose tissue extract V. Programmed temperature analysis: 5 min at 180°, to 210° at 5°/min, isothermal at 210°.



50. HGC chromatogram of human adipose tissue extract B. Programmed temperature [illegible]

sec Levels. The mass spectra were recorded at 80 eV electron energy with 2500 V accelerating voltage and the biasing emission current was 500 μA. Helium carrier gas flow rate was 2 ml/min, and the injector temperature was 175°. The mass spectra were scanned magnetically over the range *m/e* 5 to *m/e* 500 in 6 sec. Figs. 49 and 50 illustrate the total ion current monitor chromatograms of human adipose tissue. Previous analysis by microcoulometric and electron capture GLC had indicated the presence of PCB residues and a modified Mills' procedure[7] was used for the preparation of the samples. Samples designated A and B (Figs. 49 and 50, respectively) were estimated to contain 200 p.p.m. and 600 p.p.m. total PCB as determined by electron capture GC.

This study indicated the enhancement of separation efficiency for the individual PCB components when analyzed by capillary column GC–mass spectrometry with no evidence of thermal degradation of any of the PCB components. All components of the Aroclors gave molecular ion groups of high intensity as would be expected from highly chlorinated biphenyl structures. The characteristic isotopic distribution pattern[8] corresponding to the number of chlorine atoms in the parent ion and chlorine-containing fragment ions was observed.

Two noteworthy features of the spectra were the relatively intense fragment ions produced by consecutive loss of chlorine atoms from the parent ion and the cross-




Fig. 51. Mass spectrum of a [illegible] contained in standard Aroclor 1232






Fig. 52. Mass spectrum of a heptachlorobiphenyl found in standard Aroclor 1260.

ence of intense doubly charged fragments within the mass spectra of most of the PCB compounds.

Figs. 51 and 52 show mass spectra obtained for a trichlorobiphenyl isomer in Aroclor 1232 and a heptachlorobiphenyl isomer in Aroclor 1260, respectively. Thus, in the chromatograms of Figs. 47 and 48 peaks 1 and 2 were shown to be tetrachlorobiphenyls, peaks 3–6, hexachlorobiphenyls, peaks 7 and 8–12, heptachlorobiphenyls, peaks 13 and 14, octachlorobiphenyls, peaks 15 and 16, nonachlorobiphenyls, and peak 17, decachlorobiphenyl.

The mass spectra of the components of adipose tissue of sample extract A (Fig. 49) revealed that peaks a, b, c, and d were polychlorinated biphenyls identical with those obtained respectively for peaks 6, 7, 9 and 10 of Aroclor 1254 (Fig. 47).

The mass spectrum of peak d (an isomer of hexachlorobiphenyl, mol. wt. 358) is shown in Fig. 53.

The mass spectra of components of adipose tissue B (Fig. 50) showed that peaks a through j were identical to those obtained for peaks 7 and 8–17 in the Aro-




Fig. 53. Mass spectrum of a hexachlorobiphenyl contained in human adipose tissue sample A (peak d, Fig. 49).

clor 1260 standard (Fig. 48). In addition, *p,p'*-DDE and *p,p'*-DDT, which were present at higher levels than those encountered in sample A, were also confirmed in this adipose tissue sample.

A system for the identification of multiple pesticide residues in foods by combined GC–mass spectroscopy was described by Beroza and Bowman[9] and this technique was used to demonstrate the absence of PCBs as an intermediate in the quantitation of *p,p'*-DDT and *p,p'*-DDE in fresh and smoked chubs. A Perkin Elmer Model 1 gas chromatograph was used equipped with a flame ionization de-

MONS 202137

...3.5 mV (full scale recorder). The column was a glass U tube, 6 ft ... ID, packed with 3% OV-17 on Supelcoport (100–120 mesh HP Chromosorb W ... The liquid phase consisted of 2% OV-17 on Supelcoport (100–120 mesh HP Chromosorb ... carrier gas. Separations were performed isothermally with column temperatures of 180–215° depending on the pesticides being separated with the ... effluent split between the mass spectrometer and the flame ionization detector.

The mass spectrometer was an AEI MS-12 with an electron bombardment ion source. The electron trap was set at 100 eV and the source temperature maintained at ca. 200° by the heat of the electron emitting filament. The electron bombardment energy of 24 eV has the advantage over the more customarily used 70 eV of increasing the relative intensity of the molecular ion chlorine isotope cluster. The ion accelerating voltage was 8 kV with the electron multiplier detector set at a gain of $7 \times 10^5$ with mass spectra scanned magnetically from *ca. m/e* 500 to *m/e* 12 in about 12 sec. A total ion current monitor detector was situated between the ion source and the mass analyzing magnet. (This detector intercepts a portion of the ion beam and displays it as a voltage signal on a 10-mV potentiometric strip chart recorder.) The interface between the GLC and the mass spectrometer was essentially as reported by MARKEY[38].

Cheese and cocoa beans were prepared for analysis by the fatty food procedure described in the *Pesticide Analytical Manual*[37]. The pesticides in the isolated fish sample (fresh and smoked chubs) were extracted with petroleum ether and transferred into acetonitrile and cleaned up successively over Florisil and silicic acid columns. The latter utilized the procedure of ARMOUR AND BURKE[35] to remove any PCB that might have been present. Fig. 54 illustrates the flame ionization detection (FID) and total ion current monitor (TICM) chromatograms obtained from a smoked chub sample after silicic acid chromatography and shows several peaks, two of which have the same retention times as those of *p,p'*-DDE and *p,p'*-DDT. It was determined




...54. FID and TICM chromatograms obtained from a smoked chub sample after silicic acid chromatographic separation. Solvent front (▲) appeared at which glass valve suspended. The FID was at 25 MS sensitivity. ... DDE ... DDT ...

... FID chromatograms of Aroclor 1254 standard, *p,p'*-DDE, and *p,p'*-DDT mixture. ... Width of *p,p'*-DDE ... DDT peak defined by ...



(using the procedure of ARMOUR AND BURKE[35]) that the original PCB interference in the chubs was caused by Aroclor 1254. An FID chromatogram of Aroclor 1254 (Fig. 55) shows several peaks that can interfere with the determination of *p,p'*-DDE and *p,p'*-DDT. The regions A and B represent the width of the *p,p'*-DDE and *p,p'*-DDT peaks respectively. The centers of the regions represent the peak maximum for these pesticides. Mass spectra were obtained for the GLC components of Aroclor 1254 in the center of regions A and B, i.e., Aroclor 1254 components with the same retention




Fig. 56. Mass spectra of (a) Aroclor 1254 components with the same retention time as that of *p,p'*-DDE and (b) Aroclor 1254 components with the same retention times as that of *p,p'*-DDT.

time as that of *p,p'*-DDE (Fig. 56a) and Aroclor 1254 components with the same retention time as that of *p,p'*-DDT (Fig. 56b). Fig. 57 shows that mass spectra of (a) *p,p'*-DDE standard at 24 eV and (b) *p,p'*-DDE in a sample showing PCB carry-over after a silicic acid column separation procedure. The *m/e* 326 peak shows the presence of PCB. This carry-over of PCB into the pesticide-containing fraction occurred when extracts representing more than 0.4 g of fat were chromatographed over silicic acid.

Another objective of the study of BUTLER AND BARRY[36] was to determine if eluates obtained from the *Pesticide Analytical Manual*[37] procedure could be used to confirm the identification of pesticide residues by combined GLC–MS. The FID and TICM chromatograms (Fig. 58) were obtained from a 6% eluate of a frog leg sample and show that the mass spectra of the four GLC peaks are identical. (The mass spectra

MONS 202138



Fig. 57. Mass spectra of (a) a *p,p'*-DDT standard recorded at 21 eV and (b) *p,p'*-DDE in a sample showing PCB carry-over after a silicic acid column separation procedure. The *m/e* 326 peak was the presence of PCB.

of standard α-, β-, γ- and δ-isomers of HCH[4] also appear to be identical and match the mass spectra of the four GLC peaks). Hence identification of which isomers of HCH were present in the sample required the combination of GLC retention time and the mass spectra produced.




Fig. 58. GLC traces (ECD and FID) of a hexane solution (from the fatty tissue of rat) containing [illegible] PCB peaks (marked MS [illegible] (SE-30, 1.5%, OV-1, 1.5%) GLC column (6 ft × 1/8 in.) at 180°.



The occurrence and distribution of PCBs in the River Rhine and the coastal areas of The Netherlands has been reported by KORMAN *et al.*[5]. The identification was carried out by GC and mass spectrometry. For this purpose a Varian MAT Model CH 4 mass spectrometer was coupled to a Varian Aerograph Type A550-B oven utilizing an interface under conditions described by TENNISON[37] or BRAUMAN and BRUNNER[38]. The relative retention times ($R_r$) were estimated with a Varian Aerograph Model 204-1B gas chromatograph with electron capture detection. Pyrex glass columns (5 ft × 1/8 in.) filled with 10% DC-200 on 80–100 mesh Gas-Chrom Q were used in both gas chromatographs, and the columns were operated at 200° with helium or nitrogen as carrier gas. The tissues of the indicator organisms used were extracted with petroleum ether in a Soxhlet extraction apparatus after drying with anhydrous sodium sulfate. The samples were cleaned up by dimethylformamide partition[39] and column chromatography over activated Florisil. The apolar compounds including the PCB's were eluted with hexane and then a 10% diethyl ether in hexane solution was used to elute dieldrin and aldrin. Tables 31 and 32 list the mass numbers and numbers of chlorine atoms per molecule of peaks with relative retention times (relative to dieldrin = 1.00) longer than 0.20 and shorter than 1.00 and of peaks longer than 1.00, respectively. Mass spectra were taken from all peaks present in the total ion current chromatogram obtained with the double ion source of the CH 4 mass spectrometer. In Figs. 59a and b gas chromatograms are shown of an eider extract and a technical PCB mixture (Phenoclor DP6), illustrating a strong resemblance between the electron capture chromatograms and the ion current chromatograms. Both from the mass numbers and the numbers of chlorine atoms per molecule, it was concluded that most of the compounds present in the extracts could be identified as PCB's.




Fig. 59. Gas chromatograms of (a) an eider extract, (b) a technical PCB mixture (Phenoclor DP6) and (c) [illegible] of the eider extract.

MONS 202139

TABLE 31

MASS NUMBERS AND NUMBERS OF CHLORINE ATOMS PER MOLECULE (IN PARENTHESES) OF PEAKS WITH RELATIVE RETENTION TIMES (RELATIVE TO DIELDRIN = 1.00) LONGER THAN 0.20 AND SHORTER THAN 1.00

With the exception of HCB, telodrin and compound 3,6 (8) all compounds identified are conformable to PCB's. A 10% [illegible] Gas Chrom Q column was used; temperature, [illegible] Mass number calculations are based on [illegible] HCB [illegible] chlorobenzene, telodrin [illegible] 1,3,4,5,6,7,8,8-octachloro-[illegible] hexahydro-3,7-methano-isobenzofuran [illegible] at the given retention time [illegible] present in the chromatogram.

| Sampling area | Type of sample | $R_x$ 0.25 | 0.32 | 0.33 | 0.34 | 0.40 | 0.48 | 0.57 | 0.61 | 0.65 | 0.71 | 0.7[illegible] | 0.8[illegible] | 0.8[illegible] |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | Roach (*Leuciscus rutilus*) total body extracts 1967, 1968 | HCB 282 (6) | 256 (3) | 256 (3) | | 256 (3) + 290 (4) | 290 (4) | 290 (4) | 290 (4) | | 290 (4) | | 324 (5) | 324 (5) |
| 1 | Groundling (*Gobio gobio*) total body extracts 1968 | + | + | + | | + | + | | + | | + | | + | + |
| 2 | Mussel (*Mytilus edulis*) 1968 | + | | | + | + | | | | + | | | + | |
| 3 | Sand eel (*Ammodytes lanceolatus*) total body extracts 1966 | + | | + | | + | + | + | + | | + | | | + |
| 3 | Sandwich tern (*Sterna sandvicensis*) tissue and egg extracts 1965, 1966 | HCB 282 (6) | | | | 256 (3) | 290 (4) | 290 (4) | Telodrin 406 (8) | | 290 (4) + 376 (8) | | 324 (5) | |
| 3 | Eider (*Somateria mollissima*) tissue and egg extracts 1966, 1967 | HCB 282 (6) | | | | 256 (3) | | | Telodrin 406 (8) | | 290 (4) + 376 (8) | | 324 (5) | |
| | Phenochlor DP6 | | | | | | | | | | | 324 (5) | 324 (5) | |
| | Arochlor 1260 | | | | | | | | | | | 324 (5) | 324 (5) | |
| | Clophen A60 | | | | | | | | | | | 324 (5) | 324 (5) | |

TABLE 32

MASS NUMBERS AND NUMBERS OF CHLORINE ATOMS PER MOLECULE (IN PARENTHESES) OF PEAKS WITH RELATIVE RETENTION TIMES (RELATIVE TO DIELDRIN = 1.00) OF PEAKS LONGER THAN 1.00

With the exception of HDE all compounds identified are conformable to PCB's. + = At the given retention time a peak is present in the chromatogram. For further information, see Table 31.

| Sampling area | Type of sample | $R_x$ 1.01 | 1.17 | 1.23 | 1.45 | 1.57 | 1.69 | 1.96 | 2.27 | 2.67 | 3.20 | 3.58 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | Roach (*Leuciscus rutilus*) total body extracts 1967, 1968 | DDE 316 (4) + 324 (5) | 358 (6) | 358 (6) | 358 (6) | 392 (7) | 358 (6) 392 (7) | 392 (7) | 392 (7) | 392 (7) | 392 (7) | 426 (8) |
| 1 | Groundling (*Gobio gobio*) total body extracts 1968 | + | + | + | + | + | + | + | + | + | + | + |
| 2 | Mussel (*Mytilus edulis*) 1968 | + | + | + | + | + | + | + | + | + | + | + |
| 3 | Sand eel (*Ammodytes lanceolatus*) 1966 | + | + | + | + | + | + | + | + | + | + | + |
| 3 | Sandwich tern (*Sterna sandvicensis*) tissue and egg extracts 1965, 1966 | DDE 316 (4) + 324 (5) | | 358 (6) 324 (5) | 358 (6) | | 358 (6) | 392 (7) | 392 (7) | 392 (7) | 392 (7) | 426 (8) |
| 3 | Eider (*Somateria mollissima*) tissue and egg extracts 1966, 1967 | DDE 316 (4) + 324 (5) | | 358 (6) + 324 (5) | 358 (6) | | 358 (6) | 392 (7) | 392 (7) | 392 (7) | 392 (7) | 426 (8) |
| | Phenochlor DP6 | 358 (6) | 358 (6) | 358 (6) | 358 (6) | 392 (7) | 358 (6) | 392 (7) | 392 (7) | 392 (7) | 392 (7) | 426 (8) |
| | Arochlor 1260 | 358 (6) | 358 (6) | 358 (6) | 358 (6) | 392 (7) | 358 (6) | 392 (7) | 392 (7) | 392 (7) | 392 (7) | 426 (8) |
| | Clophen A60 | 358 (6) | 358 (6) | 358 (6) | 358 (6) | 392 (7) | 358 (6) | 392 (7) | 392 (7) | 392 (7) | 392 (7) | 426 (8) |



MONS 202140

[illegible] metal compounds formed were hexachlorobenzene (HCB), p,p'-DDE and dieldrin.

The results indicate that the lower chlorinated PCB components are more readily metabolized than the more highly chlorinated ones. This suggests the presence of [illegible] PCBs are less persistent and this was confirmed in an experiment in which [illegible] were fed a diet containing 2 000 ppm of Aroclor 1254 for seven days and then given a diet containing 2 000 ppm of Aroclor DF6 for a further seven days. A chromatogram representing the fat extract of breast of these birds is shown in Fig. 5(a) and shows that a number of components present in the original mixture were metabolized, particularly the more highly chlorinated ones. Residues in the livers and fat of the quail were measured on a semi-quantitative scale by peak height comparison of peak $R_F = 1.45$ in the Aroclor DF6 mixture as a standard. The residue levels calculated in this manner were about twenty times higher in the quail tissues than in the livers and breasts of the birds. The quail developed hydropericardium with a dose of 2 000 ppm. Lucas *et al.*[7] have reported hydropericardium in White Leghorn cockerels dosed with 1 000 ppm in their diet.

Bowes *et al.*[8] described the identification of polychlorinated biphenyls in two analogues by combined GLC–mass spectrometry. The separation and identification of the unknown components were obtained with an LKB Model 9000 GLC–MS equipped with the stainless steel molecule separator system of Ryhage[10]. The 9 ft × 0.25 in. glass column was packed with 1% SE-30 on 100–120 mesh Gas-Chrom Q. The operating temperatures were flask heater 220°, GLC oven 180°, separator 240°, and ion source 270°. The carrier gas was helium at a rate of 35 ml/min, the ionization potential was 70 eV, trap current 60 μA, accelerating voltage 3.5 kV and the scan time was 12 sec from *m/e* 2–400. TLC on silica gel as described by Mulhern[9] was




Fig. 5. (a) Total ion current chromatogram of (1) standard PCB mixture (Aroclor 1254, 1 g/l), (b) eagle carcass 65 056, TLC zone IV, (c) eagle carcass 65 056, TLC zone III, and (d) eagle carcass 65 50, TLC zone IV. GLC conditions: 0.5% × 0.25 in. 1% SE-30 column, oven temperature 180°, helium flow 35 ml/min.






Fig. 6a. Mass spectrum of peak 1 (Figs. 6a, b and d), empirical formula $C_{12}H_4Cl_4$.




Fig. 6b. Mass spectrum of peak 2 (Figs. 6a, b and d), empirical formula $C_{12}H_5Cl_5$.




Fig. 6c. Mass spectrum of peak 3 (Figs. 6a–c), empirical formula $C_{12}H_6Cl_4$.




Fig. 6d. Mass spectrum of peak 4 (Figs. 6a and c), empirical formula $C_{12}H_5Cl_5$.



MONS 202141



Fig. 65. Mass spectrum of peak 5 (Figs 60a, b and d), empirical formula $C_{12}H_6Cl_4$.



Fig. 66. Mass spectrum of peak 5 (Fig. 60c: 1254-III (total ion current chromatogram not shown for Aroclor 1254, TLC zone III)), empirical formula $C_{12}H_6Cl_4$.



Fig. 67. Mass spectrum of peak 7 (Figs 60a, b and d), empirical formula $C_{12}H_5Cl_5$.



Fig. [illegible]. Mass spectrum of peak 7 (Figs 60c, 1254-III (total ion current chromatogram not shown for Aroclor 1254, TLC zone III)), empirical formula $C_{12}H_5Cl_5$.





Fig. 69. Mass spectrum of peak 9 (Figs 60a and c), empirical formula $C_{12}H_5Cl_5$.



Fig. 70. Mass spectrum of peak 10 (Figs 60a, b, and d), empirical formula $C_{12}H_4Cl_6$.



Fig. 71. Mass spectrum of peak 11 (Fig. 60c: 1254-III (total ion current chromatogram not shown for Aroclor 1254, TLC zone III)), empirical formula $C_{12}H_4Cl_6$.



Fig. 72. Mass spectrum of peak 12 (Figs 60a, b, and d), empirical formula $C_{12}H_3Cl_7$.

MONS 202142




Fig. 74. Mass spectrum of peak 12 (Figs. 60a, b and d), empirical formula $C_{12}H_4Cl_6$




Fig. 74. Mass spectrum of peak 14 (Figs. 60b and d), empirical formula $C_{12}H_4Cl_6$




Fig. 75. Mass spectrum of peak 15 (Figs. 60a, b and d), empirical formula $C_{12}H_3Cl_6$

used to separate interfering organochlorine pesticides. The major peaks as shown in the total ion current chromatogram for Aroclor 1254 (Fig. 60a) are numerically labeled in order of their retention time. The peaks in total ion current chromatograms for eagle samples (Fig. 60d) are labeled as they relate to the Aroclor standard and show that peaks 1–16 have approximately the same retention times as identically numbered peaks in Aroclor 1254. Figs. 61–75 show mass spectra for peaks in the Aroclor standard and corresponding peaks in eagle tissue samples.

Table 33 lists the relative retention time of PCB's and some commonly occurring organochlorine pesticides obtained using a 1% SE-30 column. The similarity of relative retention times (compared to p,p'-DDE) of a number of PCB peaks and the organochlorine pesticides prohibits the utility of GLC identification alone.



TABLE 33

RELATIVE RETENTION TIME OF PCB'S AND SOME COMMONLY OCCURRING ORGANOCHLORINE PESTICIDES (p,p'-DDE = 1.00)

| PCB | | Relative retention time | Compound | Relative retention time |
|---|---|---|---|---|
| Peak No. | Mol. wt. | | | |
| 1 | 256 (3)* | 0.39 | Heptachlor | 0.43 |
| 2 | 290 (4) | 0.49 | | |
| 3 | 290 (4) | 0.53 | Aldrin | 0.53 |
| 4 | 290 (4) | 0.58 | | |
| 5 | 290 (4), 324 (5) | 0.70 | Heptachlor epoxide | 0.66 |
| 6 | 324 (5) | 0.79 | | |
| 7 | 324 (5) | 0.83 | Dieldrin | 0.96 |
| 8 | 324 (5) | 0.97 | o,p'-DDD | 1.02 |
| 9 | 358 (6) | 1.04 | Endrin | 1.08 |
| 10 | 358 (6) | 1.23 | p,p'-DDD | 1.27 |
| 11 | 358 (6) | 1.46 | o,p'-DDT | 1.35 |
| 12 | 358 (6) | 1.72 | p,p'-DDT | 1.69 |
| 13 | 358 (6), 392 (7) | 2.01 | | |
| 14 | 392 (7) | 2.32 | | |
| 15 | 392 (7) | 2.80 | | |
| 16 | 392 (7) | 3.38 | | |

* The number in parentheses following the molecular weight is the number of chlorine atoms in the molecule.




Fig. 76. 70 eV mass spectra (C.E.C. 21 cyclo-instrument) of a 1:1 mixture of Aroclor 1242 and DDT at different sample temperatures. Isotope clusters at m/e 222, 256, 290 and 324 result from individual components of Aroclor; the group of peaks at m/e 186 are mainly fragment ions from the [illegible]

MONS 202143

TABLE 34

MASS SPECTRA OF POLYCHLORINATED BIPHENYLS

The first line in the heading is sample, the second peak number and the third molecular weight with the number of chlorine atoms in the molecule in parentheses

| m/e | 1254 III 6 324 (5) | 68-050 III 9 324 (5) | 66-050 III 10 358 (6) | 1254 14 392 (7) | 68-056 II 14 392 (7) | 1254 14 392 (7) | 68-056 II 16 392 (7) | 1254 13 392 (7) |
|---|---|---|---|---|---|---|---|---|
| 98 | 10.2 | 5.4 | 6.3 | 8.2 | 10.9 | 10.1 | 7.6 | 10.5 |
| 99 | 7.0 | 3.7 | 4.5 | 4.1 | 5.4 | 5.0 | 4.1 | 4.1 |
| 107 | 3.9 | 1.0 | 2.1 | 4.1 | | | 3.4 | 4.5 |
| 109 | 4.5 | 2.2 | 5.3 | 11.8 | 15.9 | 13.6 | 10.7 | 11.1 |
| 109 | 20.2 | 12.5 | 16.0 | 15.3 | 20.0 | 16.4 | 10.0 | 22.5 |
| 110 | 17.0 | 9.3 | 12.3 | 13.8 | 14.1 | 14.5 | 8.3 | 10.1 |
| 111 | 9.5 | 3.3 | 4.0 | 5.3 | 6.4 | 6.8 | 4.1 | 5.5 |
| 119 | | | 1.2 | | 4.1 | 4.5 | 3.1 | 4.5 |
| 120 | 3.2 | 1.3 | 2.5 | 5.9 | 6.4 | 6.4 | 5.2 | 5.4 |
| 121 | 3.9 | 1.5 | 2.8 | 5.3 | 5.4 | 7.3 | 4.8 | 5.3 |
| 122 | 6.4 | 3.4 | 5.7 | 7.1 | 8.2 | 6.8 | 5.9 | 7.1 |
| 123 | 6.6 | 3.0 | 3.1 | | 5.9 | | 4.1 | 3.4 |
| 125 | 3.6 | | | | | | | |
| 130 | | 1.9 | 9.1 | 17.1 | 18.9 | 19.1 | 15.2 | 19.2 |
| 137 | 29.5 | 22.9 | 11.2 | 22.3 | 20.7 | 25.4 | 16.2 | 23.9 |
| 138 | 27.3 | 21.2 | 6.0 | 14.1 | 9.5 | 14.5 | 6.7 | 9.7 |
| 139 | 9.3 | 5.6 | 1.8 | | | 5.4 | | |
| 141 | | | 1.7 | | 5.9 | | 3.4 | 3.2 |
| 151 | | 1.6 | 3.9 | 8.5 | 10.0 | 10.0 | 7.2 | 9.1 |
| 153 | 4.8 | 2.4 | 3.7 | 8.2 | 9.1 | 10.0 | 5.9 | 7.2 |
| 154 | | 1.4 | 2.1 | 4.4 | 5.9 | | 3.4 | 5.9 |
| 155 | 1.9 | 1.6 | 2.8 | 2.9 | 4.5 | | 3.1 | 3.2 |
| 156 | | | | | | | | |
| 157 | 2.7 | | | | | | | |
| 159 | 5.2 | | | | | | | |
| 163 | | | | 10.3 | 13.1 | 11.8 | 10.3 | 8.6 |
| 164 | 6.6 | 4.6 | 27.4 | 15.2 | 29.5 | 20.9 | 11.7 | 37.5 |
| 165 | 9.6 | 5.8 | 34.0 | 16.2 | 29.4 | 27.3 | 11.0 | 35.1 |
| 166 | 6.9 | 4.2 | 18.9 | 14.4 | 16.5 | 13.6 | 9.3 | 21.3 |
| 167 | 7.3 | 3.7 | 8.5 | 6.5 | 8.6 | 10.0 | 5.2 | 11.1 |
| 168 | 5.0 | 2.9 | 5.3 | 3.2 | | | 2.1 | 3.3 |
| 169 | 13.4 | 5.0 | 2.8 | 2.9 | | | 3.1 | 2.8 |
| 170 | 5.7 | | | 3.5 | | | | 3.5 |
| 50 | | | 4.5 | 4.4 | | | 4.5 | 4.2 |
| 157 | | | 2.1 | 2.9 | | | 2.8 | 3.1 |
| 158 | | | 1.7 | 2.3 | | | | 2.9 |
| 191 | | | 4.5 | 25.3 | 29.1 | 27.3 | 26.7 | 17.5 |
| 192 | 9.3 | | 3.8 | 42.9 | 42.7 | 43.6 | 48.3 | 27.1 |
| 193 | 13.5 | | 3.7 | 30.6 | 31.4 | 30.0 | 31.0 | 21.1 |
| 194 | 9.3 | | 1.9 | 12.3 | 13.4 | 10.9 | 9.5 | 7.3 |
| 195 | 3.2 | | | | | | | |
| 197 | 3.2 | | | | | | | |
| 168 | | | | | | | | |
| 169 | | | | | | | | |
| 170 | | | | | | | | |
| 171 | | | | | | | | |
| 179 | | | 9.1 | 5.9 | 9.1 | 8.2 | 6.0 | |
| 180 | | | 10.0 | 7.1 | 13.2 | 7.7 | 6.9 | |
| 181 | | | 15.1 | 8.2 | 10.9 | 9.1 | 6.5 | |
| 182 | 2.0 | 1.5 | 11.1 | 8.8 | 8.7 | 10.0 | 9.5 | |
| 183 | 6.1 | 4.6 | 8.4 | 5.9 | 7.4 | 7.3 | | |
| 191 | 25.2 | 17.5 | 5.7 | 9.1 | 5.9 | 10.0 | 4.3 | |
| 193 | 6.8 | 3.7 | 1.8 | 4.2 | | 5.1 | | |



TABLE 34 (continued)

| m/e | 1254 III 6 324 (5) | 68-050 III 9 324 (5) | 68-050 III 10 358 (6) | 1254 14 392 (7) | 68-056 II 14 392 (7) | 1254 16 392 (7) | 68-056 II 16 392 (7) | 1254 13 392 (7) |
|---|---|---|---|---|---|---|---|---|
| 196 | 9.1 | 5.8 | 1.1 | 3.2 | | | | 1.9 |
| 196 | | | | 9.8 | 9.1 | 8.2 | 7.6 | 5.0 |
| 197 | | | | 17.6 | 19.5 | 12.7 | 13.8 | 11.4 |
| 198 | | | | 15.6 | 17.3 | 13.6 | 13.4 | 11.7 |
| 199 | | | | 9.1 | 8.6 | 6.4 | 9.0 | 5.1 |
| 216 | | | 1.2 | 6.5 | 7.7 | 5.4 | 5.5 | 5.0 |
| 217 | 7.9 | | 3.7 | 9.2 | 8.6 | 7.3 | 6.5 | 7.5 |
| 218 | 9.5 | 6.7 | 23.6 | 11.5 | 16.4 | 10.9 | 6.0 | 25.8 |
| 219 | 16.6 | 6.2 | 6.6 | 7.1 | 7.7 | 8.2 | 5.5 | 8.3 |
| 220 | 7.3 | 3.8 | 16.0 | 12.3 | 11.8 | 15.4 | 4.6 | 19.2 |
| 221 | 8.6 | 4.2 | 2.9 | 3.8 | | 4.5 | | 4.4 |
| 222 | 2.7 | 1.4 | 2.8 | 6.8 | 4.5 | 7.3 | | 5.5 |
| 251 | | | | | | | | 2.1 |
| 252 | | | 5.7 | 25.9 | 28.2 | 25.4 | 25.5 | 20.8 |
| 253 | | | 6.6 | 8.2 | 9.8 | 9.1 | 6.9 | 9.3 |
| 254 | 59.1 | 34.2 | 11.3 | 40.0 | 32.7 | 40.0 | 45.5 | 28.6 |
| 255 | 10.2 | 7.5 | 7.5 | 8.8 | 7.7 | 9.1 | 3.7 | 10.4 |
| 256 | 61.4 | 53.7 | 8.5 | 21.5 | 14.1 | 25.4 | 10.7 | 15.5 |
| 257 | 9.3 | 7.3 | 3.4 | 5.3 | 4.5 | 6.4 | 2.2 | 5.3 |
| 258 | 20.4 | 15.8 | 2.7 | 7.1 | 3.6 | 9.1 | 2.4 | 4.7 |
| 259 | 4.1 | 3.3 | 1.4 | | | | | |
| 260 | 4.2 | 2.7 | | | | | | |
| 261 | 1.8 | 1.1 | | | | | | |
| 261 | 1.9 | | | | | | | |
| 286 | | | | 3.5 | 4.5 | 4.5 | 4.6 | 5.3 |
| 287 | | | | 5.9 | 7.3 | 7.3 | 5.9 | 5.8 |
| 288 | | | 46.2 | 12.3 | 24.5 | 13.6 | 9.0 | 13.6 |
| 289 | 29.5 | 14.2 | 7.5 | 12.3 | 13.2 | 11.8 | 9.3 | 69.4 |
| 290 | 20.4 | 3.7 | 59.4 | 23.5 | 34.1 | 26.4 | 10.7 | 14.2 |
| 291 | 43.2 | 17.9 | 20.4 | 10.0 | 10.0 | 10.9 | 5.5 | 43.0 |
| 292 | 23.9 | 3.7 | 3.0 | 15.8 | 20.9 | 23.6 | 7.1 | 8.9 |
| 293 | 22.7 | 0.9 | 5.7 | 5.9 | 5.7 | 6.8 | 3.1 | 12.2 |
| 294 | 11.4 | 1.5 | 8.0 | 9.4 | 7.0 | 10.0 | 2.8 | 3.6 |
| 295 | 5.7 | 2.1 | 2.6 | 3.3 | | | | 3.0 |
| 296 | 2.9 | | 1.7 | 2.9 | | | | |
| 297 | 0.7 | | | | | | | |
| 322 | | | | 38.8 | 40.9 | 40.9 | 40.3 | 24.5 |
| 323 | | | 12.3 | 9.7 | 9.1 | 10.4 | 8.6 | 48.6 |
| 324 | 63.6 | 62.5 | 13.2 | 82.3 | 68.2 | 82.1 | 69.0 | 52.8 |
| 325 | 10.4 | 9.2 | 20.7 | 17.6 | 13.6 | 17.3 | 13.6 | 33.3 |
| 326 | 100.0 | 100.0 | 19.8 | 86.5 | 50.4 | 74.5 | 48.3 | 43.4 |
| 327 | 13.9 | 12.9 | 15.1 | 14.7 | 10.0 | 16.4 | 9.0 | 19.7 |
| 328 | 63.6 | 62.5 | 12.3 | 32.9 | 20.9 | 36.4 | 17.9 | 21.4 |
| 329 | 8.9 | 7.9 | 5.7 | 7.1 | 5.0 | 9.2 | 4.0 | 8.6 |
| 330 | 21.6 | 18.3 | 3.8 | 10.6 | 6.4 | 13.3 | 5.2 | 6.4 |
| 331 | 3.7 | 2.6 | 1.2 | 3.5 | | | | |
| 332 | 3.4 | 2.7 | | 2.9 | | | | |
| 333 | 0.6 | | | | | | | |
| 357 | | | | 14.1 | 3.6 | 13.6 | 9.0 | 55.3 |
| 358 | | | 52.8 | 11.2 | 3.2 | 11.8 | 5.9 | 21.1 |
| 359 | | | 9.4 | 27.6 | 7.2 | 24.5 | 17.9 | 100.0 |
| 360 | | | 100.0 | 20.0 | 3.7 | 19.1 | 10.2 | 23.6 |
| 361 | | | 10.0 | 12.0 | 5.4 | 20.0 | 10.2 | 81.9 |

(Continued on p. 418)

MONS 202144

TABLE 34 (*Continued*)

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | 1-54 (1) | 6%-050 (1) | 08-050 (1) | 15-050 (1) | 1251 | 58-056 (1) | 125 |
| | 6 | 9 | 10 | 14 | 16 | 16 | 13 |
| | 354 (5) | 354 (5) | 358 (6) | 392 (7) | 392 (7) | 392 (7) | 392 |
| 82 | | | 80.2 | 15.0 | 17.0 | 17.3 | 7.9 |
| 83 | | | 32.3 | 10.6 | 23.6 | 20.0 | 7.2 |
| 91 | | | 34.0 | 7.1 | 6.5 | 7.3 | 3.4 |
| 95 | | | 4.7 | 3.2 | 6.4 | 3.6 | 1.7 |
| 190 | | | 8.5 | 2.1 | | | |
| 192 | | | 1.1 | | | | |
| 258 | | | 1.2 | | | | |
| 141 | | | | 47.1 | 43.2 | 47.3 | 51.7 |
| 293 | | | | 10.6 | 10.4 | 13.6 | 10.7 |
| 361 | | | | 100.0 | 100.0 | 100.0 | 100.0 |
| 195 | | | | 17.6 | 17.3 | 20.0 | 17.6 |
| 196 | | | | 97.1 | 95.4 | 90.0 | 96.5 |
| 197 | | | | 14.7 | 13.9 | 16.4 | 15.2 |
| 198 | | | | 52.9 | 46.8 | 51.2 | 53.4 |
| 199 | | | | 7.6 | 7.7 | 10.0 | 7.9 |
| 109 | | | | 15.9 | 16.6 | 17.3 | 16.5 |
| 191 | | | | 2.6 | 2.5 | 2.7 | 2.4 |
| 40 | | | | 3.5 | 3.4 | 3.6 | 2.9 |

peaks have similar retention times although in some cases mass spectral data show the compounds to be isomers rather than the same compound.

The identification of PCB's and DDT in mixtures by mass spectrometry was reported by Herzinger *et al.*[101]. Fig. 76 illustrates 70 eV mass spectra of a 1:1 mixture of Aroclor 1242 and DDT at different sample temperatures. The peaks arising from the individual major components of the Aroclor mixture can easily be distinguished from major peaks found in the most common chlorinated insecticides. A useful spectrum of Aroclor 1242 could be obtained at low temperatures by direct introduction with the aid of a temperature-controlled probe[101], and fractional sublimation from many chlorinated pesticides directly in the ion source seems possible (see Fig. 76).

## VII. ANALYSIS OF CHLORINATED NAPHTHALENES AND DIBENZOFURANS

Chlorinated naphthalenes possess a variety of suggested uses related to their electrical, flame-retardant and fungus resistant properties, stability, and compatibility with other materials[102]. Many of their physical properties and suggested uses are similar to those of polychlorinated biphenyls[3]. The behavior of chlorinated naphthalenes on analytical methods for organochlorine pesticides and PCB's was studied by Armour and Burke[103]. The chlorinated naphthalenes studied were representative Halowax (Hoppers Co., Inc.) samples, e.g., Halowax 1013, a mixture of tri- and tetrachloronaphthalenes; Halowax 1051, octachloronaphthalene; and Halowax 1099, a mixture of trichloro- and tetrachloronaphthalenes. GLC of the tested naphthalenes showed that their behavior is similar to that of the organochlorine pesticides. Retention time and response data for the three Halowax [illegible] are given in Table [illegible].



TABLE 35

GLC DATA FOR CHLORINATED NAPHTHALENES

GLC columns: glass 6 ft. × 4 mm I.D. 10% DC-200 on 80–100 mesh HP Chromosorb W and 10% DC-200/15% QF-1 (1:1) on 80–100 mesh HP Chromosorb W. Operating conditions: nitrogen flow rate, 120 ml/min; column and detector temperature, 200°; injector temperature, 225°. Concentric design electron capture detector operated at dc voltage to produce 1% full scale deflection [f.s.d.] for 1 ng of heptachlor epoxide when full scale deflection is 1 × 10⁻⁹ A.

| *Compound* | *Response (ng) (approx. amount for 1% f.s.d.)* | *Retention times relative to aldrin* | |
|---|---|---|---|
| | | *10% DC-200 column* | *10% DC-200/15% QF-1 (1:1) column* |
| Halowax 1099 | 25 | 0.45, 0.57, 0.65, 0.71, 0.79, 0.86, 0.91, 1.04, 1.14, 1.23, 1.34, 1.57, 1.88 | 0.49, 0.65, 0.73, 0.82, 0.96, 1.14, 1.26, 1.46, 1.62, 1.96 |
| Halowax 1014 | 25 | 0.45, 0.79, 1.04, 1.14, 1.57, 1.82, 2.85, 3.12, 3.55, 3.73, 6.0 | 0.49, 0.83, 1.15, 1.45, 1.61, 1.93, 2.80, 3.16, 3.68, 3.84, 5.9 |
| Halowax 1051 | 10 | 11.3 | 10.8 |

GC conditions are those used for organochlorine pesticide residue analysis[100]. The components of Halowax 1014 chromatograph throughout the retention time region of the organochlorine pesticides (retention times relative to aldrin, 0.5 to 6). Halowax 1099 exhibits peaks in the region of early eluting pesticides (retention times relative to aldrin, 0.5 to 2), and Halowax 1051 chromatographs beyond the retention times of common pesticides (retention time relative to aldrin, 11). Study of the Florisil column chromatographic clean-up[100] showed that Halowax 1014 and 1099 were completely eluted by 200 ml of 6% diethyl ether–petroleum ether. Halowax 1051 was found in both eluates from successive 200 ml elution with 6% and 15% diethyl ether–petroleum ether. A number of the organochlorine pesticides, including DDT and its analogs are also eluted by 6% diethyl ether–petroleum ether. The chromatographic procedure on silicic acid columns developed for separating PCB from organochlorine pesticides[43] was evaluated for the separation of the chlorinated naphthalenes from pesticides. Each of the Halowax samples (*e.g.*, 1014, 1051 and 1099) was completely eluted by 250 ml of petroleum ether eluent. (PCB's are also recovered in this eluate.) No Halowax was found in the acetonitrile–hexane–methylene chloride eluate, where most of the common organochlorine pesticides are recovered.

Fig. 77 shows gas chromatograms of brown trout extract fortified with 2.5 p.p.m. Halowax 1014, 0.3 p.p.m. *p,p'*-DDT and 0.2 p.p.m. *p,p'*-TDE and containing 0.19 p.p.m. *p,p'*-DDE residues before and after separation on silicic acid columns.

The identification and toxicological evaluation of chlorinated dibenzofuran and chlorinated naphthalene in two commercial polychlorinated biphenyls was described by Vos *et al.*[38]. In a previous study[104] a significant difference in toxicity was found between three commercial PCB preparations, high mortality, liver necrosis and clutch edema like lesions being associated with two of the compounds tested. In the study of Vos *et al.*[38] column chromatography and GLC demonstrated the presence of polychlorinated [illegible] diethyl ether fraction of each of two PCB

MONS 202145




Fig. 77. GLC curves of brown trout extract fortified with 2.5 p.p.m. Halowax 1014, 0.3 p.p.m. p,p'-DDT and 0.2 p.p.m. p,p'-TDE and containing 0.19 p.p.m. p,p'-DDE residue, before and after separation on a silicic acid column. (a) Before separation. (b) petroleum ether eluate from silicic acid column containing chlorinated naphthalene. (c) Polar eluate from silicic acid column containing (1) p,p'-DDE, (2) p,p'-TDE and (3) p,p'-DDT. 10-mg sample injected for each curve. GLC conditions same as those given in Table 35.

preparations which contained an average of 60% chlorine (Phenoclor DP6 and Clophen A60), and a chick embryo assay[109] confirmed the high toxicity of this fraction. The polar compounds were not found in the corresponding fraction of a third PCB (Aroclor 1260). The identity of the polar compounds which included tetra- and pentachlorodibenzofuran was confirmed by mass spectrometric and microcoulometric analyses.

Aliquots obtained from the column chromatography (Florisil) of the P[illegible]




Fig. 78. Gas chromatogram of the third fraction of Clophen A60 (showing a pattern identical [illegible] the third fraction of Phenoclor DP6. [illegible] in Table [illegible]



TABLE 36

RETENTION DATA AND COLLECTED MASS SPECTROMETRIC DATA OF PEAKS SHOWN IN FIG.

| *Peak No.* | *Relative retention*[a] | *Mass Nos.*[b] *and No. of chlorine atoms/molecule* | *Identity of compounds* |
|---|---|---|---|
| 1 | 1.40 | 304 (4 Cl) | Tetrachlorodibenzofuran |
| 2 | 1.58 | 332 (6 Cl) | Hexachloronaphthalene |
| 3 | 1.74 | 358 (6 Cl) | Hexachlorobiphenyl |
| | | 392 (7 Cl) | Heptachlorobiphenyl |
| 4 | 2.42 | 338 (5 Cl) | Pentachlorodibenzofuran |
| | | 392 (7 Cl) | Heptachlorobiphenyl |
| 5 | 3.48 | 366 (7 Cl) | Heptachloronaphthalene |

[a] Retention related to dieldrin = 1.00.
[b] Mass number calculations based on Cl = 35.

were analyzed by GLC employing a 204-1B Varian Aerograph gas chromatograph with electron capture detection. The 5 ft. × 1/8 in. Pyrex glass column was filled with 10% DC-200 on 80–100 mesh Gas-Chrom Q and operated at 200° with nitrogen as carrier gas at 50 ml/min. Mass spectrometry was used to identify compounds present in the polar fraction (25% diethyl ether-hexane) from the column fractionation of PCB's. For this purpose a CH 4 mass spectrometer (Varian MAT) was coupled to a Varian Aerograph type A550B oven, with a 5 ft. × 1/8 in. Pyrex glass column filled with 10% DC-200 on 80–100 mesh Gas-Chrom Q. The design of the interface between the above gas chromatograph and the mass spectrometer has been described by TENNOEVER DE BRAUW AND BRUNNEE[98].

High-resolution mass spectra were taken on a double-focus spectrometer type SM (Varian MAT). Exact mass measurements were made by the peak-matching method and measurements were also made from spectra which were recorded on photographic plates processed by a Varian MAT on a Leitz comparator SAM1.

Quantitative GLC information was obtained via microcoulometric analysis of the third polar Florisil column fractions using a Dohrmann microcoulometer coupled to a Microtek Model MT220 gas chromatograph with a 6 ft. × 1/4 in. glass column containing the same 10% DC-200 columns packing described above. The




Fig. 79. Mass spectrum of tetrachlorodibenzofuran (mass No. 304) obtained from the third fractions of Clophen A60 and Phenoclor DP6 (peak 1 in Fig. 78).

MONS 202146

...ed to 1 ml as described above and the hydrochloric acid produced by chlorinated compounds was detected and measured quantitatively on a Tracor 310 S gas chromatograph.

The microcoulometric gas chromatogram of the third fraction (25% diethyl ether hexane eluates) obtained with a 10-mm Florisil column is shown in Fig. 8 and reveals the presence of polar compounds in Phenoclor DP6 and Clophen A60. (Data for peaks 1–5 are given in Table 30.)

Fig. 79 shows the mass spectrum of tetrachlorodibenzofuran (mass No. 304) obtained from the third fractions of Clophen A60 and Phenoclor DP6 (peak 3 in Fig. 78). Fig. 80 depicts a mass spectrum of pentachlorodibenzofuran (mass No. 338) obtained from the third fractions of Clophen A60 and Phenoclor DP6 (peak 4 in Fig. 78). The spectrum of heptachlorobiphenyl (mol. wt. 392) is also present.




Fig. 80. Mass spectrum of pentachlorodibenzofuran (mass No. 338) obtained from the third fractions of Clophen A60 and Phenoclor DP6 (peak 4 in Fig. 78). The spectrum of heptachlorobiphenyl (mol. wt. 392) is also present.

The peak of the tetrachlorodibenzofuran (mass No. 304) was too masked by the appearance of the hexachloronaphthalene peak (peak No. 2, Fig. 78) to allow quantitative calculations. Similarly, peak No. 4 in the chromatogram (Fig. 78), found to be a mixture of heptachlorobiphenyl and pentachlorodibenzofuran, could not be measured directly. However, by assuming that the whole peak was produced by the dibenzofuran compound alone, a certain maximum level could be indicated. This level is found to be 5.4 ppm of pentachlorodibenzofuran in Clophen A60 and 20 ppm in Phenoclor DP6. The maximum level could be indicated with the estimated lowest limit of detection (1 ppm) of the mass spectrometric equipment.

The halogenated dibenzofurans are extremely toxic. Tri- and tetrachlorodibenzofuran in a single oral dose of 0.5–1.0 mg/kg caused severe and often lethal necrosis of the liver[110,111] and chloracne[108]. Table 37 shows the results of the chick embryo assay of the chlorinated biphenyls Aroclor 1260, Phenoclor DP6 and Clophen A60 and the 25% diethyl ether fraction of Clophen A60. The similarity between the mortality curves in groups 3 and 5 (Table 37) suggests strongly indeed that the polychlorinated dibenzofurans, the toxicity of the commercial PCBs. Chlorinated naphthalenes are far less toxic than chlorinated dibenzofurans[111] and the latter may thus be assumed to determine the toxicity of the third fraction. The occasional occurrence of hydropericardia in chicks fed Aroclor[39] may indicate the presence of a very small quantity of a toxic factor in this preparation.



TABLE 37

CHICK EMBRYO ASSAY OF THE CHLORINATED BIPHENYLS AROCLOR 1260, PHENOCLOR DP6, AND CLOPHEN A60 AND THE 25% DIETHYL ETHER FRACTION (THE THIRD FRACTION) OF CLOPHEN A60

| Test compound | Group No. | Dose* (mg/egg) | No. of eggs treated | No. of infertile eggs | Percentage hatch (% of fertile eggs) | Remarks |
|---|---|---|---|---|---|---|
| Aroclor 1260 | 1 | 3.5 | 15 | 0 | 80 | |
| Phenoclor DP6 | 2 | 3.5 | 15 | 0 | 0 | Most embryos died during first half of incubation period. |
| Clophen A60 | 3 | 3.5 | 20 | 1 | 5 | |
| Clophen A60 (third fraction) | 4 | 3.5 | 15 | 0 | 0 | All embryos died during first four days of incubation period. |
| | 5 | 3.5 | 15 | 0 | 7 | Most embryos died during first half of incubation period. |
| | 6 | 0.35 | 15 | 2 | 92 | |
| | 7 | 0.035 | 15 | 2 | 100 | |
| – | 8 | 0 | 20 | 2 | 94 | Ethanol control |
| – | 9 | — | 20 | 0 | 90 | Untreated control |

* The doses, each dissolved in 0.05 ml of ethanol, were injected into the air cell of fresh eggs before incubation.

For example PCB can react with sodium hydroxide at elevated temperatures to yield phenolic compounds which may be saponified to polychlorohydroxybiphenyls by sodium hydroxide in a polyhydric alcohol medium[112]. Dibenzofuran derivatives could then be formed via a further loss of hydrochloric acid.

## VIII. SUMMARY

The chromatographic (column, thin-layer and gas–liquid chromatography and combined gas–liquid chromatography and mass spectroscopy) and biological aspects of the polychlorinated biphenyls have been reviewed with primary focus as to their utility, ecological distribution and toxicity as well as diverse techniques for their separation from chlorinated pesticides and subsequent analysis.

## REFERENCES

1 *Monsanto Tech. Bull. (PP) 9A*. Aroclor Physicals.
2 J. H. Koeman, M. C. Ten Noever de Brauw and R. H. de Vos, *Nature, Lond.* 221 (1969) 1126.
3 G. L. Baughman, W. L. Bowers and J. Swineford, *J. Ass. Off. Agr. Chem.* 54 (1971) 1.

MONS 202147

4 F [illegible] and W N Sullivan *J Econ Entomol* 46 (1953) 937
5 C [illegible] Sullivan and I Bornstein *J Econ Entomol* 46 (1953) 58
6 [illegible] and T T Liang *J Econ Entomol* 62 (1969) 791
7 S Jensen *New Scientist* 32 (1966) 612
8 R W Risebrough in Reiche D B Pearsall S G Hervan and M N Kirven *Nature* 210 (1966) 1095
9 D C Holmes J H Simmons and J O G Tatton *Nature* 216 (1967) 227
10 A V Holden and K Marsden *Nature* 216 (1967) 1274
11 D W Anderson J J Cooley R W Risebrough D F Hughes and R E Christiansen *Can Field Nat* 84 (1970) 89
12 L G Reynolds *Biochem Soc* 31 (1971) 27
13 S Jensen A G Johnels A Olsson and G Otterlind *Nature* 224 (1969) 247
14 I J Bross A C Walker and A Meidrey *Bull Environ Contam Toxicol* 5 (1970) 347
15 R Risebrough and V Brodine *Environment* 12 (1970) 16
16 G Westöö K Noren and M Anderson *Var Föda* 2–3 (1970) 10
17 D B Pearsall and J L Lincer *BioScience* 20 (1970) 958
18 L Greenberg M R Myers and A R Smith *J Ind Hyg Toxicol* 21 (1939) 29
19 L Schaffer *J Amer Med Ass* 122 (1943) 158
20 L Schwartze and F A Barlow *US Pub Health Rep* 57 (1942) 1747
21 L Schwartze and S M Peck *NY State Med J* 43 (1943) 1711
22 J W Jones and H S Alden *Arch Dermatol Syphilol* 33 (1936) 1021
23 H von Weidel W A Holla and J. Denton, *Rubber Age (New York)* 53 (1943) 401
24 I N Sax *Dangerous Properties of Industrial Materials* 2nd Ed., Reinhold Pub Corp New York 1935 p 396
25 J W Miller *US Pub Health Rep*, 39 (1924) 1085
26 C K Drinker M F Warren and G A Bennett, *J Ind Hyg Toxicol* 19 (1937) 283
27 G A Bennett C K Drinker and M F Warren, *J Ind Hyg Toxicol* 20 (1938) 97
28 M Nishizumi *Arch Environ Health* 21 (1970) 620
29 E L McCune J E Savage and B L O Dell, *Poultry Sci* 41 (1962) 295
30 D J Flick C D Douglass and L Galld, *Poultry Sci* 42 (1963) 855
31 J McLaughlin Jr J P Marliac M J Verrett, M K Mutchler and O G Fitzhugh *Toxicol Appl Pharmacol* 5 (1963) 760
32 J C Street F M Urry D G Wagstaff and V D Beau, *Amer Chem Soc Meet New York Sept 8–12 1969*
33 D Presst D J Jefferies and N W Moore *Environ Pollution* 2 (1970) 3
34 J G de Vos and J H Koeman *Toxicol Appl Pharmacol*, 17 (1970) 656
35 *FDA Status Rept Chemistry and Toxicology of Polychlorinated Biphenyls or Aroclors* June 1970
36 J Bitman and H C Cecil *J Agr Food Chem* 18 (1970) 1108
37 D B Peakall *Nature* 216 (1967) 505
38 J L Lincer and D B Peakall *Nature* 228 (1970) 783
39 D C Villeneuve D L Grant W E J Phillips, M L. Clark and D J Clegg *Bull Environ Contam Toxicol* 6 (1971) 120
40 K Vermeer and L M Reynolds, *Can Field Nat*, 84 (1970) 117
41 D B Peakall *Bull Environ Contam Toxicol* 6 (1971) 100
42 M Friend and D O Trainer *Science* 170 (1970) 1314
43 C L Litterst and E P Lichtenstein *Arch Environ Health* 22 (1971) 454
44 L M Reynolds *Bull Environ Contam Toxicol* 4 (1969) 128
45 J A Armour and J A Burke *J Ass Offic Anal Chem* 53 (1970) 761.
46 B M Mulhern, E Cromartie W L Reichel and A A Belisle, *J Ass Anal Chem* 54 (1971) 548
47 H L Faller *J Amer Chem Soc* 67 (1945) 1591
48 W L Reichel T G Lamont and E Cromartie *Bull Environ Contam Toxicol*, 4 (1969) 24
49 J Arnold and J Burke *FDA Lab Inform Bull No 918* (1969) 11
50 N V Feiringer and J F Westfall *J Chromatogr* 57 (1971) 397
51 O Hutzinger S Safe and V Zitko *Bull Environ Contam Toxicol* 6 (1971) 269
52 C G Gustafson *Environ Sci Technol* 4 (1970) 814
53 S Jensen and G Widmark *Rept OECD Pesticide Conf Unintended Residues in the Environment, Stockholm 1967*
54 R W Risebrough P Reiche and H S Olcott, *Bull Environ Contam Toxicol* 4 (1969) 192
55 H Evans *Analyst (London)*, 87 (1962) 569
56 M J de Albert Mallett H Egan E W Godly E W Hammond J Roburn and J Thomson *Analyst (London)* 89 (1964) 197
57 J F Simmons and J O G Tatton *J Chromatogr* 27 (1967) 153
58 W C Krantz B M Mulhern G E Bagley A Sprandt F J Ligas and W B Robertson Jr *Pestic Monit J* 4 (1970) 136
59 R W Risebrough in M W Miller and G G Berg (Editors) *Chemical Fallout* Thomas Springfield Ill 1969 p 5.
60 R W Risebrough D B Menzel D J Martin Jr and H S Olcott *Nature*, 216 (1967) 589
61 F Ferro A Bevenue and H Beckman *Bull Environ Contam Toxicol* 2 (1967) 373
62 A V Holden *Nature* 228 (1970) 1220
63 A V Holden and K Marsden *J Chromatogr* 44 (1969) 481
64 D J Hansen, P R Parrish, L E Lowe A J Wilson Jr and P D Wilson *Bull Environ Contam Toxicol*, 6 (1971) 113
65 P A Mills J F Onley and R A Gaither *J Ass Offic Agr Chem*, 46 (1963) 186
66 H F Enos, personal communication in ref 64
67 D J Hansen and A J Wilson *Pestic Monit J* 4 (1970) 51
68 J E Keil, L E Priester and S H Sandifer *Bull Environ Contam Toxicol* 6 (1971) 136
69 T W Duke, J I Lowe and A J. Wilson Jr *Bull Environ Contam Toxicol*, 5 (1970) 171
70 S Lofstrand A Södergren and J Rabol, *Nature* 232 (1971) 467
71 G Westöö and K Noren *Acta Chem Scand* 24 (1970) 1039
72 K Noren and G Westöö, *Acta Chem Scand*, 22 (1968) 2187
73 M F Kovacs, *J Ass Offic Agr Chem* 46 (1963) 884
74 D L Grant, W E J Phillips and D C Villeneuve *Bull Environ Contam Toxicol*, 6 (1971) 202
75 J A Burke and W Holswade *J Ass Offic Anal Chem*, 49 (1966) 547
76 M Kunitsune *Fukuoka Acta Med* 62 (1969) 513
77 T Kojima, H Fukumoto and N Makisumi *Jap J Legal Med* 25 (1971) 415
78 L C Mitchell, *J Ass Offic Agr Chem* 40 (1957) 999
79 K R Tarrant and J O G Tatton *Nature* 219 (1968) 725
80 S Jensen A Jernelov R Lange and K H Palmork *Fir WG/70/E 83 FAO Technical Conf on Marine Pollution, Rome Dec 9–18 1970*
81 B Ahling and S Jensen *Anal Chem* 42 (1970) 1483
82 T T Schmidt, R W Risebrough and F Gress, *Bull Environ Contam Toxicol* 6 (1971) 255
83 R I Stanley and H T LeFavour *J Ass Offic Anal Chem* 48 (1965) 666
84 V Zitko *Bull Environ Contam. Toxicol* 5 (1970) 279
85 H. Weingarten W D Ross J M Schlabper and G G Wheeler, *Anal Chim Acta* 26 (1962) 391
86 N L. Gregory *J Chem Soc (B)* (1968) 295
87 E. M Emery and G M Gasser *US Pat.*, 3,520 108 July 14 1970
88 V Zitko, O Hutzinger and S Safe, *Bull Environ Contam Toxicol*, 6 (1971) 260
89 R H de Vos and E W Peet, *Bull Environ Contam Toxicol*, 6 (1971) 164
90 R I Asai F A Gunther W E Westlake and Y Iwata *J Agr Food Chem* 19 (1971) 396
91 R I Asai F A Gunther and W E Westlake, *Residue Rev*, 19 (1967) 57
92 M Beroza and R Sarmiento *Anal. Chem*, 36 (1964) 1744
93 P Mills *J Ass Offic Anal Chem*, 42 (1959) 734
94 J H Beynon, *Mass Spectrometry and Its Applications to Organic Chemistry* Elsevier New York, 1960 p 298
95 S W Bellman and T L. Barry, *J Ass Offic Anal Chem* 54 (1971) 490
96 S. P Markey *Anal Chem*, 42 (1970) 396
97 *Pesticide Analytical Manual*, Food and Drug Administration Washington 1969, Sect 211
98 M C Tenaglever deBrauw and C Brunner, *Anal Chem*, 220 (1967) 321
99 D F Flick R G O Dell and V. A. Childs, *Poultry Sci* 44 (1965) 1460
100 R Rymage, *Anal Chem*, 36 (1964) 739
101 B M Mulhern *J Chromatogr*, 34 (1968) 536.
102 O Hutzinger W D Jamieson and V Zitko, *Nature*, 226 (1970) 664
103 W D Jamieson and F G Mason, *Res Dev. Technol* in press.
104 *Halowax Chlorinated Naphthalenes* Koppers Co., Pittsburgh Pa.
105 J A Armour and J A Burke, *J Ass. Offic. Anal. Chem* 54 (1971) 175
106 *Pesticide Analytical Manual*, Vol I, Food and Drug Administration, Washington 1969




*J Chromatogr* 68 (1972) 345–426

MONS 202148

107 J. G. VOS, J. H. HOFMAN, H. C. VANDERMAAS, M. C. TENNOEVER-DEBRAUW AND R. H. DE VOS, *Food Cosmet. Toxicol.*, 8 (1970) 625.
108 J. G. VOS AND J. H. HOLMAN, *Toxicol. Appl. Pharmacol.*, 17 (1970) 656.
109 M. VERRETT, J. P. MARIAC AND J. MCLAUGHLIN, *J. Ass. Offic. Anal. Chem.*, 47 (1964) 1003.
110 H. BAUER, K. H. SCHULZ AND G. SPIEGELBERG, *Arch. Gewerbepathol. Gewerbehyg.*, 18 (1961) 518.
111 H. J. HOFMANN, *Arch. Exp. Pathol. Pharmakol.*, 232 (1958) 228.
112 J. W. J. CAY AND J. M. RICHARDS, *Office Tech. Serv. P. B. Rept.* 75850, *Bus. Final Rept.* (1947) 803.
113 PROGIL S.A., *Brit. Pat.*, 779 221, July 17, 1957.



MONS 202149