

Please return to:
Mr. Homer P. Wheeler
Monsanto Chemical Co.
800 N. Lindbergh Blvd.
St. Louis 68, Missouri

The Toxicity of the Fogs Formed
by Dropping Pydraul F-9,
Aroclor 1248 and Tricresyl
Phosphate, Upon the Surface
of a Heated Inconel
(Project 49)

by
Treon, Cleveland, Shaffer, Cappel,
Gahegan, Wagner
March 11, 1953

PLAINTIFF'S
EXHIBIT

1156

NEV 016470

715722

Table of Contents

	Page
Scope	1
Conclusions	1
Materials	5
Experimental Methods	7
Equipment	7
Volatilization of the Materials Without Decomposition	7
Generation of a Fog From a Heated Inconel Surface	9
Sampling and Analysis of Contaminated Air . .	11
Experimental Results	19
Exposure to Undecomposed Vapors	19
Exposure to Products Arising From Pydraul F-9, Arcolor 1248, or Tricresyl Phosphate When in Contact with Inconel at 1250° or 1050° F.	21
Mortality	21
Signs and Pathology of Intoxication	23
The Concentrations of Identified Decomposition Products Formed at the Higher and Lower Temperatures	25
Discussion	28
Supplementary Statement - The Application of Pydraul F-9, Arcolor 1248 and Tricresyl Phosphate Upon the Skin of Rabbits	31
Experimental Methods	31
Experimental Results	32

CONFIDENTIAL

Sequence of Tables

1. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to the Vapor of Pydraul P-9 in Air.
 2. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to the Vapor of Arcolor 1248 in Air.
 3. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to the Vapor of Tricresyl Phosphate in Air.
 4. Concentrations of Various Decomposition Products in Air of Chamber in the Several Experiments.
 5. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to Aerosols Formed by Dropping Pydraul P-9 Into an Inconel Tube Heated to a Temperature of 1250° F. (677° C.).
 6. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to Aerosols Formed by Dropping Pydraul P-9 Into an Inconel Tube Heated to a Temperature of 1050° F. (566° C.).
 7. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to Aerosols Formed by Dropping Arcolor 1248 Into an Inconel Tube Heated to a Temperature of 1250° F. (677° C.).
 8. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to Aerosols Formed by Dropping Arcolor 1248 Into an Inconel Tube Heated to a Temperature of 1050° F. (566° C.).
 9. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to Aerosols Formed by Dropping Tricresyl Phosphate Into an Inconel Tube Heated to a Temperature of 1250° F. (677° C.).
 10. The Mortality, Length of Survival, and Changes in Weight of Animals Exposed to Aerosols Formed by Dropping Tricresyl Phosphate Into an Inconel Tube Heated to a Temperature of 1050° F. (566° C.).
- NEV 016472