



CHEMICAL MANUFACTURERS ASSOCIATION

June 1, 1995

Dear Vinyl Chloride Health Committee Members:

A draft protocol for the vinyl chloride combined 2-generation and developmental toxicity study prepared by Bill Breslin of Dow is enclosed for your review. Please bring your comments on the draft to the next Committee meeting scheduled for June 7 from 10:00 a.m. to 3:30 p.m. at the CMA offices in Washington, D.C. If you are unable to attend the meeting, please send your written comments to me as soon as possible. Please call me at (202) 887-1192 if you have any questions. I look forward to seeing you next week.

Sincerely,

A handwritten signature in cursive script, appearing to read "Has Shah", is written in dark ink.

Hasmukh C. Shah, Ph.D.
Manager, Vinyl Chloride Panel

Enclosure

BFG42852



THE DOW CHEMICAL COMPANY

HEALTH AND ENVIRONMENTAL SCIENCES

1803 Building

Midland, Michigan 48674

FAX Number: 517/638-9863

FAX TRANSMITTAL SHEET

Date: 6/1/95

To: Dr. Has Mukh Shah

Company Name: CMA

Fax Number: 202-887-5427

From: William J. Breslin

Phone: 517-636-1311

Number of Pages: 23
(including this page)

Material sent by: W.J. Breslin

Phone: 517-636-1311

Comments:

Has Mukh, please find in the subsequent pages of this FAX the draft protocol for the vinyl chloride combined 2-generation and developmental toxicity study. I look forward to your comments following the Panel's review. At this time it is unlikely that I will be able to make the Panel meeting next week. I was also unable to reach J. Ramlow as he is still ill and out of the office.

Bill

If you do not receive all material being transmitted, please call the sender listed above.

Updated 8/89
clatton

BFG42853

BFG 01888

HEALTH AND ENVIRONMENTAL SCIENCES
THE DOW CHEMICAL COMPANY

PROTOCOL

THE TOXICOLOGY RESEARCH LABORATORY,
1803 BUILDING, MIDLAND, MICHIGAN 48674

TITLE: VINYL CHLORIDE: COMBINED INHALATION TWO-GENERATION
REPRODUCTION AND DEVELOPMENTAL TOXICITY STUDY IN CD RATS

DATE:

FILE NUMBER:

PROPOSED EXPERIMENTAL
START DATE:

PROJECT NUMBER:

TASK NUMBER:

PROPOSED EXPERIMENTAL
TERMINATION DATE:

OSD NUMBER:

COST CENTER:

ESTIMATED DATE
FINAL REPORT:

SPONSOR:

CAS NUMBER:

SIGNATURES:

STUDY DIRECTOR:

/DATE

COINVESTIGATOR(S):

/DATE

INHALATION SPECIALIST:

/DATE

STUDY PATHOLOGIST:

/DATE

APPROVAL(S):

/DATE

SPONSOR:

/DATE

DISTRIBUTION -- SEE ATTACHED LIST

BFG42854

BFG 01889

MATERIALS AND METHODS

The test material used for this study was obtained from X(name), X(city), X(state) and was identified as (Lot number of the test material). The purity of the test material was reported to be xx.x% (reference). A sample of the test material will be taken and stored in a manner consistent with the reference sample retention policy of this laboratory. The test material will be reanalyzed at approximately 6-month intervals to confirm purity and stability.

The test substance has the following properties:

Chemical Name:

Synonyms:

Molecular Formula:

Molecular Weight:

Structures:

Appearance:

Vapor Pressure:

Saturated Atmosphere:

Vapor Density:

Flash Point:

Boiling Point:

Specific Gravity:

Conversion Factors:

Test Species and Husbandry.

Male and female CD rats (Charles River Breeding Laboratory, Kingston, NY or Portage, MI) approximately four weeks of age will be purchased for the exposure phase of this study. An additional group of male CD rats (Charles River Breeding Laboratory, Kingston, NY or Portage, MI) approximately ten weeks of age will be purchased for breeding stock for the developmental portion of this study and will not be exposed to the test material. Animals

DRAFT

BFG 01891

BFG42856

will be ordered to ensure that a sufficient number of animals of acceptable health and weight are available to conduct the study as designed. This strain of rat has been selected because of its general acceptance and suitability for toxicity testing and the availability of a reliable commercial source. Upon arrival at the laboratory¹, all rats will be examined for health status by a veterinarian and acclimated to the laboratory environment for approximately two weeks, according to the Standard Operating Procedures of the Reproductive Toxicology Group. For the randomization procedure, the rats will be weighed and ranked according to body weight and those from the extremes of the distribution will be identified and removed from the population until only the number of animals required for the study remain. These rats will be randomly assigned by weight to the treatment groups to increase the probability of uniform group mean weights and standard deviations at the initiation of the study. Rats not placed on test will be removed from the test room and the disposition of these animals will be documented in the study file.

Identification of all rats on test will be accomplished by inserting a uniquely coded alphanumeric metal tag in one ear of each rat. In the event that an ear tag becomes dislodged during the course of the study, it will be replaced with one having the same alphanumeric code (i.e., a new alphanumeric code will not be assigned) and noted in the study file.

Rats will be housed singly in wire mesh, stainless steel cages in racks provided with deotized cage board to minimize odor and aid in maintaining a clean environment. Prior to and following daily exposures during late gestation and throughout lactation, females will be housed in plastic nesting boxes provided with ground corn cob nesting material (further details provided in experimental design section). The animal rooms of the facility are designed to maintain humidity at approximately 40-60%, temperature at approximately 22 °C, photoperiod at 12 hrs light:12 hrs dark and air flow at 12-15 changes/hour. A feed crock and a pressure-activated stainless steel water

¹Fully accredited by the American Association for Accreditation of Laboratory Animal Care (AAALAC).

DRAFT

BFG42857

BFG 01892

nipple will be components of all cages. A basal diet of Purina Certified Rodent Chow No. 5002 (Purina Mills Inc., St. Louis, MO) will be provided *ad libitum* except during the six hour per day exposures, at which time feed will be withheld. Municipal drinking water will be available *ad libitum* throughout the prestudy and study periods. Analysis of the chow will be performed by Purina Mills Inc. to confirm that the diet provides adequate nutrition, and to quantify the levels of selected contaminants associated with the formulation process. Drinking water obtained from the City of Midland, MI will be analyzed for chemical parameters and biological contaminants by the City of Midland Water Department. In addition, specific analyses for chemical contaminants will be conducted at periodic intervals as stated in the Standard Operating Procedures of The Toxicology Research Laboratory, The Dow Chemical Company.

Exposure Chamber.

Animals will be housed and exposed to test material vapors in a 14.5 m³ chamber (2.4 m wide x 2.4 m high x 2.4 m deep with a pyramidal top) under dynamic airflow conditions. Chamber airflows will be maintained at approximately 2900 liters/minute, which is sufficient to provide the normal concentration of oxygen to the animals.

Vapor Generating System.

The various concentrations of test material will be generated using a glass J-tube method (Miller et al., 1980). Liquid test material will be metered into a glass J-tube assembly and vaporized by a preheated stream of compressed air (up to 100 l/min) passing through the J-tube. Compressed air will be heated to the minimum extent necessary to facilitate complete vaporization of the test material. The compressed air and test material vapors will be diluted and mixed with room air to achieve the desired concentration of test material vapors at a total flow rate of 2900 l/min.

Chamber Monitoring.

The concentration of the test material in each chamber will be measured at least once per hour using a MIRAN 1A infrared spectrophotometer (Foxboro

DRAFT

BFG42858

BFG 01893

Analytical, Norwalk, CT). Analytical equipment will be calibrated using standards having a known test material vapor concentration contained in 90 liter SARAN^{*} film gas bags prior to the first exposure and at least monthly thereafter. Daily checks of the analytical equipment will be performed prior to each exposure period using a single test material standard concentration. In addition, the amount of test material used each day will be recorded and the nominal concentrations (amount of the test material used/total chamber airflow) of test material will be calculated. Prior to the start of the study, each of the chambers to be used will be checked to ensure that a uniform distribution of vapors occurs within the breathing zone according to Standard Operating Procedures.

Airflow through each chamber will be determined at hourly intervals using a differential pressure transducer (Model C264, Setra Systems, Inc., Acton, MA) or a Universal Venturi tube (Series 180, BIF, 345 Harris Ave., Providence, RI). The manometer and differential pressure transducer will be calibrated with a gas meter (Singer Aluminum Diaphragm Meter, Model AL-1400, American Meter Division, Philadelphia, PA) prior to the start of the study. The Universal Venturi tubes have been calibrated at the factory.

Chamber temperatures will be measured with a thermometer or resistance temperature device (RTD) and relative humidities will be measured with relative humidity gauges or humidity sensors (HMP 112A, Vaisala, Helsinki, Finland) at least once each hour. Calibration of the hygrometers, RTD's and humidity sensors will be documented in the study file. The temperature and relative humidity in each chamber will be controlled by a system designed to maintain temperature at approximately $22 \pm 2^{\circ}\text{C}$ and relative humidity at approximately 40-60%.

Output from the differential pressure transducer, RTD, humidity sensor and infrared spectrophotometer will be collected by the CAMILE^{*} Data Acquisition and Control System.

* Trademark of The Dow Chemical Company

* Trademark of SAGIAN Indianapolis, Indiana.

DRAFT

BFG42859

BFG 01894

Exposure Concentrations.

Rats will be exposed to target concentrations of 0, X, Y or Z ppm of the test material. These exposure levels correspond to an oral equivalent dose of approximately 0, X, Y and Z mg/kg/day assuming ventilation rates of 1 l/min/kg, 100 percent absorption and a 6 hour/day exposure. The chosen concentrations were selected by the sponsor and were based upon the results of the inhalation toxicity studies previously discussed.

Reproduction - Experimental Design.

Groups of 30 male and 30 female rats will be exposed to 0, X, Y or Z ppm of the test material via inhalation, for 6 hours/day, 5 days/week prior to mating and 6 hours/day, 7 days/week during mating, gestation and lactation. The overall chronology of events and study design for this study are depicted in Tables 1 and 2, respectively. The treatment of the first generation parental (P1) rats will begin at approximately 6 weeks of age. After approximately 10 weeks of exposure (5 days/week, excluding holidays), P1 rats will be mated (one male to one female of the respective treatment group) to produce the F1 litters. Following weaning (3 weeks of age) of the F1 litters, 30 males and 30 females from each treatment group will be randomly selected and assigned to the respective treatment group to become the second generation parents (P2). After approximately 10 weeks of treatment following weaning of the last F1 litter, the P2 adults will be bred to produce the F2 litters. Exposures of P1 and P2 adults rats to the test material will continue until the adults are sent to necropsy. All rats will be housed continuously in exposure chambers following the initial exposure to the test material, except during late gestation and throughout lactation periods, when female rats will be housed outside of the exposure chambers. Maternal rats will not be exposed to the test material after day 20 of gestation (as calculated from day 0 of gestation via sperm-positive vaginal lavage) through the fourth day postpartum, in order to allow for parturition and initiation of lactation. During the lactation period, pups will not be placed in the exposure chambers, but will remain in the nesting boxes separated from the dam for approximately 7 hours/day on lactation days 5 through 21.

DRAFT

BFG42860

BFG 01895

Reproduction - Breeding Procedure

Breeding of the P1 and P2 adults will commence after approximately 10 weeks of treatment. Each female will be placed with a single male from the same dose level (1:1 mating) until pregnancy occurs or two weeks have elapsed. During each breeding period, daily vaginal lavage samples will be evaluated for the presence of sperm as an indication of mating. The day on which sperm are detected or a vaginal plug is observed *in situ* will be considered day 0 of gestation. Sperm- and plug-positive females will then be separated and placed back into wire mesh, stainless steel cages. If mating has not occurred after two weeks, the animals will be separated without further opportunity for mating. For the P2 mating, cohabitation of male and female litter mates will be avoided.

Reproduction - Culling and Weaning

To reduce the variation in the growth of the pups, the F1 and F2 litters with a total number of pups exceeding eight will be culled on day 4 postpartum. Culled litters will be reduced to a total of eight pups, four males and four females, if possible. Pups to be culled will be selected using a computer generated randomization procedure. Litters with eight or fewer pups will not be culled. Preferential culling of runts will not be performed. Culled pups will be examined grossly for abnormalities and euthanized by the deposition of sodium pentobarbital (Succumb Butler, Columbus, OH) into the oral cavity. Weaning of all litters will be done 21 days after delivery. Weanlings not held for prospective generations or selected for necropsy will be examined grossly for abnormalities and euthanized by CO₂ inhalation.

Reproduction - Physical Observations

Each rat on study will be observed twice daily (a.m. and p.m.) for mortality, morbidity and moribundity as well as availability of feed and water. Changes in behavior or demeanor and indications of overt toxicity will be evaluated during the a.m. or p.m. observation. In addition, a thorough clinical examination will be conducted on all animals prior to the start of the study and weekly thereafter. This examination will include thorough evaluations

DRAFT

BFG42861

BFG 01896

of the skin and fur, mucous membranes, respiration, nervous system and behavior pattern. All adult rats found dead or in moribund condition will be submitted for a gross pathologic examination. Adult rats found dead after normal working hours will be refrigerated until a necropsy can be performed. All pups found dead or pups that are euthanized in moribund condition will be examined to the extent possible for defects and/or cause of death and preserved in neutral, phosphate-buffered 10% formalin. Cannibalized pups will be examined to the extent possible and discarded.

Reproduction - Body Weights and Feed Consumption

All P1 animals will have body weights and feed consumption (optional) recorded weekly during the 10-week pre-breeding treatment period, beginning on or before the first week of the study. Body weights for males will be recorded weekly throughout the course of the study. Sperm and plug positive females will be weighed on Days 0, 7, 14 and 21 of gestation. Females that deliver litters will be weighed on Days 1, 4, 7, 14, and 21 of lactation. During breeding, feed consumption will not be measured in males or females due to cohousing. Following completion of the breeding periods, weekly feed consumption again will be measured in males. During gestation, feed consumption will be measured at weekly intervals in sperm and plug positive females. After parturition, feed consumption will be measured twice during the first and second week of lactation and at 2 - 3 day intervals during the last week of lactation. A similar schedule will be followed for the P2 generation.

Reproduction - Litter Data

All litters will be examined as soon as possible after delivery. The following parameters will be recorded for each litter: total litter size on the day of parturition (day 0), the number of live and dead pups on days 0, 1, 4, 7, 14, and 21 postpartum, and the sex and the weight of each pup on days 1, 4 (before and after culling), 7, 14, and 21 of lactation. Any visible physical abnormalities or demeanor changes in the neonates will be recorded during the lactation period.

DRAFT

BFG42862

BFG 01897

Reproduction - Physical Maturation Landmarks

All F1 weanlings selected for mating will be observed daily for vaginal opening beginning on postnatal day 30 (Adams *et al.*, 1985) or preputial separation beginning on day 35 (Korenbrod *et al.*, 1977). If there is a treatment-related effect observed on the F1 sex ratio, age of vaginal opening or age of preputial separation, then anogenital distance will be measured on post natal day 4 for all F2 pups.

Reproduction - Estrous Cycling

Estrous cycle length and normality will be evaluated daily by vaginal lavage (Cooper *et al.*, 1993) for all P1 and P2 females starting three weeks prior to mating and continuing throughout cohabitation.

Reproduction - Pathology - Adult Rats

A complete necropsy will be conducted by a team of trained individuals under the direct supervision of a veterinary pathologist on all P1 and P2 adults. The scheduled necropsy will be performed after the last litter of the respective generation has been weaned. Adult males will be fasted overnight, anesthetized with methoxyflurane and euthanized. Adult females will be necropsied on day 2 of diestrus whenever possible. This will be accomplished by monitoring (by vaginal lavage) for the occurrence of at least one estrous cycle after which time females found to be in day one of diestrus will be fasted overnight and necropsied on the following morning. The expected, subsequent stage of the estrous cycle (day 2 of diestrus) will be confirmed by vaginal lavage on the day of necropsy, prior to euthanasia. Based upon the results of these smears, exclusion of appropriate data parameters used for statistics will be performed for females not found to be in the appropriate stage of the estrous cycle (day 2 of diestrus) on the day of necropsy. The fasted females will be euthanized as described for the males. The eyes of both males and females will be examined *in situ* by gently pressing a moistened glass slide against the cornea and observing the eyes under fluorescent light. The uteri of all cohabited females will be examined for the presence and number of implantation sites. Tissues routinely collected (Table 3) will be saved from

DRAFT

BFG 01898

BFG42863

these rats and preserved in neutral, phosphate-buffered 10% formalin, with the following exceptions. The testes and epididymides will be preserved in Bouin's fixative. The lungs will be infused with formalin to their approximate normal inspiratory volume. The nasal cavity will be flushed with formalin via the pharyngeal duct to ensure rapid fixation of the tissue. Moribund rats and those dying spontaneously will be necropsied in a similar manner. However, body and organ weights will not be recorded.

Reproduction - Organ Weights - Adult Rats

The following organs of all P1 and P2 parental animals will be weighed: uterus, ovaries, testes, single epididymis (total and cauda), seminal vesicles (with coagulating glands and their fluids), prostate, brain, liver, kidneys, lungs, adrenal glands, spleen, and thymus, and the organ-to-body weight ratios calculated.

Reproduction - Histology - Adult Rats

Histologic examination of potential target organs and reproductive tissues (Table 3) will be performed on the control and high dose groups. Examination of tissues from the low and middle groups will be limited to those tissues which demonstrate treatment-related histologic changes in the high dose group. Only the right ovary will be routinely processed for standard microscopic examination. The left ovary will be saved for possible oocyte quantification. If deemed necessary by the study sponsor, oocyte quantification will include evaluation of a minimum of ten sections, randomly selected from one completely sectioned ovary per female of the high-dose and control groups. Ovarian follicles will be placed into one of three categories as described by Plowchalk et al., (1993). The total number of follicles and the number of follicles in each of the three categories will be evaluated. Ovaries from the low and middle dose groups may be evaluated if treatment-related changes are observed in the high dose group. A complete set of tissues (excluding the left ovary for females), encompassing all organs listed in Table 3, will be prepared from all rats dying spontaneously or euthanized in a moribund condition and examined in an attempt to determine cause of death.

DRAFT

BFG42864

BFG 01899

Reproduction - Sperm Count, Motility and Morphology

For all P1 and P2 males at termination, samples of sperm from the distal cauda epididymis or the proximal vas deferens will be collected for evaluation of percent progressively motile sperm and possible evaluation of sperm morphology. The entire right cauda epididymis will be weighed and then minced in saline to enumerate the total number of sperm (cauda reserves). Sperm motility and count will be determined with the use of the Hamilton-Thorn (HTM) Integrated Visual Optical System (IVOS) motility analyzer (Hamilton-Thorn Research, Beverly, Massachusetts). All samples for motility analyses will be video recorded or recorded digitally on disk and the recording kept as raw data. Sperm samples will be prepared for morphological evaluation and saved, but will not be evaluated unless deemed necessary by the study sponsor.

Reproduction - Pathology - Weanling Rats

At the time of weaning, 1 pup/sex/litter/dose from the F1 and F2 litters will be randomly selected for a complete necropsy by a team of trained individuals under the direct supervision of a veterinary pathologist. In order to control for variation in body and organ weight, all F1 and F2 pups selected for a complete necropsy will be euthanized at the same age. Pups will be anesthetized with methoxyflurane and euthanized. Terminal body weights will be recorded. Gross pathologic examination and preservation of tissue samples (Table 3) will be performed as described above for adults.

Reproduction - Organ Weights - Weanling Rats

For all F1 and F2 pups that are examined macroscopically (one/sex/litter), the following organs will be weighed: ovaries, testes, brain, liver, kidneys, adrenal glands, spleen and thymus.

Reproduction - Histology - Weanling Rats

Organs that demonstrate treatment-related effects in weanlings will be examined microscopically in the control and high-dose groups. Examination of tissues from the low and middle groups will be limited to those tissues

DRAFT

BFG 01900

BFG42865

which demonstrate treatment-related histologic changes in the high dose group. Microscopic examination will also be made of all tissues showing gross pathologic changes.

Developmental - Experimental Design.

Additional groups of 30 female rats will be exposed to 0, X, Y or Z ppm of the test material via inhalation, for 6 hours/day, 5 days/week prior to mating and 6 hours/day, 7 days/week during mating and gestation. The overall chronology and study design for the developmental portion of this study is depicted in Tables 1 and 2, respectively. Treatment of the rats will begin at approximately 6 weeks of age. After approximately 10 weeks of exposure (5 days/week, excluding holidays), the females will be mated with unexposed stock males, approximately 12 weeks of age. On gestation day 21 the females will be euthanized and maternal and developmental parameters evaluated.

Developmental - Breeding Procedure

Breeding of the adults will commence after approximately 10 weeks of treatment and continue until pregnancy occurs or two weeks has elapsed. Each female will be placed with a single unexposed stock male following the daily exposure period and left with the male until the following morning. In the morning prior to exposure, each female will be evaluated for the presence of sperm via vaginal lavage, and returned to her exposure cage. The day on which sperm are detected or a vaginal plug is observed *in situ* will be considered day 0 of gestation. Sperm- and plug-positive females will be separated from the male and will remain in their exposure cage. If mating has not occurred after two weeks, the animals will be separated without further opportunity for mating.

Developmental - Maternal Observations

All animals will be observed daily during the study for alterations in behavior or demeanor as previously described under the Reproduction Study "Physical Observation" section, with the exception that a thorough weekly clinical examination will not be conducted. Any animal which dies, appears moribund or shows indications of early termination of pregnancy will be

DRAFT

BFG 01901

BFG42866

submitted for a complete necropsy as described for the Reproduction Study. Body weights will be recorded weekly during premating and gestation as described for the Reproduction Study.

On Day 21 of gestation, all surviving animals assigned to the developmental study will be euthanized by carbon dioxide inhalation and given a limited necropsy. Any obvious structural or pathologic changes noted in the adult will be recorded and the weight of the liver, kidneys and gravid uteri will be recorded. Liver, kidneys and gross lesions will be preserved in neutral, phosphate-buffered 10% formalin, but microscopic examination of tissues will not be conducted unless deemed necessary to interpret other observations made during the study or requested by the sponsors.

Developmental - Fetal Observations

At necropsy, the uterine horns will be exteriorized through an abdominal incision and the following data recorded: 1) the number and position of fetuses *in utero*, 2) the number of live and dead fetuses, 3) the number and position of resorptions, 4) the number of corpora lutea, 5) the sex and body weight of each fetus, and 6) any gross external alteration. The uteri of apparently non-pregnant animals will be stained with a 10% aqueous solution of sodium sulfide (Kopf et al., 1964) and examined for evidence of early resorptions. Corpora lutea will not be recorded for females that are not visibly pregnant at c-section or for females that are submitted for necropsy prior to Day 21. At least one-half of the fetuses in each litter, selected using a table of random numbers, will be examined immediately by dissection under a low power stereo-microscope for evidence of visceral alterations (Staples, 1974). The heads of rat fetuses examined by dissection will be removed, placed in Bouin's fixative and examined by the serial sectioning technique of Wilson (1965). All fetuses will then be preserved in alcohol, eviscerated and stained with alizarin red-S (Dawson, 1926). Skeletal examination will be conducted on all fetuses that were not given visceral examinations.

DRAFT

BFG42867

BFG 01902

Statistical Evaluation.

Descriptive statistics (means and standard deviations) will be reported for feed consumption. Body weights, gestation/lactation body weight gains, organ weights, and sperm count per gram cauda epididymis and percent motile sperm will first be evaluated by Bartlett's test for equality of variances. Based upon the outcome of Bartlett's test, either a parametric or nonparametric analysis of variance (ANOVA) will be performed. If the ANOVA is significant, a Dunnett's test or the Wilcoxon Rank-Sum test with Bonferroni's correction will be performed.

Gestation length, average time to mating, number of corpora lutea, number of implants, litter size, age at vaginal opening and age at preputial separation will be analyzed using a nonparametric ANOVA. If the ANOVA is significant, the Wilcoxon Rank-Sum test with Bonferroni's correction will be performed. Statistical outliers will be identified by the method of Grubbs (1969) and will be routinely excluded from analysis for feed consumption only. Outliers for other endpoints will only be excluded from analysis for documented, scientifically sound reasons. Fertility indices and pregnancy rate will be analyzed by the Fisher exact probability test and Bonferroni's correction will be used for multiple testing of groups in comparison to a single control. Evaluation of the neonatal sex ratio will be performed by the binomial distribution test. Survival indices will be analyzed using the litter as the experimental unit by the Wilcoxon test as modified by Haseman and Hoel (1974). Statistical evaluation of the frequency of pre-implantation loss, resorptions and fetal alterations among litters and the fetal population will be performed using a censored Wilcoxon test with Bonferroni's correction. Nonpregnant females, females pregnant following staining or females having totally resorbed litters will be excluded from the appropriate analyses.

The nominal alpha levels to be used are as follows:

Bartlett's Test (Winer, 1971)	$\alpha=0.01$
Parametric ANOVA (Steel and Torrie, 1960)	$\alpha=0.10$
Nonparametric ANOVA (Hollander and Wolfe, 1973)	$\alpha=0.10$

DRAFT

BFG 01903

BFG42868

Dunnett's Test (Winer, 1971)	$\alpha=0.05$, two-sided
Wilcoxon Rank-Sum Test (Hollander and Wolfe, 1973)	$\alpha=0.05$, two-sided with Bonferroni correction (Miller, 1966)
Fisher's Test (Siegel, 1956)	$\alpha=0.05$, two-sided
Censored Wilcoxon Test (Haseman and Hoel, 1974)	$\alpha=0.05$, two-sided
Outlier Test (Grubbs, 1969)	$\alpha=0.02$, two-sided
Binomial Distribution Test (Steel and Torrie, 1960)	$\alpha=0.05$, two-sided

Because numerous measurements are statistically compared in the same group of animals, the overall false positive rate (Type I errors) will be much greater than the cited alpha levels would suggest. Thus, the final interpretation of numerical data will consider statistical analyses along with other factors such as dose-response relationships and whether the results are significant in the light of other biologic and pathologic findings.

Safety Precautions.

Standard safety precautions will be followed during the conduct of this study.

Quality Assurance.

Permanent records of all data generated during the course of this study, the protocol, any addenda to the protocol, and the final report will be available for inspection by the Quality Assurance Unit. All data generated including the protocol, addenda, and final report will be archived at Health and Environmental Sciences, The Dow Chemical Company, Midland, Michigan.

DRAFT

BFG 01904

BFG42869

REFERENCES

Adams, J., Buelke-Sam, J., Kimmel, C.A., Nelson, C.J. (1985) Collaborative Behavioral Teratology Study: Protocol Design and Testing Procedures. *Neurobehavioral Toxicology and Teratology* 7, 579-586.

Cooper, R. L., Goldman, J. M. and Vandenberg, J. G. (1993). Monitoring of the estrous cycle in the laboratory Rodent by vaginal lavage. In Methods in Toxicology, Volume 3, Part B, Female Reproductive Toxicology. (J. J. Heindel and R. E. Chapin and, Eds.). Academic Press, Inc., New York.

Dawson, A. B. (1926). A Note on the Staining of the Skeletons of Cleared Specimens with Alizarin Red-S. Stain Tech. 1: 123-124.

EEC (1988). European Economic Community. Methods for the Determination of Toxicity. Official Journal of the European Communities, Vol., 31, No. L 133, May 30, 1988. ISSN 0378-6978.

EPA (1985). Environmental Protection Agency Toxic Substances Control Act Test Guidelines, Final Rule. 40 CFR Part 798, September 27, 1985, pp 39426-39433.

EPA (1990). Environmental Protection Agency. Toxic Substances Control Act; Good Laboratory Practice Standards. 40 CFR Part 792 (1 July 1990 Edition).

FDA (1988). Food and Drug Administration Good Laboratory Practice for Nonclinical Studies. 21 CFR Part 58 (April 1, 1988 Edition).

Grubbs, F. E. (1969). Procedures for Detecting Outlying Observations in Samples. *Technometrics* 11, 1-21.

Haseman, J. K. and Hoel, D. G. (1974). Tables of Gehan's Generalized Wilcoxon Test with Fixed Point Sensoring. *J. Statis. Comput. Simul.* Vol. 3, 117-135.

DRAFT

BFG42870

BFG 01905

Hollander, M. and Wolfe, D. A. (1973). Nonparametric Statistical Methods, John Wiley, New York, NY.

Johanson, G. (1990). NEG and NIOSH Basis for an Occupational Health Standard: Propylene Glycol Ethers and Their Acetates. *Arbete och Halsa*, 32.

Kopf, R., Lorenz, D. and Salewski, E. (1964). [The Effect of Thalidomide on the Fertility of Rats: In an Examination of Two Generations.] - Containing the Procedure for Staining Implantation Sites of Fresh Rat Uteri. *Naunyn-Schmiedebergs Arch. Exp. Path. Pharmacol.* 247:121-135.

Korenbrot, C. C., Huhtaniemi, I T. and Weiner, R. I. (1977) Preputial separation as an external sign of pubertal development in the male rat. *Biology of Reproduction* 17, 298-303.

Miller, R. G., Jr. (1966). Simultaneous Statistical Inference. McGraw-Hill, New York, NY.

Miller, R. R., Letts, R. L., Potts, W. J. and McKenna, M. J. (1980). Improved Methodology for Generating Controlled Test Atmospheres. *Am. Ind. Hyg. Assoc. J.* 4, 844-846

OECD (1981). Organisation for Economic Co-Operation and Development-Guidelines for Testing of Chemicals, Section 4-Health Effects, Paris.

OECD (1982). Organisation for Economic Co-Operation and Development-Principles of Good Laboratory Practice, ISBN 92-64-12367-9, Paris.

Siegel, S. (1956). Non-Parametric Statistics for the Behavioral Sciences. McGraw-Hill, New York, NY.

Plowchalk, D. R., Smith, B. J. and Mattison, D. R. (1993). Assessment of toxicity of the ovary using follicle quantitation and morphometrics. In

DRAFT

BFG42871

BFG 01906

Methods in Toxicology, Volume 3, Part B, Female Reproductive Toxicology.
(J. J. Heindel and R. E. Chapin and, Eds.). Academic Press, Inc., New York.

Staples, R. E. (1974). Detection of Visceral Alterations in Mammalian Fetuses.
Teratology 9: 37 (Abstract).

Steel, R. G. D. and Torrie, J. H. (1960). Principles and Procedures of Statistics.
McGraw-Hill, New York, NY.

Wilson, J. G. (1965). Method for Administering Agents and Detecting
Malformations in Experimental Animals. In: Teratology: Principles and
Techniques (J.G. Wilson and J. Warkany, eds.). University of Chicago Press,
Chicago.

Winer, B. J. (1971). Statistical Principles in Experimental Design, 2nd edition.
McGraw-Hill, New York, NY.

DRAFT

BFG42872

BFG 01907

TABLE 1
VINYL CHLORIDE: COMBINED INHALATION TWO-GENERATION REPRODUCTION
AND DEVELOPMENTAL TOXICITY STUDY IN CD RATS

CHRONOLOGY OF EVENTS

WEEKS ON STUDY	F1	F1/P2	F2
1-10	Exposure of P1 males and females prior to first mating.		
11-12	P1 mating period for F1 litters. Mating period for developmental toxicity evaluation		
14-15		F1 born and litters culled on day 4 post-partum to 8 pups each. Necropsy of Developmental toxicity dams and fetal evaluations	
17-18		F1 litters weaned on day 21 post-partum; offspring selected for P2 adults; 1 F1 pup/sex/dose/litter selected for necropsy; remaining pups euthanized.	
19-28	Necropsy P1 adults.	Exposure of P2 males and females prior to first mating.	
29-30		Mating period of P2 for F2 litters.	
32-33			F2 born and litters culled on day 4 post-partum to 8 pups each.
35-36			F2 litters weaned on day 21 post-partum; 1 F2 pup/sex/dose/litter selected for necropsy; remaining pups euthanized.
37		Necropsy P2 adults.	

DRAFT

BFG 01908

BFG42873

TABLE 2

VINYL CHLORIDE: COMBINED INHALATION TWO-GENERATION
REPRODUCTION AND DEVELOPMENTAL TOXICITY STUDY IN CD RATS

STUDY DESIGN

<u>EXPOSURE LEVELS (PPM)</u>	<u>NO. NO. MALES</u>	<u>NO. FEMALES</u>
0	30	30 Reproductive 30 Development
X	30	30 Reproductive 30 Development
Y	30	30 Reproductive 30 Development
Z	30	30 Reproductive 30 Development
TOTAL=		120 240

PARAMETERS

REPRODUCTION STUDY

Animal Observations
Body Weights (parental and neonatal)
Feed Consumption (optional)
Necropsy (parental and weanlings)
Organ Weights (parental and weanlings)
Gross Pathology (parental and weanlings)
Histopathology (parental and weanling)
Fertility Indices
Estrous Cycling and Sperm Analyses
Developmental Landmarks
Neonatal Survival

DEVELOPMENTAL STUDY

Animal Observations
Body Weights (maternal and fetal)
Feed Consumption (optional)
Necropsy
Limited Organ Weights
Gross Pathology
Uterine Examine
External, Visceral and Skeletal Examination

DRAFT

BFG42874

BFG 01909

TABLE 3

VINYL CHLORIDE: COMBINED INHALATION TWO-GENERATION REPRODUCTION AND DEVELOPMENTAL TOXICITY STUDY
IN CD RATS

TISSUES COLLECTED AND PRESERVED AT NECROPSY

ADRENALS	KIDNEYS*	PROSTATE*
AORTA	LACRIMAL/HARDERIAN GLANDS	RECTUM
AUDITORY SEBACEOUS GLANDS	LARYNX	SALIVARY GLANDS
BONE (INCLUDING JOINT)	LIVER*	SEMINAL VESICLES*
BONE MARROW	LUNGS*	SKELETAL MUSCLE
BRAIN (CEREBRUM, BRAINSTEM, CEREBELLUM)	MAMMARY GLAND*	SKIN
CECUM	MEDIASTINAL LYMPH NODE	SPINAL CORD (CERVICAL, THORACIC, LUMBAR)
CERVIX*	MEDIASTINAL TISSUES	SPLEEN
COAGULATING GLANDS*	MESENTERIC LYMPH NODE	STOMACH
COLON	MESENTERIC TISSUES	TESTES*
DUODENUM	NASAL TISSUES*	THYMUS
EPIDIDYIMIDES*	ORAL TISSUES	THYROID GLAND
ESOPHAGUS	OVARIES*	TONGUE
EYES	OVIDUCTS*	TRACHEA*
CROSS LESIONS*	PANCREAS	URINARY BLADDER
HEART	PARATHYROID GLANDS	UTERUS*
ILEUM	PERIPHERAL NERVE	VAGINA*
JEJUNUM	PITUITARY*	

*TISSUE SELECTED FOR HISTOPATHOLOGIC EVALUATION.

DRAFT

BFG42875

BFG 01910