

UNUSUAL SPLENOMEGALIC LIVER DISEASE AS EVIDENCED BY PERITONEOSCOPY AND GUIDED LIVER BIOPSY AMONG POLYVINYL CHLORIDE PRODUCTION WORKERS

H. J. Marsieller and W. K. Leibach

*Medizinische Universitätsklinik
D-53 Bonn, Federal Republic of Germany*

R. Müller and P. Geddigk

*Pathologisches Institut der Universität Bonn
D-53 Bonn, Federal Republic of Germany*

INTRODUCTION

Economic Considerations, Technology

As early as 1833¹ and 1835,² respectively, Regnault in France succeeded in preparing vinyl chloride (VC) and polyvinyl chloride (PVC). However, the road to large-scale production was not opened until vinyl chloride could be synthesized from acetylene and hydrogen chloride (Klatte and Zacharias, 1912)^{3,4} and organic peroxides were used for catalyzing VC polymerization (Klatte and Rollett, 1914).⁵ In 1928, PVC production was started in the United States and in 1933 in Germany.⁶ The enormous growth of VC and PVC production capacities over the past 45 years ran parallel to the rising economic importance of plastic materials⁷ (TABLE 1).

PVC-producing industries in West Germany increased their production, parallel to the development in the United States (TABLE 2), at a rate of 10–20 percent per year from 1960 (172,673 tons of PVC) to 1972 (930,701 tons of PVC and polyvinylidene chloride).^{8–11} Total sales of PVC amounted to 850,140 German marks in 1972.⁹ This enormous increase in production capacity, especially in flexible PVC (U.S., 1965: flexible PVC, 90 percent; rigid PVC, 10 percent)¹² can be explained by the almost unlimited use that can be made of PVC. Diversification of PVC markets exceeds that of any other large-volume thermoplastic and is one of the reasons for the increasingly favorable market situation (TABLE 3).¹³

Rigid PVC is used for tubing and fittings (including insulation material and drainage pipes), foils, films, and sheeting (packaging, lining, recording tapes), profiles (blinds, window frames), tiles, sound records, and fibers. Flexible PVC is used for cables, foils (decoration, roof covering), profiles, tubing, artificial leather, flooring, foam rubber, paint, varnish (lacquer), and toys.¹⁴ Thus, industries that manufacture the end product operate on a much larger scale than those producing the polymer, but the companies are often identical.^{15,16}

Technologically, vinyl chloride was synthesized mainly by hydrochlorination of acetylene, a process which was almost exclusively in operation up to 1960, but in the United States now accounts for only 7 percent of the production.¹⁷ Vinyl chloride can also be produced by oxychlorination of ethylene to 1,2-dichloroethane which is then cracked to VC and HCl by subsequent pyrolysis (now 93 percent of total production in the United States in 1972).^{18,19,20,21} Under

TABLE 1

WORLD PRODUCTION OF PLASTICS IN MILLION TONS PER YEAR¹¹

Year	Million Tons
1950	<0.1
1960	1.5
1965	6.9
1970	30.0

TABLE 2

WORLD PRODUCTION AND PRODUCTION CAPACITY FOR PVC RESINS (MILLION TONS)^{12,13}

	Production			Annual Capacity		
	1950	1960	1972	1960	1972*	1975*
Western Europe	0.053	0.578	2.948	0.583	3.583	
Western Germany	0.014	0.173	0.930	0.195		
United States			1.950	0.610	1.995	3.175
Japan			1.088	0.270	1.587	
Eastern Europe			0.861	Not stated	0.816	
Other areas			0.544	0.152	0.816	
Total			7.391	1.615	8.797	

* Estimated capacities.

TABLE 3
DIVERSIFICATION OF PVC MARKETS¹⁴

PVC Market	Consumption (%)
Calendering	18
Extruded products	14
Wire and cable	12
Calendered flooring	10
Film and sheet	6
Paper and textile coating	5
Sound records	5
Plastisols	5
Protective coatings and adhesives	4
Injection and blow molding	3
Coated flooring	2
Other domestic uses	11
Exports	5

ambient conditions, vinyl chloride ($\text{CH}_2=\text{CHCl}$, mol. wt. = 62.5)¹⁵ is a colorless inflammable gas of faintly sweet odor. At -13.8°C and 760 torr it condenses to a colorless liquid of low viscosity.¹⁶ VC reactivity is almost exclusively due to its double bond; the chlorine atom, as second functional site, does not react easily. The most important property of VC is its ability to polymerize. It also should be mentioned that peroxidation was repeatedly observed to occur when VC came into contact with atmospheric air due to leakage.¹⁷ Commercially produced VC monomer is in general highly purified; degrees of purity of 99.99 per cent are reported. Specific declarations of individual plants state only marginal

amounts of acid (free HCl), acetaldehyde, iron, acetylene, butadiene, water, and stabilizers.¹⁸ Avoiding impurities is a prerequisite since they retard polymerization.¹⁹ Liquid VC is stored and transported in steel containers.^{2,3,4,10} Vinyl chloride is used mainly for polymerization, but it is also used as a refrigerating substance, as a propellant for sprays and aerosols, and for a small number of other chemical reactions.¹⁸

Four methods of polymerization are used for industrial fabrication of polyvinyl chloride: emulsion, suspension, bulk, and solvent polymerization.^{2,3,4,10} Commercially most important were the two older methods of emulsion and suspension polymerization, but now bulk polymerization has been added.²⁰ In 1972, emulsion polymerization accounted for 78 percent, bulk polymerization for 6 percent, and solution polymerization for 3 percent of the total amount of all PVC resins produced in the United States.²¹ Initiation of polymerization is accomplished with compounds that form free radicals at relatively low temperatures, e.g., water-soluble compounds such as peroxide, ammonium, and potassium persulfate for emulsion polymerization or soluble organic compounds, such as lauroyl and benzoyl peroxide, azobisisobutyronitrile and isopropylpercarbonate.^{2,4,10} Formation of free radicals also occurs by thermal decomposition of peroxide bonding ($-\text{O}-\text{O}-$).²² Physical basis for initiation of free radical chain-polymerization is the homolytic dissociation of a pair of π -electrons (covalent bond) resulting in free radicals that contain single unpaired electrons. The single unpaired π -electron of the free radical obtained from the catalyst then reacts with the π -electrons of the double bond contained in the VC monomer, transforming the VC monomer itself into a free radical and thus propagating the growth of a chain of molecules with a terminal free radical. Chain growth is interrupted by saturation of the terminal free radical, often involving a reaction between two growing chains.^{2,4,10,23} The random character of the termination steps accounts for the production of chains of different length and hence different degrees of polymerization with molecular weights statistically distributed around a mean value. One of the operating variables determining specific type of PVC produced is temperature²⁴ which influences degree and velocity of polymerization as well as vapor pressure. Ordinarily, emulsion polymerization proceeds at temperatures of 40–70°C and total reaction pressure of 8–10 atm.

In emulsion polymerization reactors (size: 2,000–5,000 gallons) are charged with the watery phase, VC is added, the agitator is started, and the emulsion is heated. At termination of the polymerization process the reactor content is emptied into a dump tank. The slurry can then be processed by centrifuging and drying. The dry polymer is separated, screened, and bagged. In suspension polymerization water-soluble protecting colloids are added as dispersing agents for stabilization of the suspended vinyl chloride droplets. Subsequent procedures are similar. Unreacted monomer is driven out of the slurry and recovered for recycling.^{2,25} Suspension PVC is almost pure polymer since additives amount to less than 1 percent and most of that leaves when the slurry is centrifuged. For compounding, the powdery or granular polymer is dry-blended under heat and pressure using plasticizers, light and heat stabilizers, and sometimes fillers, lubricants, and pigments or dyes.² PVC dry blends are compounded by hot mixing at fusing temperatures. Depending on the type of end product, fusion temperature lies below or within the softening range (120–160°C). In calendaring (production of endless sheets) temperatures of 100–120°C for rigid PVC and of 150–180°C for flexible PVC are reached. Tiles are fabricated by welding of several layers of calendared sheets in a heated press. In conversion to end products by extrusion

(profiles, pipes, conduit, ribbons) temperatures of 150–300°C can be reached. All this shows that temperature ranges, as a rule, from 100–200°C in converting the thermoplastic PVC resin by various procedures.^{10,11,12}

Toxicology, Occupational Medicine, Clinical Experience (TABLES 4–6)

Simultaneous with the start of large-scale production of PVC in 1930, pharmacologic properties of vinyl chloride^{13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} and problems of occupational health^{101,102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200} were investigated. At first, vinyl chloride appeared to be a suitable narcotic which, even after Oster in 1947¹⁰¹ and Carr in 1949¹⁰² had observed severe cardiac arrhythmia (believed to be due to potentiation of adrenaline effect), was considered to be one of the least dangerous chlorinated hydrocarbons.^{103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200}

With the exception of two communications concerning acute VC intoxication without serious consequence in 2 workers^{101,102} no other data on occupational health hazards in PVC production were reported until 1949 when Tribukh *et al.*¹⁰³ published results of an investigation at a Russian plastics factory. On examination of a group of 73 workers (48 males, 25 females) engaged mostly in processing PVC polymer (compounding at temperatures up to 160°C and ending with release of trapped VC, use of chlorinated naphthalenes and diphenylene as plasticizers, effects of PVC dust), the authors found evidence of what they termed a "more or less marked hepatitis," presenting as a slight enlargement of the tender liver but with conspicuous absence of characteristic complaints, jaundice or bilirubinuria. Other findings were: hypotension, anemia, "chronic gastritis," skin lesions, and a peculiar whitish coating of the upper respiratory tract mucosa. This spectrum of symptoms was attributed to the action of plasticizers, and long-term follow-up studies were thought to be necessary, particularly for prevention of later development of serious liver disease. With the exception of liver disease, similar but badly documented findings, also attributed to additives, were reported by Devigneville,¹⁰⁴ Parmeggiani,¹⁰⁵ and Hervieux and Tessier¹⁰⁶ between 1953 and 1959. Acute intoxication was again reported, this time in Great Britain, with acute collapse during VC polymerization¹⁰⁷ and a second degree burn after direct contact with vinyl chloride.¹⁰⁸ Filatova *et al.*¹⁰⁹ were the first to point out a prevalence of "toxic angioneuropathy" in workers engaged in VC polymerization in spite of working area concentrations below Russian maximal allowable concentration (MAC) values (1 mg of VC per liter = 391 ppm). In 1960, Danziger¹¹⁰ described 3 cases of acute VC intoxication occurring in VC polymerization, 2 of them fatal. Suciu *et al.*¹¹¹ were the first to give a more detailed analysis of the disease spectrum encountered in 168 workers of two Rumanian PVC-producing plants who had been studied during a 4-year period. Their paper ranks as the earliest description of vinyl chloride disease. Gastrointestinal symptoms dominated; in addition, central nervous system disturbances were found in a number of cases, a Raynaud-like syndrome in 10 workers, contact dermatitis and pseudoscleroderma in another 10, and alteration of thyroid function in 3. The authors found hepatomegaly in 51 and splenomegaly in 10 workers. BSP retention proved to be elevated in 2 of 11 cases examined. Liver biopsy in 2 cases revealed chronic hepatitis. A combination of various symptoms was seen in quite a number of workers. In 1965, Filatova *et al.*¹¹² again reported on a Raynaud-like syndrome and central nervous system symptoms in PVC workers. Pushin¹¹³ found liver and biliary tract disease in 15 percent of those workers of the Sverdlovsk chemical plant who were engaged in PVC production. On the basis of a clinically slightly enlarged tender liver with mild hyperbilirubinemia occurring in these cases, he diagnosed amietic, chronic

"epithelial" hepatitis. As possible etiologic factors, primary ingredients for polymerization, compounds released from the polymer during calendaring, and plasticizers (phthalates), were discussed. In 1966, Cordier *et al.*¹¹⁴ were the first to describe acroosteolysis of the distal phalanges combined with a Raynaud-like symptomatology in 2 reactor cleaners (VC polymerization). Subsequently, similar cases were reported in 1967 by Chatelein and Matillon,¹¹⁵ Marin *et al.*,¹¹⁶ Bouteiron,¹¹⁷ Wilson *et al.*,¹¹⁸ and Harris Adams.¹¹⁹ With this clustering of publications the disease became known as "occupational acroosteolysis" with typical radiogenic findings, often combined with Raynaud's syndrome and accompanied rarely by scleroderma-like skin lesions; the disease seemed to occur only in reactor cleaning ("polyelester disease").¹²⁰ Between 1969 and 1972, Anghel *et al.*,¹²¹ Filatova and Antonovchenko,¹²² Dinnan *et al.*,¹²³ on the basis of a large-scale epidemiologic study, Lelievre,¹²⁴ Markowitz *et al.*,¹²⁵ and Dugues *et al.*¹²⁶ observed further cases of this syndrome, occurring only in workers engaged in VC polymerization. It appeared as if processing the polymer would not entail any health hazard (in spite of reflections on the content of trapped substances^{127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200}). Single deviant observations made in 1970 by McCord¹³¹ and by Almgeld *et al.*¹³² in 1973 as well as warnings voiced in the past by Tribukh *et al.*¹⁰³ in 1949 and by Danishevskii *et al.*¹³³ in 1961 obviously were not sufficient to cast doubt on this concept, especially since many standard textbooks had stressed the absolute harmlessness of processing the polymer,^{134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200} apart from the minimal risk attributed to VC inhalation^{139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200} (TABLE 4).

Animal experiments between 1930 and 1963 (TABLE 5) were first limited to the assessment of acute VC toxicity. However, after 1961 four research groups presented results on chronic toxicity of vinyl chloride (TABLE 6) that were not repeated on a large-scale basis. The importance of these results seems to have been underrated, in part even by the authors (e.g., Lester *et al.*¹³⁵) themselves. The experiments of Maltoni *et al.*¹³⁶ that up to now were made known only in part, deal essentially with the carcinogenicity of vinyl chloride. These results had been preceded by the experimental work of Viola and co-workers.^{137,138} However, results published by Torkelson *et al.*¹³⁹ in 1961 induced the German Standards Advisory Committee to reduce MAC standards for vinyl chloride from 500 ppm (in operation since 1966) to 100 ppm in 1970.¹⁴⁰ In 1972, a group of dermatologists at Bonn University were the first in Germany to report on cases of "occupational acroosteolysis," with Raynaud's syndrome and scleroderma-like skin lesions.¹⁴¹ They found the symptomatology of this disease, however, to be much more complex than was so far assumed and hence proposed calling this syndrome "vinyl chloride disease."¹⁴²

LIVER DISEASE IN PVC WORKERS

Clinical Observations; General Outline

It seems quite remarkable that after the findings of Tribukh *et al.*¹⁰³ in 1949 liver damage was only mentioned marginally in subsequently published papers and that no intensive investigation of this feature of vinyl chloride-polyvinyl chloride toxicity had been initiated before 1972. One of the reasons for this late detection of life-threatening liver disease may be the minimal degree of functional hepatic abnormalities generally encountered in these workers which contrasts strikingly with gross morphologic changes, seen at peritoneoscopy and with the development of portal hypertension and angiocarcinoma of the liver which both are now recognized to be associated with exposure to vinyl chloride.

TABLE 4
CLINICAL MANIFESTATIONS AND FATALITIES AMONG WORKERS IN VINYL CHLORIDE POLYMERIZATION (PM),
AND POLYVINYL CHLORIDE PROCESSING (PC) (REVIEW OF LITERATURE)

Author	No. of Workers		Acrosteo- lysis	Raynaud- like Phenomenon	Skin Lesions	Liver Disturbances	Acute Intoxication (Fatalities)
	PM	PC					
Dublin and Vane (1933, 1941)	2						Slight intoxication
Tribukh <i>et al.</i> (1949)	73				—*	21	
Chief Inspector of Factories (1953)	2						Collapse
Harris (1953)	1				1		Second-degree burn
Devigneville and Flucher (1953)	—†				—*	—*(?)	
Parneggiani and Sassi (1955)	13					—*(?)	
Filatova <i>et al.</i> (1957, 1958)	—†			—*			
Hervieux and Tessier (1959)		145			2		
Danziger (1950)	3						1 Slight narcosis, 2 fatalities
Suciu <i>et al.</i> (1963, 1967)	168			10	10	51‡	
Filatova <i>et al.</i> (1965)	163			—*			
Pushin (1965)	350§					15%‡	
Cordier <i>et al.</i> (1966)	123		2	2	2		
		178					
Chatelain and Motillon (1967)	441		5	12			
Martin <i>et al.</i> (1967)	528		5	12	5		
Spain	1		1				
Italy	1		1				
Benitochon	1(¶)		5				
Wilson <i>et al.</i> (1964)	1 (¶¶)		11	22	8		
Harris and Adams (1965)	300	1,000	1	1	1	10	
Asplund <i>et al.</i> (1965)	900		1	8	1		
McCord (1966)		1	1		1		1 Fatality
Filatova and Antonyuzhenko (1971)	1			"Chronic occupational poisoning"			
Duman <i>et al.</i> (1971)	3,751		41	24			
		1,257		1			
Kramer and Mutchler (1972)	98					Slight**	
Lefèvre (1972)	780		19				
Markowitz <i>et al.</i> (1972)	2		1	2	2		
Dugois <i>et al.</i> (1972)	1		1	1	1		
Misgeld <i>et al.</i> (1973)		1	1	1	1		

Total	10,056	2,582	118	97	40	73	5 Acute intoxications 3 Fatalities
	86 ± x						
	12.724						

* --, number of abnormal findings not stated.

† --, number of workers not stated.

‡ Liver biopsy in 2 cases—diagnosis "chronic hepatitis;" splenomegaly in 10 cases.

§ 350 workers employed in the Swerdlowsk plastics factory, number of PVC workers contained therein not stated.

¶ Liver disturbances in 6.57 percent (23/350) of 350 chemical plant workers, in 15 percent of PVC workers.

|| Palpable liver and persistently raised serum bilirubin.

** "Some changes in liver function," with statistically significant correlation of BSP retention and exposure.

TABLE 5
VINYL CHLORIDE EXPOSURE IN LABORATORY ANIMALS—ACUTE TOXICITY (REVIEW OF LITERATURE)

Author	Species*	Concentration (ppm)	Length of Exposure	Effect	Pathology
Patty <i>et al.</i> (1930)	g†	200,000-400,000	"Very short time"	Fatal (lethal range)	Lung: edema Liver: hyperemia Kidney: hyperemia
		100,000 5,000	30-60 min Several h	"Dangerous to life" No severe acute disturbances No cardiac failure	
Schaumann (1934)	Starling heart-lung preparation (cat)†	250,000-300,000	?		
Schaumann (1938)	m + r†	Slight narcosis	4 h/5-8 d; 1 h/d; 4 wk	Narcosis	Liver and kidney: no changes
	d†	100,000	3 h/7 d during 3 wk	Narcosis	Liver and kidney: no considerable changes
Peoples and Leake (1933)	m†	245,000-285,000 85,000-125,000	10 min 10 min	Fatal (lethal range) Minimal anesthetic range	
	d + rb†	170,000	?	Narcosis	"No apparent untoward effects even after prolonged anesthetization"
Oster <i>et al.</i> (1947)	d (6)	100,000 (narcosis)	?	Serious cardiac arrhythmias	
Carr <i>et al.</i> (1949)	d (7)	? (narcosis)	?	Narcosis: sensitization of myocardium	

Mastromattico <i>et al.</i> (1960)	m (5)	100,000	30 min	Deep narcosis—side position with tremors	Fatal outcome: Lung: vascular engorgement with hemorrhages and edema Liver: congestion, partly fatty infiltration (g) Kidney: congestion Clotting defect Survivors: "Little difference from control animals (liver, kidney, spleen, brain, heart, adrenals, lymph nodes, eyes) except for pulmonary congestion"	
	r (5)					
	g (5)					
	m (5)	200,000	30 min	Deep narcosis (1 mouse dead)		
	r (5)					
	g (5)	300,000	30 min	Dead; g in deep narcosis, 1 died hours after exposure		
m (5)						
g (5)	400,000	30 min	Deep narcosis, 1 died during, 1 after exposure			
r (5)						
g (5)						
Lester <i>et al.</i> (1963)	rt	50,000		Moderate intoxication Righting reflex lost Corneal reflex disappeared	"No visible gross pathology"	
		70,000				
		100,000				
	r (2)	150,000		Deep narcosis, 1 rat died of respiratory failure		Lung: congestion/edema

* Abbreviations used: r, rats; g, guinea pigs; rb, rabbits; d, dogs; m, mice. Number of animals stated is given in parentheses.
† Number of animals not stated.

TABLE 6
VINYL CHLORIDE EXPOSURE IN LABORATORY ANIMALS—CHRONIC TOXICITY (REVIEW OF LITERATURE)

Author	Species*	Concentration (ppm)	Length of Exposure	Response	Pathology
Torkelson <i>et al.</i> (1961)	r (20)	500	7 h/d, 5 d/wk for 4.5 mo	Normal growth	Liver: average weight increase, centrilobular granular degeneration Kidney: interstitial and tubular changes
	r (24) g (9) rb (3) d (1)	200	7 h/d, 138–144 times in 204 d	Normal in appearance, mortality, and growth	Liver: microscopically normal in rats, guinea pigs, and dogs; centrilobular granular degeneration in rabbits (together with necrosis and some foamy vacuolation in male and necrosis with periportal cellular infiltration in female); weight increase of rat livers
	r (5) r (5) r (5) r (5)	200	4 h/d 2 h/d for 6 mo 1 and 0.5 h/d for 6 mo	Normal in appearance, mortality, and growth "Entirely normal"	Liver: average weight increase
	(same procedure as in 200 ppm)	100		Normal in appearance, mortality, and growth	Liver: slight increase of average weight in rats after exposure 7 h/d, 138–144 times in 204 d and either 4 or 2 h/d for 6 mo
	r (48) g (24) rb (6) d (2)	50	7 h/d, 130 times in 189 d	Normal in appearance, mortality, and growth	Normal
Lester <i>et al.</i> (1963)	r (18)	100,000 80,000	8 h/d for 2 d 8 h/d for 13 d	Narcosis, 10 rats died, initial loss of weight, followed by normal growth ("tolerance to the gas")	Lung: acute focal necrotizing pneumonia, edema, metaplasia Liver: parasitic cysts, moderate to marked swelling of cells, definite fine to medium or large irregular vacuoles, compression of sinusoids Spleen: lymphocytic hyperplasia including most of the interstitium
	r (10)	50,000	8 h/d for 19 d	Initial loss of weight, followed by normal growth ("tolerance to the gas")	Liver: parasitic cysts, marked swelling of cells, large irregular vacuoles or clear spaces, compression of sinusoids, changes focal to diffuse

	r (30)	20,000	8 h/d, 5 d/wk for 3 mo	Normal in appearance	Spleen, kidney: no differences between control and experimental animals Liver: parasitic cysts, moderate to marked swelling of cells, definite fine to medium or large irregular vacuoles, compression of sinusoids Spleen: no differences between control and experimental animals Conclusions: 1. The only finding suggesting a specific toxic action of VC is the increase in liver weight on exposure to 50,000 ppm for 19 d and to 20,000 ppm over 92 d 2. Morphological alterations without pathologic significance
Viola (1970)	r (25)	30,000	4 h/d, 5 d/wk for 12 mo	Slightly soporific, normal in appearance and growth After 10 mo loss of weight 13 died from cardiorespiratory complications, 2 from hematoperitonium	Liver: increased volume, smooth surface, sometimes subicteric, somewhat more brittle consistency than normal Microscopically: Alteration of hepatocytes with marked cytoplasmic and nuclear polymorphism (torpid degeneration of the cytoplasm to necrosis) Proliferation and hypertrophy of Kupffer-cells Round cell infiltration of portal tracts Blockade of portal capillaries, centrilobular veins and sinusoids by numerous centrally located areas of partial necrosis Intensive fibrosclerotic reaction along areas of degenerative processes Fatty change Kidney: signs of tubulonephrosis sometimes accompanied by chronic interstitial nephritis

TABLE 6—Continued

Author	Species	Concentration (ppm)	Length of Exposure	Response	Pathology
Viola <i>et al.</i> (1971)	r (26)	30,000	4 h/d, 5 d/wk for 12 mo	Slightly soporific, normal in appearance and growth After 10 mo development of paraneuritic tumors	Arteries: signs of endothelial fibrosis (with thickening and dissociation of the elastic membrane), often with partial hyaline degeneration and homogenization of the walls and proliferation of endothelium Skin: superficial hyperkeratosis, thickening of epidermis, dissociation of collagen bundles, diminution and fragmentation of elastic reticulum Bones: extensive periosteal proliferation of cartilage-like material (small metatarsal bones) Brain: degenerative lesions of grey and white matter Nerves: surrounded and invaded by fibrotic processes Liver: enlarged, very fragile, "severe chronic hepatitis" Kidney: swelling of parenchyma, often tubulonephrosis Lung: interstitial pneumonia Brain: degenerative lesions Tumors: skin in 17 cases; lung in 7; bone in 5 Cardiovascular disorders Changes in bioelectric activity of hypothalamus Hyperadrenalinemia Osteoporosis and resorption of bone tissue
Basalacv <i>et al.</i> (1972)	r + rb†	11.7-15.6	?		

* Abbreviations used: r, rats; g, guinea pigs; rb, rabbits; d, dogs; m, mice. Number of animals stated is given in parentheses.
† Number of animals not stated.

In December 1972, our colleagues from the Department of Dermatology alerted us to the possible occurrence of serious liver disease in workers of a nearby chemical plant who were engaged in vinyl chloride polymerization and to a smaller degree in processing the polymer to the final plastic product. We started an extensive examination program that included peritoneoscopy and guided liver biopsy because we felt that this procedure might yield deeper insight into nosology and pathomorphology of vinyl chloride-associated hepatic disease than merely blind percutaneous biopsy, since a larger proportion of cirrhotic or fibrotic changes may escape detection if only blind biopsy is employed.

According to information furnished by the plant physician, a total of roughly 180 workers had in the past been exposed to vinyl chloride in this plant, either as reactor cleaners or in further processing of vinyl chloride (centrifuging, drying, sizing, and bagging the polymer).

To date, we have been able to collect data from a total of 50 workers of this plant (Table 7). Their age distribution is shown in Figure 1. More than half of them were of non-German origin (nationality: German, 23; Greek, 21; Turkish, 6). Among the 50 workers examined 44 had been working in the polymerization process (39 of them alternatively as reactor cleaners) and 6 in processing the polymer into the final plastic products. The length of exposure (Figure 2) ranged from 9 months to 21½ years for the 44 polymerization workers, with 15 of them having had an exposure of 7 or more years' duration. For the 6 men working in the postpolymerization phase the length of exposure was 13, 9, 5½, 4½, 4, and 2 years, respectively. Case 50 had been working in the postpolymerization phase for 2 years and subsequently in the manufacture of high-strength paper (hardening by epoxy compounds) at the same plant for 10 years.

Medical History

In the overwhelming majority of cases previous history was uneventful. One worker (case no. 1) had a previous history of jaundice of unknown etiology 18




FIGURE 1. Age distribution of 50 PVC production workers (age at examination).

SYNOPSIS OF ANAMNOSTIC, CLINICAL, BIOCHEMICAL, PATHOLOGIC, AND HISTOLOGIC DATA OF 50 PVC WORKERS
 Abbreviations used are: *Nationality*: Ger, German; Gr, Greek; Turk, Turk; *Job Classification*: PM, VC polymerization; PC, processing of the polymer; *Pathology*: N, normal; P, patchy; *Histology*: (+), minimal; ++, slight; +++ moderate; ++++, marked; *Surface relief*: U, slightly irregular or undulated; G, granular to finely nodular; S, coarsely nodular; C, comma-like or stellate; R, finely to coarsely reticular; S, S.

No.	Nationality	Age (yr)	Job Classification	Exposure (yr/mo)	Alcohol Intake (g/d)	Central Nervous System	Upper Abdominal Symptoms	Hepatic Enzymes*	Splenomegaly*	Serum Bilirubin (>1.0 mg/100 ml)	BSP Retention (>5% after 45 min)	SCOT (>12 IU/mU/ml)	SCPT (>12 IU/mU/ml)	Masson's-like Phenomenon	Stain Lesions	Peritonoscopy	Histology	Spiders Size by Scintigraphy (cm)	Relationship of Spleen	Esophageal Varices	Thrombocytopenia (<150 X 10 ⁹ /mm ³)
1	Ger	50	PM	21/9	1	+	+	1.5	1.5		8.7	15	15 (21)	+				Normal (liver scan)	++	+	+
2	Ger	56	PM	18/6	10-32	+	+	2	2		6.1	13	19 (24)	+				Markedly enlarged (liver scan)	++	+	+
3	Ger	47	PM	18	10-32	+	+	2	2		15.6	13	19 (24)	+				Markedly enlarged (liver scan)	++	+	+
4	Ger	46	PM	17/6	8	+	+	1.5	1.5	1.1	10.2	13 (22)	15 (27)	+				Markedly enlarged (liver scan)	++	+	+
5	Ger	51	PM	17	48-64	+	+	1.5	1.5	1.1	9.1	19	17	+				Markedly enlarged (liver scan)	++	+	+
6	Ger	51	PM	16	8	+	+	1.5	1.5	1.1	11.3	25	22	+				Markedly enlarged (liver scan)	++	+	+
7	Ger	52	PM	15/3	8	+	+	2	2	1.2	16.7	25	22	+				Markedly enlarged (liver scan)	++	+	+
8	Ger	53	PM	14/6	8	+	+	1.5	1.5	2.0	5.6	15	13	+				Markedly enlarged (liver scan)	++	+	+
9	Ger	44	PM	13/9	8	+	+	1.5	1.5	1.1	17.6	31	13	+				Markedly enlarged (liver scan)	++	+	+
10	Ger	39	PM	13/9	16-32	+	+	1.5	1.5	1.1	6.2	15	13	+				Markedly enlarged (liver scan)	++	+	+
11	Ger	48	PM	13	16-32	+	+	1.5	1.5	1.1	6.2	15	13	+				Markedly enlarged (liver scan)	++	+	+
12	Ger	27	PM	12/6	28	+	+	1.5	1.5	1.1	6.2	15	13	+				Markedly enlarged (liver scan)	++	+	+
13	Gr	40	PM	12/3	28	+	+	1.5	1.5	1.1	6.2	15	13	+				Markedly enlarged (liver scan)	++	+	+
14	Gr	52	PM	0/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
15	Gr	35	PM	9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
16	Gr	45	PM	8/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
17	Gr	42	PM	6/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
18	Gr	54	PM	6/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
19	Turk	31	PM	5/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
20	Turk	33	PM	5/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
21	Turk	33	PM	5/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
22	Gr	36	PM	5	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
23	Gr	36	PM	5	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
24	Gr	30	PM	5	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
25	Gr	35	PM	4/9	16-32	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
26	Gr	30	PM	4/9	16-32	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
27	Gr	31	PM	4/9	16-32	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
28	Gr	31	PM	4/9	16-32	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
29	Turk	33	PM	4/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
30	Turk	30	PM	3/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
31	Turk	36	PM	3/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
32	Turk	33	PM	3/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
33	Gr	32	PM	3/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
34	Gr	40	PM	3/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
35	Gr	31	PM	3/3	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
36	Gr	38	PM	3/3	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
37	Gr	47	PM	3/0	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
38	Gr	30	PM	3/0	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
39	Turk	30	PM	2/9	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
40	Gr	33	PM	2/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
41	Gr	41	PM	1/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
42	Gr	41	PM	1/6	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
43	Gr	33	PM	0/3	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
44	Turk	33	PM	0/3	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
45	Gr	48	PM	13/0	14	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
46	Gr	44	PM	9/0	28	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
47	Gr	39	PM	5/3	16	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
48	Gr	45	PM	4/6	44	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
49	Gr	37	PM	4/0	44	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+
50	Gr	36	PM	2/0	32	+	+	2	10	1.4	13.0	16	16	+				Markedly enlarged (liver scan)	++	+	+

* Centimeters below costal margin.

† For explanation see Table 11.

‡ For explanation see Table 11.

SPI-03086


TABLE 8
ALCOHOL CONSUMPTION IN 50 PVC WORKERS

Reported Alcohol Intake (in g ethanol/dl)	No.
None	10
Up to 8 g	15
Up to 16 g	16
Up to 32 g	6
Up to 64 g	2
Not stated	1
Total	50

TABLE 9
GASTROINTESTINAL BLEEDING (ESOPHAGEAL VARICES) AND SURGICAL INTERVENTION IN 50 PVC WORKERS

	No.
Gastrointestinal bleeding	4/50
Splenectomy	3/50
Splenorenal shunt	1/50
Gastric resection for anaplastic carcinoma	1/50

FIGURE 2. Length of exposure in 44 workers engaged in t polymerization process.



years prior to examination (length of exposure: 21% years disease" was diagnosed in 1968. In 3 cases a symptomatic culus was found on cholecystography (case nos. 43, 45, 50) (case no. 48), who had been engaged in the postpolymerization for 4½ years and who in the past 2 years had repeated for gastric ulcer, the present examination by barium meal revealed a small anaplastic carcinoma of the stomach for which total gastric resection. A German worker (case no. 18) had in 1971, shortly before splenectomy was performed, been of bleeding from an endoscopically diagnosed pyloric ulcer. One Greek worker (case no. 38) suffered from hemophilia A, discovered at appendectomy in 1971; he had taneous hemorrhage and factor VIII levels ranged between 18 and 50 percent. Another Greek worker (case no. 49) was admitted to the hospital because of deafness of sudden onset, combined with because of elevated transaminase levels he was later referred to the Medical Department.

None of the workers had a history of alcoholism. Ter any form (TABLE 8). Alcohol intake was not stated in worker (case no. 36) had formerly consumed 110-160 g entering the plant had reduced his intake to 8-16 g. Several an aversion to alcoholic beverages or an alcohol intolerance after entering the plant.

Complaints

No clear-cut abdominal symptomatology was found. Complaints were usually abdominal discomfort.

Another 24 claimed to have experienced temporary nervous symptoms, such as dizziness, slight orientation, disturbance of vision, and mild headache, often combined with nausea and mild upper abdominal discomfort. Twelve workers spontaneously reported to have smelled "the gas" occasionally which they claimed had a faintly sweet odor. A Raynaud-like symptomatology was reported in 8 cases. Four had experienced episodes of bleeding from esophageal varices during the past 2 years (TABLE 9).

Methods of Investigation

Besides physical examination, a broad spectrum of laboratory tests was applied to each case (TABLE 10). X-ray examination of the upper gastrointestinal tract and intravenous cholecystography were carried out in all cases as were a plain chest film and an ECG with 12 leads. In 3 cases esophagogastrosocopy was done. Scintigraphy of liver and spleen was performed in 41 workers.

Peritoneoscopy and guided liver biopsy were performed by us in 41 workers, peritoneoscopy without biopsy in one, and percutaneous blind needle biopsy in two other cases (total: 44 cases). Data of case 14 were put at our disposal by courtesy of Dr. Harald Henning* who, during a 3-year follow-up including three peritoneoscopies, had previously classified the case as noncirrhotic portal fibrosis with portal hypertension of unknown etiology. We were given access to data of cases 10, 24, 29, and 31 which had been examined by peritoneoscopy and biopsy at the Medizinische Universitäts-Poliklinik†; 1 of these 4 cases (case no. 10) later

* Head, Klinik Föhrenkamp of the Federal Board for Employee's Insurance, D-241 Mülin, Germany.

† We would like to thank Prof. Dr. F. Krück, Direktor der Medizinischen Universitäts-Poliklinik, Bonn, and Prof. Dr. A. Guigemann, Direktor der Chirurgischen Universitätsklinik, Bonn, for permission to include data of 3 and 2 cases, respectively.

TABLE 10
SPECTRUM OF LABORATORY TESTS PERFORMED IN 50 PVC WORKERS

1. Blood sedimentation rate; hemoglobin; red cell count; white cell count; differential cell count; hematocrit reading; number of platelets	
2. Total bilirubin	
3. Glutamic oxaloacetic transaminase	
4. Glutamic pyruvic transaminase	
5. Alkaline phosphatase	
6. Lactic dehydrogenase	
7. Cholinesterase	
8. Thymol turbidity	
9. Total protein	
10. Serum electrophoresis	
11. Bromsulphalein retention	
12. Urinalysis	
13. Blood urea nitrogen	
14. Serum creatinine	
15. Serum uric acid	only in some cases
16. Serum iron	
17. Serum copper	
18. Blood glucose	
19. Total cholesterol	
20. Serum triglycerides	
21. β -Lipoproteins	
22. Blood coagulation factors (II, V, VII, X, antithrombin, thrombin time, thrombelastogram)	
23. Hepatitis-associated antigen and antibody	

underwent splenectomy after repeated hemorrhage from esophageal varices. In a last case (case no. 7) splenectomy and lienorenal anastomosis were done at the Chirurgische Universitätsklinik.† Histology of the spleen was available in 3 cases after splenectomy; in another 8 cases we performed guided biopsy of the spleen for light microscopy; in 5 of these cases material for electron microscopy was also obtained. In addition, liver biopsy material was obtained for electron microscopy from 8 patients.

Results

Physical Examination

*On palpation, the liver was found to be slightly to markedly enlarged in 31 cases. Palpable splenomegaly was noted in 16 cases, in 4 of which it was not accompanied by hepatomegaly. Jaundice, spider angiomas, palmar or plantar erythema or physical demonstrable ascites were not found.

Laboratory Tests

*The most consistently positive biochemical test was 45-min BSP retention (Table 11). A marginally to markedly pathologic retention was seen in 38 patients with a mean value of 10.5 percent. In 12 cases serum bilirubin levels were marginally elevated (maximum: 2.2 mg/100 ml). Likewise, activities of serum alkaline phosphatase were increased slightly in only one-fifth of the total. Activities of serum aminotransferases GOT and GPT were unequivocally pathologic

TABLE 11

RESULTS OF LIVER TESTS IN 50 PVC WORKERS (CLASSIFICATION BASED ON MAXIMAL VALUES OBTAINED IN MULTIPLE DETERMINATIONS)

	45-Min BSP Retention (%) ^a	Serum Bilirubin (1.0 mg/100 ml) ^a	Alkaline Phosphatase (15 mU/ml) ^a	No. of Cases
Normal	12	38	40	
Pathologic (Range)	38	12	10	
(Mean)	(5.1-25.6%) (10.47%)	(1.1-2.2 mg/100 ml) (1.4 mg/100 ml)	(50-110 mU/ml) (71.9 mU/ml)	
	GOIT	No. of Cases	GPT	No. of Cases
Definitely normal	-12 mU/ml -19 mU/ml	4	-12 mU/ml -22 mU/ml	11
Marginally normal	12-20 mU/ml 20-30 mU/ml	31	12-20 mU/ml 23-30 mU/ml	22
Pathologic	20-50 mU/ml 31-60 mU/ml >50 mU/ml >60 mU/ml	13 15 2 50	20-50 mU/ml 31-60 mU/ml >50 mU/ml >60 mU/ml	12 17 5 50
Total				

^a Upper limit.

† Modification of the method described by O. A. H. Bessey *et al.* J. Biol. Chem. 164: 221, 1946; upper normal limit (adults): S. J. Walter and R. Glockner. Ärztl. Lab. 10: 220, 1964.

‡ Owing to a change in methods of determination of enzyme activities during the investigation (later use of a substrate-optimized method) two sets of reference values had to be used.

§ E. and F. W. Schmidt. Enzymol. Biol. Clin. 3: 1, 1963.

¶ Substrate-optimized standard method, W. Thefeld *et al.* Deut. Med. Wochschr. 99: 343, 1974.

in 15 and 17 cases, respectively; however, marginal values may occasionally be the only and earliest biochemical indication of toxic liver damage.¹⁻³

A more or less marked thrombocytopenia (less than 150,000 platelets/mm³) was observed in 42 cases. Except for 1 patient with splenomegaly (case no. 14) who had leukopenia besides thrombocytopenia, hematologic results were otherwise within normal limits. Tests for hepatitis-associated antigen and antibody were negative in all cases. All other biochemical tests including initially introduced but later abandoned immunologic reactions failed to contribute to the problem in question.

Röntgenology; Endoscopy of Upper Gastrointestinal Tract; Scintigraphy

A diagnosis of esophageal varices, in some cases combined with varices located in the cardiac region of the stomach, could be made in 10 workers; in 5 of them this was done by barium meal, in 2 by x-ray plus endoscopic examination, in 1 by barium meal and splenography, in 1 by endoscopy and splenogram, and in the last patient by indirect splenopneumatography. Only in one case (case no. 7) was intraoperative measurement of portal pressure possible, and

TABLE 12
SPLENOMEGALY IN 50 PVC WORKERS
Spleen size as measured by scintigraphy in 48 cases

Methods:
Splenic scanning by ^{197}Tl -BxIHP (40 cases) (normal diameters: $11 \times 7 \times 4$ cm)
Liver scan by $^{99\text{m}}\text{Tc}$ sulfur colloid (46 cases)
A. Within normal limits (4 of the cases = PVC-processing workers): 11/48*
B. Exceeding normal limits (2 of the cases = PVC-processing workers): 37/48†
C. Not examined (previous splenectomy): 2/50

Splenectomy in 3 cases

- Diameters, weight:
1. $15 \times 12 \times 7$ cm (300 g)
 2. Ruptured at splenopneumography (500 g)
 3. $23 \times 13 \times 6$ cm (1250 g) (scintiscan = $24 \times 12 \times 7$)
- * In 6 of the 11 cases by liver scan.
† In 2 of the 37 cases by liver scan.

this proved to be only slightly elevated. Otherwise, no determinations of intra-splenic pressure or of wedged hepatic venous pressure have been carried out so far.

In 5 cases a diagnosis of acroosteolysis was established by x-ray examination of hands and feet.

Approximate spleen size was determined by means of selective scintigraphy using labeled mercury (^{203}Hg -labeled bromomercure hydroxy propane); in 40 cases. Measurement of three diameters (length, breadth, depth) allowed detection of even minor degrees of splenomegaly.^{22,23,24} The organ proved to be slightly to markedly enlarged in 37 out of 48 cases (total assessment of spleen size: 39 out of 50), a proportion that was considerably larger than that obtained by either physical or peritoneoscopic examination (TABLE 12).

Peritoneoscopy and Liver Biopsy

In internal medicine, the paramount advantage of peritoneoscopy lies in the fact that it permits direct inspection of the abdominal cavity and of size, coloring, surface structures, and circumscribed lesions, especially of the two main hepatic lobes and the spleen which, otherwise, can only be accomplished by exploratory laparotomy. Besides inspection of abdominal organs, the method also allows, although only to a limited extent, minor intraabdominal manipulations such as guided biopsy of liver and spleen, determination of organ consistency by probe, and removal of strand-like adhesions. If contraindications are strictly adhered to, serious complications are rare.^{25,26} Based upon a compilation of 63,845 cases in which peritoneoscopy was performed with 48,766 guided liver biopsies, Brühl²⁶ calculated a mortality rate of 0.029 percent. The rate of complications was 2.49 percent, two-thirds of which, however, had been only of minor importance.

In the cases presented here, peritoneoscopic findings were categorized according to the following criteria:

1. Symptoms of hepatic enlargement. These were measured by the relation of

* We wish to thank Prof. Dr. C. Winkler, Direktor des Instituts für Klinische und Experimentelle Nuklearmedizin an der Universität Bonn, for permission to use results of scintigraphic examinations carried out at his institute.




FIGURE 3. Granular appearance of liver surface (conspicuous scattering of light reflection), pronounced coarsely reticular fibrosis of Glisson's capsule, patchy cicatricial fibrosis, distinctly increased capsular vessels. At left upper corner site of guided liver biopsy can be seen. (Case no. 41; length of exposure, $2\frac{1}{4}$ years.)

lower edge of right hepatic lobe to costal margin and a rounding-off of the normally sharp edge.

2. Changes in hepatic surface relief. Hepatic surface relief is normally smooth and glossy with well-defined lobular configuration, mirror-like capsule, and sharply defined light reflection; slightly uneven (irregular) or undulated appearance due to shallow concavities in connection with slight initial cicatrization; granular (FIGURE 3) appearance as initial stage of distortion of hepatic lobular architecture, characterized and recognizable by scattering of light




FIGURE 4. a. Finely nodular surface of the liver. Close-up view of lower edge of left lobe. (Case no. 4; length of exposure, 17½ years.) b. Finely nodular appearance of underside of right hepatic lobe with stellate capsular fibrosis and small scars in a worker engaged in processing the polymer. (Case no. 48; duration of job assignment, 4½ years.)



FIGURE 5. Coarsely nodular hepatic surface resembling advanced cirrhosis; left lobe. Histologically there was, however, only septal fibrosis and collagenization of sinusoidal walls. No evidence of portal hypertension. (Case no. 42; length of exposure, 1½ years.)

reflection, proceeding through finely nodular (FIGURE 4) to eventually coarsely nodular surface relief (FIGURE 5).

3. Presence of different degrees of local or diffuse hepatic capsular fibrosis. The capsule is a very subtle indicator of pathologic processes taking place in the underlying tissue.¹⁰ Capsular fibrosis (FIGURE 6) was usually irregularly distributed but mostly less pronounced toward the convexity; in its mildest form it was characterized by a strand-like or membranous opacity. In some cases we noticed a peculiar comma-like or stellate fibrosis (FIGURE 7), probably corresponding to a delicate scarring process in the underlying hepatic tissue. A more pronounced lesion seemed to be the development of a thinly or coarsely outlined more or less elevated network of thickened capsular connective tissue covering the surface (FIGURE 8). In other cases there were several or numerous small scar-like lenticular patches (FIGURE 9) or broader slightly concave cicatricial lesions probably indicating postnecrotic scarring. In 4 cases focal whitish capsular thickening resembled what is known as focal "perihepatitis cartilaginea" (FIGURE 10).

4. Appearance of capsular vessels. Capsular vessels are normally not visible; pathologically, they may be slightly angulated or conspicuously increased (FIGURE 11) as an indication of underlying inflammatory processes of various activity.

5. Symptoms of splenic enlargement and perisplenitis. Visualization of the spleen by peritoneoscopy is not always possible, even after extreme tilting of the table because the organ may be covered by adipose greater omentum; in this case an effort to bare the lower pole by inserting a palpation probe is some-




FIGURE 8. Coarsely reticular fibrosis of Glisson's capsule; right lobe. (Case no. 9; length of exposure, 13 1/2 years.)



FIGURE 9. Lentil-shaped and patchy irregular milky-white capsular thickening near lower edge of right hepatic lobe. (Case no. 16; length of exposure, 8 1/2 years.)




FIGURE 6. Types of capsular fibrosis of the liver observed in 12 workers.




FIGURE 7. Typical comma-like capsular fibrosis on convexity near falciform ligament (top center); in contrast to finely nodular (see FIGURE 4a) of left lobe, surface on convexity is comparative (4; length of exposure 17 1/2 years.)




FIGURE 10. Coarsely reticulated fibrosis and patchy focal whitish capsular thickening (focal "perihepatitis cartilaginea"), resembling capsular alterations seen in thorotrastosis of the liver.⁶ (Case no. 6; length of exposure, 16 years.)




FIGURE 11. Particularly pronounced increase of capsular vessels in case no. 14 (length of exposure, 11 years) with finely to coarsely nodular surface. The case was formerly diagnosed as "noncirrhotic portal fibrosis and portal hypertension of unknown etiology." Close-up view shot with a special magnifying peritoneoscope (Courtesy of Dr. Hermann Schöberl).



FIGURE 12. Crenate margin of enlarged spleen; lent splenic fibrosis and millet-sized subcapsular hemorrhage in capsule. (Case no. 5; length of exposure, 17 years.)



FIGURE 13. Shaped thick milk-white peritoneal fibrosis. (Case no. 5; length of exposure, 17 years.)

times successful. For assessing the spleen size, the of the spleen to the left costal margin and to the as the inspection of the crenate margin are useful. The enlargement cannot be recognized by peritoneally the crenate margin is deeply indented. Th equally sensitive to processes going on in the un capsule of the liver. In a number of cases minor present in the form of pinhead-sized to somewhat delineated white capsular plaques (Figure 12); a of thick white coating resembling focal "perisple worthy that patchy perisplenitis was found only less marked splenomegaly.

6. *Symptoms of portal hypertension.* Quite early i lateral circulation due to portal hypertension a t tortuosity of preformed peritoneal venous ves falciform ligament, on the anterior stomach wa testine, and especially accompanying intra-ab amounts of ascites can be observed to accumu lobe and the abdominal wall if the table is tilted distinct etiology.

All the lesions described above are non-specific of

• Changes in hepatic surface relief, degree of cap

ilation of the inferior pole renocolic ligament as well Minor degrees of splenic copy. In marked spleno- plenic capsule seems to be lying tissue as is Glisson's grees of perisplenitis were larger lentil-sized sharply rarely were there patches is cartilaginea." It is noie- connection with more or

the development of a col- rked dilatation and later a s can be detected in the on the small and large in- ninal adhesions. Minimal between the right hepatic accordingly.

ations not suggestive of a ar fibrosis, and appearance

TABLE 13
HEPATIC SURFACE ALTERATIONS OBSERVED ON PERITONEOSCOPY (n = 47) OR
AT LAPAROTOMY (n = 2)

Alterations	No.
Surface relief	
A. Smooth	24/49
B. Slightly irregular or undulated	7/49
C. Granular to finely nodular	13/49
D. Coarsely nodular	5/49
Capsular fibrosis (irregularly distributed)	25/49
A. None	6/46
Predominantly:	
B. Conium-like or stellate	3/46
C. Finely to coarsely reticular	20/46
D. Small scar-like patches to broader concave post-necrotic scars	14/46
E. Patchy "perihepatitis cartilaginea"	40/46
Capsular vessels	3/46
A. None (normal)	23/46
B. Slightly augmented	18/46
C. Conspicuously increased	5/46
	23/46

of capsular vessels on the liver surface were recorded and graded on a severity scale. Table 13 shows that in about half of the cases the surface relief of the liver was altered ranging in degree from slightly uneven to coarsely nodular, the macroscopic appearance of the liver being suggestive of advanced cirrhosis in 2 cases (see Figure 5). Similarly, capsular vessels were considered to be slightly to conspicuously increased in half of the total cases. The most consistently positive feature of gross pathology was capsular fibrosis. It was unevenly distributed, sometimes more pronounced on the left lobe than on the right one, and in the majority of positive cases it was present in the form of a more or less conspicuous reticulated capsular thickening.

In cases with massive patchy perihepatitis the picture was reminiscent of hepatic "thrombotosis" (see Figure 10). The proportion of peritoneoscopically determined enlargement of liver and spleen can be seen from Table 14. Pronounced symptoms of portal hypertension were observed in 7 cases. A small amount of ascites was found in only 1 case.

In contrast to gross morphology, however, histologic lesions observed in liver biopsy material were far less pronounced than might have been expected from peritoneoscopic appearance, and this also applies to the cases which laparoscopically resembled advanced cirrhosis of the liver.

In essence, four histopathologic features could be observed (Table 15):

- Degenerative alterations of hepatocytes, occasionally accompanied by phenomena of cytoplasmic adaptation.
- Slight fibrosis and collagenization of sinusoidal walls.
- Polymorphism and polyploidy of liver cell nuclei.
- Activation and proliferation of littoral cells lining sinusoids.

Degenerative alterations of liver cells, manifest within relatively sharply defined intralobular areas (Figure 13), were predominantly characterized by hydropic swelling and granularity of cytoplasm, with occasional single cell necrosis, and were present, to some degree, in all the cases examined. In some cases adaptive changes manifested themselves as "ground glass appearance" of cyto-

TABLE 14
HEPATOSPLENOMEGALY AND SYMPTOMS OF PORTAL HYPERTENSION
OR AT LAPAROTOMY

Hepatosplenomegaly (n = 47)	Splenomegaly (n = 50)	Symptoms (n = 47)	PERITONEOSCOPY
None	31/49	None	40/49
Present	18/49	Present Spleen not visualized*	inable 2/49 7/49

* In 13 cases the spleen could not be visualized during peritoneoscopy even after extreme tilting owing to its being covered by nonremovable omentum.

TABLE 15
SYNOPSIS OF PERTINENT HISTOLOGIC FEATURES IN 49

	Collagenization of Sinusoidal Walls	Enlargement and/or Proliferation of Littoral Cells	Septal Fibrosis	TISSUES
None	10	1	34	26
Present in	39	48	15	23
Grading				
Minimal	7	25	6	6
Slight	32	20	8	14
Moderate	—	2	1	3
Marked	—	1	—	—




FIGURE 13. Degenerative changes of hepatocytes with hydropic swelling and granularity of cytoplasm. Occasional single cell necrosis. Note proliferation of littoral cells (Case no. 14; length of exposure, 11 years after last exposure). H&E (x235).



Figure 14. Micronodular appearance due to septal fibrosis with characteristic hydropic swelling of parenchymal cells (Case no. 5; length of exposure, 17 years; biopsy 2 weeks after last exposure). Trichrome Goldner stain ($\times 94$).

plasma. *Fibrosis*, observed in somewhat less than one-third of the cases, was partly septal, partly portal, and partly midzonal; in 80 percent of the cases there was collagenization of sinusoidal walls, occasionally forming a network-like intralobular fibrosis (Figure 14). *Polymerization of liver cell nuclei* was characterized by enlargement and polyploidy of nuclei. The total number of bi- or multinucleated cells, however, did not seem to be increased. A rather characteristic feature was an *activation and proliferation of sinusoidal cells*. There was a definite increase in the number of littoral cells; they were occasionally arranged in a chain-like fashion and often resembled short pegs. In 6 patients, a bizarre nuclear shape was observed with enlargement and hyperchromatism (polyploidy?), interpreted as a symptom of pronounced cytologic deviation (Figure 15). Less than half of the cases had predominantly minimal to slight fatty infiltration of hepatocytes, not correlated to either length of exposure or drinking habits. More detailed information on special features of histology may be found in Dr. Gedlig's paper.

Discussion

Our synopsis of prior studies published between 1933 and 1973 (Table 4) includes a total number of about 12,500 workers employed in VC polymerization and PVC processing. However, considering the relatively high standards of occupational hygiene in modern chemical industry, *even in the true number of workers examined routinely by plant physicians can be assumed to have been much higher*. Seen against this background and the fact that large-scale production of PVC has now been going on for 45 years, the number of reported cases with (mostly ill-defined) liver disease seems to be surprisingly small. Other mani-




Figure 15. Toxic and adaptive changes of parenchyma. Conspicuous picnotic nuclei of littoral cells (Case no. 3; length of exposure, 18 years; biopsy 2 weeks after last exposure). H&E ($\times 400$).

festations associated with VC polymerization had been seen more frequently (leucocytosis in 118, Raynaud's syndrome in 97, skin lesions in 49 of more than 12,000 workers). They evoked particular interest mainly between 1966 and 1971 and were thought to be restricted to avoidable manual cleaning of reactors.^{3,6,11}

Thus, when we first observed evidence of liver disease of possibly occupational etiology in the workers examined, we suspected plant standards to have been particularly poor, considering that liver disease was so rarely mentioned in prior publications. Intimations made by workers seemed to confirm this assumption (distinct odor of VC vapors at diverse working areas even outside the location of reactors, deteriorating quality of ventilation after windows had been pulled up, frequent symptoms of acute overexposure, and mainly manual reactor cleaning). However, with a growing number of workers examined it became evident that we dealt with a disease presenting with a multifaceted but pre-

nantly inconspicuous symptomatology (for instance, minor hepatic functional insufficiency^{10,11}) which, with reference to liver and spleen, even in advanced stages would be liable to escape detection if only conventional methods of examinations were employed. Therefore, statements given out by other plants that liver and spleen changes had never been observed should be met with scepticism unless proven by morphologic methods.

The dramatic discovery of cases of hemangioendotheliosarcoma of the liver in PVC workers throughout the world early in 1974 has shown that this is certainly not merely the problem of a specific plant.

None of the clinical or morphologic findings seemed to be mutually correlated nor did we find a recognizable correlation between a certain set of symptoms and length of exposure or alcohol consumption. The clustering of pathologic findings (liver, spleen, portal circulation) in this group now comprising 50 workers confirmed our previous impression¹² that we dealt with a symptomatology compatible with the diagnosis of chronic toxic liver damage,^{13,14,15} similar to that seen after repeated or continuous exposure to halogenated hydrocarbons. The histomorphologic appearance of the liver was essentially characterized by *degenerative lesions* (frequently observed in chronic toxic liver injury: hyperplastic smooth endoplasmic reticulum, alterations of rough endoplasmic reticulum due to toxic derangement of membrane structures^{16,17,18}), *fibrosis* (septal fibrosis or collagenization of sinusoidal walls with occasional capillarization and finally network-like interstitial fibrosis), and enlargement or *proliferation of littoral cells*.

Whereas the mostly unimpressive histologic findings permitted reflections as to possible etiology^{19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} to possible etiology^{19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} more suggestive of serious liver damage, could only be interpreted as unspecific evidence of injury; for instance, even pronounced capsular fibrosis merely indicates a repair stage of underlying tissue damage due to a variety of noxious agents.^{11,12} Although granular to nodular surface relief in a number of cases suggested incipient derangement of lobular architecture, a final diagnosis of cirrhosis of the liver could be made only in two cases.¹² Thus, the pathogenesis of portal hypertension with splenomegaly or of isolated splenomegaly at first remained unexplained, especially since no recognizable correlation seemed to exist between degree of hepatic lesions and degree of splenomegaly or of portosystemic collateral circulation. Liver disease in PVC workers reminds one of non-cirrhotic portal fibrosis, predominantly seen in India,^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} or of so-called idiopathic portal hypertension.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} In this disease entity, portal hypertension and splenomegaly are usually accompanied by similar alterations of hepatic surface and histology.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}

However, portal hypertension seemed to have been the result of sclerosis or thrombosis of the portal vein.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} Another analogy between liver disease associated with PVC production and non-cirrhotic portal fibrosis/hypertension can be seen in the frequently observed combination of a pronounced pathologic BSP retention with other laboratory parameters being normal or only slightly aberrant.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} A certain resemblance to Banti's syndrome is suggested by the histologic lesions.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} It should also be realized that the degree of fibrosis in our cases was most probably underrated on microscopy since surgical biopsy material, more suited for detection of fibrosis,^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} was not available. On the other hand, one should keep in mind that even the rather unimpressive alterations observed histologically (periportal and sinusoidal fibrosis, capillarization of sinusoids, degen-

erative cellular lesions) may be sufficient to explain an increase in portal pressure.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}

In this connection, particular attention should be paid to the enlargement and proliferation of littoral cells, as described above, because of their relation to fibrogenesis.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} Besides, proliferation of littoral cells is of diagnostic significance, indicating that the reticuloendothelial system is involved and actively participates in the process (a possibly additional pathogenetic factor in splenomegaly¹¹). In this context, recent reports on oncogenic properties of vinyl chloride^{21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} should be evaluated since proliferative alterations of littoral cells are possibly the precursor stage of hemangioendotheliosarcoma. If one looks for a pattern of injury similar to that seen in VC-associated hepatic hemangioendotheliosarcoma, gross morphology, histologic appearance^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} and conjectural course of events in the pathogenesis of hemangioendotheliosarcoma occurring after exposure to Thorotrast or arsenic can perhaps serve as an example. It could be speculated that chronic stimulation of the reticuloendothelial system (possibly also of hepatocytes¹¹) by a toxic substance¹¹ may be the common denominator.

This leads back to the question of etiology.¹¹ Quantitatively vinyl chloride monomer is the most likely suspect in VC-associated liver disease. Numerous hints (such as odor and acute symptoms of overexposure) contained in job histories point to the frequent release of considerable concentrations of VC at various working areas in VC polymerization.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} even if one considers the unreliability of olfactory sensations^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} and the wide range of olfactory threshold concentrations reported in the literature.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}

Except for Schumann's communication in 1934,¹¹ quoted by Williams¹¹ in 1959, very little is known about the kinetics and metabolism of vinyl chloride.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} Nevertheless, considering the chemical structure of VC and the course of events in polymerization, it can be speculated that free radical mechanisms (monomer, catalysts) may be responsible for damage to cell membranes and subcellular structures (endoplasmic reticulum, "drug-hydroxylating enzyme systems"¹¹), for example, by way of "lipid peroxidation."^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} Furthermore, formation of free radicals may also be involved in the oncogenic action of vinyl chloride.¹¹

Regardless of such deliberations, results of animal experiments with long-term exposure to pure VC have demonstrated considerable hepatosplenic injury^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} apart from tumorigenesis and damage to other organs.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} From these results it appears quite probable that the pattern of damage observed can be attributed to the effect of VC itself. It is, however, not established¹¹ whether and if so how far various additives¹¹ or intermediate compounds¹¹ contribute to the symptom pattern in human vinyl chloride disease.

Some comment should be made on the question of a potential health hazard for those engaged in processing the polymer to the final product. This question is even more difficult to answer and in practice of even greater importance. Detection of analogous if less pronounced lesions in some workers engaged in the manufacture of floor tiles should be reason enough also to pursue the question of possible VC-induced damage in these later stages of production. This is also suggested by the fact that VC trapped in the polymer can principally be liberated^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} during processing (e.g., calendaring, moulding, extrusion, etc.) at temperatures of 100–200°C.^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} At the moment, it is also open to discussion whether PVC dust, apart from damage to the respiratory tract,^{11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100} may cause gastrointestinal symptoms,¹¹ for instance, by persorption.¹¹

In the future, periodic control examinations of all PVC workers, especially of those engaged in VC polymerization, at short intervals employing sensitive methods should be obligatory unless it should prove feasible to eliminate completely any possible air contamination.

Some of the many scientific questions connected with the problem of vinyl chloride disease may not be answerable until results of further extensive animal experiments^{20,21} and of prospective and retrospective large-scale studies²² as well as of continuous monitoring of working areas^{23,24,25,26,27} are available.

REFERENCES

- ADAM, Y. G., A. G. HUVOIS & S. I. HADJU. 1972. Malignant vascular tumors of liver. *Ann. Surg.* 175: 375.
- ALMAGUIT, L. F. 1967. Vinyl chloride processes. *Chem. Eng.* 74 (March 27): 123.
- ALMAGUIT, L. F. 1967. Manufacture of vinyl chloride. *Chem. Eng.* 74 (April 10): 219.
- ALMAGUIT, L. F. 1967. Polymerization of vinyl chloride. *Chem. Eng.* 74 (May 8): 151.
- ALMAGUIT, L. F. 1967. Vinyl chloride polymerization by suspension processes yields polyvinyl chloride resins. *Chem. Eng.* 74 (June 5): 145.
- ALMAGUIT, L. F. 1967. Vinyl chloride polymerization by emulsion, bulk and solution processes. *Chem. Eng.* 74 (July 3): 85.
- ANGHELUSCU, FL., M. OTTOU, E. DOMINISCU, L. HAGA-PARASCHIV-DOSSIOS, G. DOMINISCU & V. GANEA. 1969. Consideratii clinico-patogenice asupra fenomenului Raynaud la muncitorii din industria policlorurii de vinil. *Med. Internu* 21: 473.
- AUSTIN, G. T. 1974. Industrially significant organic chemicals (Part 3). *Chem. Eng.* 81 (March 19): 87.
- BAADER, F. W. 1960. Klinische Grundlagen der sekundarvierzier mieldpflichtigen Berufskrankheiten. Urban & Schwarzenberg, München, West Germany.
- BADIER, F. W., Ed. 1961. Handbuch der gesamten Arbeitsmedizin. Vol. II/1: Berufskrankheiten. Urban & Schwarzenberg, München, West Germany.
- BAGHIZADEH, M. F., E. U. HIRSH & P. SCHUMACHER. 1971. Hamangioendothelium (Reticulosarcoma angiosarcoma) der Leber und alte Echinokokkusinfektion. *Acta Hepato-Splenol.* 18: 224.
- BARATTA, E. D., R. D. STEWART & J. F. MURCHER. 1969. Monitoring exposures to vinyl chloride vapor: breath analysis and continuous air sampling. *Am. Ind. Hyg. Assoc.* 30: 537.
- BASALAY, A. V., A. N. VAZIS & A. G. KODIRKOV. 1972. K patogenezu izmenenii vozmikayushchikh pri dlinnom vozdeistvii vinilklorida. *Gigiena Truda Prof. Zabolevaniya* 16(2): 24.
- BASU, A. K., J. BOYER, R. BHATTACHARYA, K. C. BASU MALLIK & K. P. SIN GUPTA. 1967. Non-cirrhotic portal fibrosis with portal hypertension: a new syndrome. Part I. Clinical and function studies and results of operations. *Indian J. Med. Res.* 55: 336.
- BASU MALLIK, K. C., K. P. SIN GUPTA, A. K. BASU, S. K. BISWAS, N. C. PAL & J. BOYER. 1967. Non-cirrhotic portal fibrosis with portal hypertension: a new syndrome. Part II. Histopathological studies. *Indian J. Med. Res.* 55: 351.
- BECK, K., W. DISCHER, M. HELMS, B. KIANG, K. SICKINGER & R. TINSER. 1968. Atlas der Laparoskopie. F. K. Schattauer Verlag, Stuttgart, West Germany.
- BERGER, V. & K. BOSSCHER. 1961. Ueber das Hamangioendothelium der Leber. *Acta Hepato-Splenol.* 8: 356.
- BISCHOFF, L. 1970. Zum Problem der Leberfibrose. *Schweiz. Med. Wochenschr.* 100: 1214.
- BOYER, J. L., K. P. SIN GUPTA, S. K. BISWAS, N. C. PAL, K. C. BASU MALLIK, F. L. INH & A. K. BASU. 1967. Idiopathic portal hypertension. Comparison with the portal hypertension of cirrhosis and extrahepatic portal vein obstruction. *Ann. Internal Med.* 66(1): 41.
- BREIT, M. D. J. 1974. Vinyl chloride and cancer. I: 599.
- BROSI-SABIR, S. F., SCHAFER & H. POPPER. 1966. Fat-storing cells (lipocytes) in human liver. *Arch. Pathol.* 82: 447.
- BROUSSARD, G. 1969. Patologia da resine poliviniliche. *Med. Bull.* 52(2): 102.
- BRÜHN, W. 1966. Zwischenfälle und Komplikationen bei der Laparoskopie und gezielten Leberpunktion (Ergebnis einer Umfrage). *Deut. Med. Wochenschr.* 91: 2397.
- CANADA. DEPARTMENT OF NATIONAL HEALTH AND WELFARE, INDUSTRIAL HEALTH DIVISION. 1949. A guide to the diagnosis of occupational diseases. King's Printer, Ottawa, Ontario, Canada.
- CARR, J., R. M. BICKSON, J. F. VITCIA & J. C. KRANTZ, JR. 1949. Anesthesia. XXIV. Chemical constitution of hydrocarbons and cardiac automaticity. *J. Pharmacol. Exptl. Ther.* 97: 1.
- CHATELAIN, A. & P. MORTILLO. 1967. Un syndrome d'aéro-ostéolyse d'origine professionnelle et de constation nouvelle en France. *J. Radiol. Electrol. Med. Nucl.* 48: 277.
- CHEMICAL ECONOMICS HANDBOOK. September 1973. Vinyl Chloride and Polyvinyl Chloride Resins. Stanford Research Institute, Menlo Park, Calif.
- CHIMIEWIRTSCHAFT IN ZAHLEN. VERBAND DER CHEMISCHEN INDUSTRIE E.V., Ed. 1973. 15th ed. Econ-Verlag, Düsseldorf, West Germany.
- CIESOWITZ, M. B. & C. L. HARR. 1962. The smaller halogenated aliphatic hydrocarbons. *Ann. Rev. Pharmacol.* 2: 363.
- CHIEF INSPECTOR OF FACTORIES. 1953. Annual Report for the Year 1951. Cmd. 8772. H.M.S.O., London, England.
- COOK, W. A., P. M. GUYER, B. D. DINMAN & H. J. MAGNUSON. 1971. Occupational acroecolysis. II. An industrial hygiene study. *Arch. Environ. Health* 22: 74.
- CORDUA, J. M., C. FURZA, M. J. LITANI & A. SEVIAN. 1966. Acro-ostéolyse et lésions cutanées associées chez deux ouvriers affectés au nettoyage d'autoclaves. *Cahiers de Médecine du Travail (Association Professionnelle belge des Médecins du Travail)* 4 janvier 1966.
- CULLERT, J. L., JR. & M. N. JOHNSON. 1974. Angiosarcoma of liver in the manufacture of polyvinyl chloride. *J. Occupational Med.* 16: 150.
- CROLL, M. N., L. W. BRADY, I. BROPSKY & L. STANTON. 1965. A new agent for splenic scanning. *BNHP. Radiology* 84: 492.
- DALLGREN, S. 1961. Thoracic tumors. A review of the literature and report of two cases. *Acta Pathol. Microbiol. Scand.* 53: 147.
- DANILEVSKII, S. L. & N. M. EGOROV. 1961. Voprosy toksikologii v khimii vysokomolekulyarnykh soedinenii. *Gigiena Truda Prof. Zabolevaniya* 5(9): 26.
- DANZIGER, H. 1960. Accidental poisoning by vinyl chloride: report of two cases. *Can. Med. Assoc. J.* 82: 828.
- DEUSE, D. E. & R. J. JOYNER. 1969. Vinyl acetate: a study of chronic human exposure. *Am. Ind. Hyg. Assoc. J.* 30: 449.
- DIEHMANN, W. B. & H. W. GIERKE. 1969. Toxicology of drugs and chemicals. Academic Press, New York, N.Y.
- DIVONVILLI, DR. & A. M. FLECHER. 1953. Etude toxicologique expérimentale des résines polyvinyls. *Service Médical des Manufactures de Saint-Marcel, Vernon, avril 1953. As cited in Truffert, 1959.*
- DIMOV, D. & T. BIRNIG. 1971. Profesionalna akroosteoлиза. *Arhiv. Hig. Rada Toksikol.* 22: 53.
- DINMAN, B. D., W. A. COOK, W. M. WHITHOUSE, H. J. MAGNUSON & TH. DIERICK. 1971. Occupational acroecolysis. I. An epidemiological study. *Arch. Environ. Health* 22: 61.
- DUNSON, V. N., B. D. DINMAN, W. M. WHITHOUSE, A. N. M. NASR & H. J. MAGNUSON. 1971. Occupational acroecolysis. III. A clinical study. *Arch. Environ. Health* 22: 83.
- DUMAS-CHATELAIN, H. 1972. Kunststoffe I. Aufbau und Eigenschaften-Kunststoffsorten-Anwendungen. 2nd ed. VDI-Verlag Düsseldorf, West Germany.
- DUMAS-CHATELAIN, H. 1973. Kunststoffe II. Aufbereitung-Oberflächenveredelung. 2nd ed. VDI-Verlag Düsseldorf, West Germany.
- DUMES, I. J. & R. J. VASI. 1933. Occupational hazards and diagnostic signs. U.S. Department of Labor, Bureau of Labor Statistics. Bulletin No. 382. U.S. Government Printing Office, Washington, D.C., p. 49. As cited in Danziger, 1960.
- DUMES, I. J. & R. J. VASI. 1941. Occupational hazards and diagnostic signs. U.S. Department of Labor, Division of Labor Standards. Bulletin No. 41 (revised). U.S. Government Printing Office, Washington, D.C., p. 65. As cited in Danziger, 1960.

48. DUGOIS, P., P. AMBLARD, B. DE BIGNICOURT & J. LEGRAND. 1972. Acropathie polyvinylque professionnelle. *Bull. Soc. Franc. Dermatol. Syphiligr.* 79: 197.
49. ECKARDT, R. E. & R. HINDIN. 1973. The health hazards of plastics. *J. Occupational Med.* 15: 808.
50. EDMONDSON, H. A., R. L. PETERS, H. H. FRANKEL & S. BOROWSKI. 1967. The early stage of liver injury in the alcoholics. *Medicine* 46: 119.
51. EUROPA-CHIMIE. 1972. Handelsblatt GmbH. Düsseldorf, West Germany.
52. FARHAT, L. T. 1957. Industrial toxicology. 2nd ed. The Williams & Wilkins Company, Baltimore, Md.
53. FILATOVA, V. S. & E. S. GRONSHERG. 1957. Sanitarno-gigienicheskie usloviya truda v proizvodstve poliklorovinilovoi smoly i neryi ikh ozdorovleniya. *Gigiena Sanit.* No. 1: 38.
54. FILATOVA, V. S., L. I. BALAKRINOVA & E. S. GRONSHERG. 1958. Gigienicheskiye kharakteristika proizvodstva kloristovo vinila. *Gigiena Truda Prof. Zabolevaniya* 2(1): 6.
55. FILATOVA, V. S., E. S. GRONSHERG, N. A. SMIRNOVA, E. A. STULOVA & I. V. ORUSHEVICH. 1965. Voprosy gigieny truda i sostoyaniya zdorov'ya rabotnikh, zanyatykh na proizvodstve latskanovo polivinilkhlorda. *Gigiena Truda Prof. Zabolevaniya* 9: 9.
56. FILATOVA, V. S. & V. A. ANTONOVYUZINSKO. 1971. Gigienicheskiye usloviya truda i professional'naya zabolevayemost' rabotnikh proizvodstva suspenszionnogo polivinilkhlorda v dinamike za ryad let. *Gigiena Truda Prof. Zabolevaniya* 15(4): 32.
57. FISCHER, J., H. MUNSCHUSK & R. WOLF. 1965. Milzszintigraphie mit 1-Bromomercuri (²⁰¹Hg)-2-Hydroxypropyl (BHMHP). *Fortsehr. Gebiete Roentgenstrahlen Nuklearmed.* 103: 349.
58. FISCHER, J., R. WOLF & H. GANN. 1973. Die Milzszintigraphie. *Dtsch. Ärztschl.* No. 7: 401.
59. GALT, E. A. 1969. Tumors of the liver. *In* Diseases of the Liver. L. Schiff, Ed. 3rd ed. J. B. Lippincott Company, Philadelphia, Pa.
60. GRISWOLD, C. T. 1971. Acro-osteolysis in PVC workers. *Med. Bull. Stand. Oil Co.* 31(1): 49.
61. GÜNTHER, O. 1936. Die Kunststoffe und ihre arbeitsmedizinische Bedeutung. *Zentr. Arbeitsmed.* 6: 156.
62. HARRIS, D. K. 1953. Health problems in the manufacture and use of plastics. *Brit. J. Ind. Med.* 10: 255.
63. HARRIS, D. K. & W. G. F. ADAMS. 1967. Acro-osteolysis occurring in men engaged in the polymerization of vinyl chloride. *Brit. Med. J.* 3: 712.
64. HESSCHUR, D., Ed. 1972/1973. Gesundheitsgefährliche Arbeitsstoffe. Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten (Maximale Arbeitsplatz-Konzentrationen). Verlag Chemie, Weinheim, West Germany.
65. HIRSH, K. 1963. Polyvinylchlorid. *In* Ullmanns Enzyklopädie der technischen Chemie. W. Foerst, Ed. 3rd ed. Vol. 14. Urban & Schwarzenberg, München, West Germany.
66. HIRVITZ & TISSER. 1959. Quelques observations d'exposition et d'intolérance aux dérivés vinyliques et aux résines éthyliques. *Arch. Maladies Prof.* 20: 61.
67. JAFFE, E. 1970. Portal hypertension in the presence of normal liver morphology. *Ann. N.Y. Acad. Sci.* 170(1): 115.
68. INTERNATIONAL LABOUR OFFICE. 1971. Encyclopaedia of Occupational Health and Safety. Geneva, Switzerland. Vol. I: 347.
69. INTERNATIONAL LABOUR OFFICE. 1972. Encyclopaedia of Occupational Health and Safety. Geneva, Switzerland. Vol. II: 1467.
70. JARIS, D. D. 1963. Halogenated hydrocarbons: I. Aliphatic. *In* Industrial Hygiene and Toxicology. Vol. II. Toxicology. D. W. Fassett & D. D. Irish, Eds. Interscience Publishers, New York, N.Y.
71. JAVITT, N. B. 1970. Clinical and experimental aspects of sulphobromophthalicin and related compounds. *In* Progress in Liver Disease. H. Popper & F. Schaffner, Eds. Vol. III. Grune & Stratton, New York, N.Y.
72. JUDAH, J. D., A. E. M. McLEAN & E. K. McLEAN. 1970. Biochemical mechanisms of liver injury. *Ann. J. Med.* 49: 609.
73. JÜNG, S. & C. E. LAWGE. 1972. Sklerodermieartige Hautveränderungen, Raynaud-Syndrom und Acroosteolysen bei Arbeitern der PVC-herstellenden Industrie. *Deut. Med. Wochschr.* 97: 1922.

74. KELLY, R. E. 1973. Vinyl chloride and acroosteolysis. *Occupational Med.* 15: 858.
75. KLAYSKIN, G. 1969. Toxic and drug induced hepatitis. *Schiff, Ed. 3rd ed.* J. B. Lippincott Company, Philadelphia, Pa.
76. KLINGE, O. & H. W. ALTMANN. 1971. Morphologie und Med. Wochschr. 113: 1529.
77. KLINGE, T., H. SOMMERSCHEID & A. FLATMARK. 1970. The correlation of portal hypertension for workers exposed to vinyl chloride. *Arch. Intern. Hyg. Assoc.* 33: 19.
78. KRAMER, C. G. & J. E. MUTCHLER. 1972. The correlation of portal hypertension for workers exposed to vinyl chloride. *Arch. Intern. Hyg. Assoc.* 33: 19.
79. LANGF, C. E., S. JÜNG, G. STIN & G. VILTMANN. 1972. Die sogenannte Vinylchlorid-Krankheit—eine berufsbedingte Systemerkrankung. *Med. Wochschr.* 97: 1922.
80. LEAKE, C. D. 1934. The role of pharmacology in the therapy. *J. Am. Med. Assoc.* 102: 1.
81. LEPAUX, R. 1966. Chemie und Toxikologie der Kunststoffe. *West Germany.*
82. LÉFÈVRE, M. J. 1972. Internationales Symposium der Industrie, 27-29 April 1972, Ludwigshafen. As cited in Stein *et al.*, 1973.
83. LEHMANN, K. B. & F. FLURY. Eds. 1938. Toxikologische Lösungsmittel. Julius Springer, Berlin, Germany.
84. LESTER, D., L. A. GREENBERG & W. R. ADAMS. 1972. Repeated exposures of humans and rats to vinyl chloride. *Arch. Intern. Hyg. Assoc.* 33: 19.
85. LINDNER, H. 1973. Laparoscopy in alcohol-induced liver disease. *Gastroenterology* 64: 842.
86. MACSWELN, R. N. M., J. M. VETTERS, S. K. ROYSTON & A. T. SANBORN. 1973. Haemangio-endothelioma. *J. Pathol.* 109: 39.
87. MALTIM, K. E. & R. L. ZIEGLER. 1964. Industrial diseases in the Production and Processing of Plastics. Elsevier, Amsterdam, The Netherlands.
88. MALTONI, C., M. CRESPI & P. J. R. BURCH, Eds. 1973. On Cancer Detection and Prevention. Bologna, Medica Int. Congr. Ser. No. 275.
89. MARIN, A., J. STRAUSS, R. MICHELIS, J. P. BENOIT, & C. PIERRE. 1967. Acro-osteolysis d'origine professionnelle (Travail contre le Rhumatisme, séance du 18 janvier 1967).
90. MARKOWITZ, S. S., C. J. McDONALD, W. FETTER, & C. PIERRE. 1967. Occupational acroosteolysis. *Arch. Dermatol.* 106: 2.
91. MARSTELLER, H. J., W. K. LEBACH, R. MÜLLER, ROHNER & G. VILTMANN. 1973. Chronisch-toxisch in der PVC-Produktion. *Deut. Med. Wochschr.* 98: 291.
92. MASTROMATTEO, E., A. M. FISHER, H. CHRISTIE & J. S. 1974. Toxicity of vinyl chloride to laboratory animals. *Bull. Hyg.* 36: 244.
93. MAYERS, M. R. 1969. Occupational Health. The Williams & Wilkins Company, Baltimore, Md.
94. MCCORD, C. P. 1970. A new occupational disease. *Occupational Med.* 12: 234.
95. MINKUS, W. P., H. A. EDMONDSON, R. L. PETER, & R. E. ECKARDT. 1965. Extra- and intrahepatic portal hypertension (hepatoportal sclerosis). *Ann. Surg.* 162: 602.
96. MISCHKE, V., H. J. STREHMANN & S. SCHULTE. 1973. Vinylchlorid-Polymerisate und/oder deren Begleitheile. *Deut. Med. Wochschr.* 98: 425.
97. MORRIS, J. S., T. HUNT & A. E. READ. 1972. Scleroderma. *Ann. Rheumatic Diseases* 31: 316.
98. NEFTLINSKY, A. & W. J. FINE. 1961. Neoplasms of the thorax. *Am. J. Clin. Pathol.* 35: 422.
99. OERTTEL, H. 1963. Gewerbetoxikologie und Physiologie. *In* Ullmanns Enzyklopädie der technischen Chemie. 6th ed. Vol. 14. Urban & Schwarzenberg, München, West Germany.

- Chemie, W. Foerst, Ed. Vol. 14: 306. Urban & Schwarzenberg, München, West Germany.
100. OSTER, R. H., C. J. CARR & J. C. KRANTZ. 1947. Anesthesia. XXVII. Narcosis with vinyl chloride. *Anesthesiology* 8: 359.
101. OSTROMAYUK, H. 1967. Vinylchlorid. In *Ullmanns Encyklopädie der technischen Chemie*, W. Foerst, Ed. Vol. 18: 87. Urban & Schwarzenberg, München, West Germany.
102. OTTO, P., P. H. KRAUSE, H. G. JUSTER, F. W. SCHMIDT & K. H. GLATZL. 1972. Idiopathische portale Hypertension. *Med. Klin. (München)* 67: 447.
103. PANCICICIANI, L., & C. SASSI. 1955. Rischio e patologia professionale nella produzione e nella lavorazione di alcune materie plastiche. *Med. Lavoro* 46: 14.
104. PATTY, F. A., W. P. YANT & C. P. WARR. 1930. Acute response of guinea pigs to vapors of some new commercial organic compounds. V. Vinyl chloride. *Public Health Rep.* 45: 1963.
105. PATTY, F. A., Ed. 1958. *Industrial Hygiene and Toxicology*. Vol. I. General Principles, 2nd ed. Interscience Publishers, New York, N.Y.
106. PATTY, F. A., Ed. 1963. *Industrial Hygiene and Toxicology*. Vol. II. Toxicology. D. W. Fassett & D. D. Irish, Eds. 2nd ed. Interscience Publishers, New York, N.Y.
107. POPLIS, A. S. & C. D. LYAKE. 1933. The anesthetic action of vinyl chloride. *J. Pharmacol. Exptl. Ther.* 48: 284.
108. POULIN, E., J. CHRISTIE, A. COHEN & B. SULLIVAN. 1962. Idiopathic presinusoidal portal hypertension (Banti's syndrome). *Ann. Internal Med.* 56: 624.
109. POPPER, H., J. PARINETTO, F. SCHAFNER & V. PLATZ. 1961. Studies on hepatic fibrosis. *Lab. Invest.* 10: 265.
110. POPPER, H. 1968. Pathology of portal hypertension. In *The Therapy of Portal Hypertension*, N. G. Markoff, Ed. Georg Thieme Verlag, Stuttgart, West Germany.
111. POPPER, H. & F. HUTTNER. 1970. Hepatic fibrogenesis and disturbance of hepatic circulation. *Ann. N.Y. Acad. Sci.* 170: 88.
112. POPPER, H. & S. UBERSKUNO. 1970. Hepatic fibrosis. Correlation of biochemical and morphologic investigations. *Am. J. Med.* 49: 707.
113. PUSIN, G. A. 1965. O porazhenii pecheni i zhenykh pri zlozhabnykh zanyatiykh v proizvodstve neftnykh widow plastmass. *Sov. Med.* 28: 132.
114. PVC producers see good business ahead. 1969. *Chem. Eng. News* August 4: 18.
115. QUORNS, H. 1959. *Gesundheitsgefahren in der Kunststoffindustrie*. Johann Ambrosius Barth Verlag, Leipzig, East Germany.
116. RAMALINGASWAMI, V., K. L. WIG & S. K. SAMA. 1962. Cirrhosis of the liver in Northern India—a clinicopathological study. *Arch. Internal Med.* 110: 350.
117. RAMALINGASWAMI, V. & N. C. NAYAK. 1970. Liver disease in India. In *Progress in Liver Disease*, H. Popper & F. Schaffner, Eds. Vol. III: 222. Grune & Stratton, New York, N.Y.
118. RAPPAPORT, A. M., M. KROTHAUCH, R. G. BLANK & S. OJIMA. 1970. Hepatic microcirculatory changes leading to portal hypertension. *Ann. N.Y. Acad. Sci.* 170: 48.
119. RAVINNA, P. 1940. Banti syndrome (fibrocongestive splenomegaly). Definition, classification and pathogenesis. *Arch. Internal Med.* 66: 679.
120. REISCHUK, H. 1970. The role of the liver in drug metabolism. *Ann. J. Med.* 49: 617.
121. RIVEROUDS, T. B. & A. G. REIDICK. 1965. Hepatic hemodynamics and portal hypertension. In *Progress in Liver Disease*, H. Popper & F. Schaffner, Eds. Vol. II: 457. Grune & Stratton, New York, N.Y.
122. RORFF, F. 1957. *Arsen, Leber, Tumoren (Hämangiendutheleiom)*. Z. Krebsforsch. 61: 468.
123. ROUSSELOT, L. M. 1940. The late phase of congestive splenomegaly (Banti's syndrome) with hematocrit but without cirrhosis of the liver. Further observations on the etiology of portal venous syndrome. *Surgery* 8: 34.
124. SAMA, S. K., S. BHASKARA, N. GANI NATH, J. K. TALWAR, N. C. NAYAK, B. N. TANIGUCHI & K. L. WIG. 1971. Noncirrhotic portal fibrosis. *Ann. J. Med.* 51: 160.
125. SARGENT, A. M., H. TAVAKIS, F. B. ISAKHIS & J. DA SILVA HORTA. 1972. Some clinical and laboratory findings in patients injected with thorium dioxide. Study of 155 cases. *Ann. J. Gastroenterol.* 57: 301.
126. SCHAFNER, F. & H. POPPER. 1963. Capillarization of hepatic sinusoids in man. *Gastroenterology* 44: 239.
127. SCHUMANN, O. 1934. Ueber die Herzwirkung einiger Inhalationsanästhetika. *Medizin & Chemie. Abhandlungen an den Chemisch-chemischen Forschungen. Leverkusen, West Germany*.
128. SCHUMANN, O. 1938. In *Toxikologie und Hygiene der technischen Lebensmittel*, K. B. Lehmann & F. Flury, Eds. p. 130 (Monochloräthylen). Julius Springer, Berlin, Germany.
129. SCHUMM, M. 1968. Zur Leberhistologie der verschiedenen Formen des portalen Hochdrucks. In *The Therapy of Portal Hypertension*, N. G. Markoff, Ed. Georg Thieme Verlag, Stuttgart, West Germany.
130. SCHUMM, F. W. 1969. Geringer Aktivitätsanstieg der Serum-Transaminasen. *Deut. Med. Wochenschr.* 94: 2230.
131. SIESSACK, H., L. STOCKINGER & F. WILKINS. 1967. Adventitious connective tissue cells in the space of Disse and their relation to fibre formation. *Rev. Intern. Hepatol.* 17: 855.
132. SIMMONS, H. R., A. L. WILHELM & M. H. BIGLOW. 1949. *Handbook of Plastics*. 2nd ed. D. Van Nostrand Co., Inc. Princeton, N.J.
133. SLATER, T. F. 1972. Free radical mechanisms in tissue injury. Pion Ltd. London, England.
134. SOROS, G. L. & H. A. BATHIFORA. 1972. Thorotrast-induced hepatoma. *Cancer* 30: 1252.
135. SUTTU, H. F., Jr. 1956. Improved communication. Hygienic standards for dairy inhalation. *Am. Ind. Hyg. Assoc. J.* 17: 129.
136. SUTOWAY, R. D., A. H. HAGENSTADT, L. J. SIMONHEIT *et al.* 1971. Observer error and sampling variability tested in evaluation of hepatitis and cirrhosis by liver biopsy. *Am. J. Digest. Diseases* 16: 1082.
137. SUTUS, G., S. JÜLL, C.-E. LANG & G. VALLMAY. 1973. Bandförmige Oskolyss in den Endphalangen des Handgelenks. *Fortschr. Gebiete Roentgenstrahlen* Nuklearmed. 118: 66.
138. SUTTOR, R. J. 1965. Fibrogenesis along the hepatic sinusoids in carbon-tetrachloride-induced cirrhosis. An electron microscopic study. *Exptl. Mol. Pathol.* 4: 357.
139. STUTZ, F. H., D. C. TOWSEY & J. BLOM. 1973. Hemangiosarcoma and pathologic rupture of the spleen. *Cancer* 31: 1213.
140. STUTZ, F. H., I. DRYANAS & M. VALASKA. 1963. Contributii la studiul imbolnavirii produse de clorura de vinil. *Med. Internu* 15: 967.
141. STUTZ, F. H., I. DRYANAS & M. VALASKA. 1967. Etude des maladies dues au chlorure de vinyle. *Med. Lavoro* 58: 261. *Abstr. Hyg.* 43: 430 (1968).
142. SUMIYASHI, W. H. J. 1974. Chronic active liver disease reexamined: prognosis hopeful. *Gastroenterology* 66: 450.
143. SYMANS, F. B., J. SYMANS & G. J. GRAY. 1974. Hemangiendutheleial carcinoma of the liver and hemochromatosis. *Arch. Pathol.* 97: 39.
144. SYMANS, H. Ed. 1963. *Handbuch der gesamten Arbeitsmedizin*. F. W. Becker, Ed. Vol. IV/2: Arbeitshygiene. Urban & Schwarzenberg, München, West Germany.
145. TANDON, R. N., R. LAKSHMINARAYANAS, S. BHASKARA, N. C. NAYAK & S. K. SAMA. 1970. Ultrastructure of the liver in non-cirrhotic portal fibrosis with portal hypertension. *Gut* 11: 905.
146. THOMAS, W. A., G. KLASIKS & W. W. L. GUNN. 1959. Portal hypertension and bleeding esophageal varices. Their occurrence in the absence of both intrahepatic and extrahepatic obstruction of the portal vein. *New Engl. J. Med.* 261: 209.
147. THOMAS, T. R., F. OWLS & V. K. ROWE. 1961. The toxicity of vinyl chloride as determined by repeated exposure of laboratory animals. *Am. Ind. Hyg. Assoc. J.* 22: 354. *Bull. Hyg.* 37: 357.
148. THOMAS, S. L., N. P. THIRUMANIAN, S. V. JAYISA & L. A. KOTTON. 1970. *Chlorvinil i meropriyatya po ikh ozonokleniya pri prevarivani i teplovoyan*. Khlorvinilovykh plasticheskikh mass. *Gigiena Sanit.* 10: 18.
149. THOMAS, L. 1959. Aspects toxicologiques de l'industrie des matières plastiques. *Arch. Maladies Profes.* 20: 41.
150. U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE—Public Health Service. *Center for Disease Control*. 1974. *Annual mortality of the liver among polyvinyl chloride workers—Kentucky*. *Morbidity Mortality Weekly Rep.* 23(6): 49.

151. VIOLA, P. L. 1970. Pathology of vinyl chloride. *Med. Lavoro* 61: 174.
152. VIOLA, P. L., A. BIGOTTI & A. CAPUTO. 1971. Oncogenic response of rat skin, lungs, and bones to vinyl chloride. *Cancer Res.* 31: 516.
153. VOLKHEIMER, G., F. SZARVAS, F. AL ARESIL, H. JOHN & S. WACHTEL. 1961. Aufsaugung von Polyvinylchlorid-Pulver durch die Dünndarmschleimhaut und deren Abtransport durch die Chylusgefäße. *Deut. Gesundheitsw.* 16: 1727.
154. VON OETTINGEN, W. F. 1964. The Halogenated Hydrocarbons of Industrial and Toxicological Importance. Elsevier Publishing Company, Amsterdam, The Netherlands.
155. WAGNER, H. N., JR., J. G. McAFEE, I. M. WEINER, M. ILIO, J. MARTINEZ & W. P. CAINE, JR. 1963. The use of Hg-203 labeled bromomercurihydroxypropene (BMHP) in radioisotope scanning of the spleen. *J. Nucl. Med.* 4: 190.
156. WEPLER, W. 1971. Zur Morphologie chronischer Leberkrankheiten. *Leber Magen Darm* 1: 45.
157. WILLIAMS, G. M., J. M. ELLIOTT & J. H. WEISBURGER. 1973. Carcinoma after malignant conversion in vitro of epithelial-like cells from rat liver following exposure to chemical carcinogens. *Cancer Res.* 33: 606.
158. WILLIAMS, R. T. 1959. *Detoxication Mechanisms. The Metabolism and Detoxication of Drugs, Toxic Substances and Other Organic Compounds.* 2nd edit. Chapman & Hall Ltd. London, England.
159. WILSON, R. H. & W. E. McCORMICK. 1960. Plastics. The toxicity of synthetic resins. *A. M. A. Arch. Ind. Health* 21: 536.
160. WILSON, R. H., W. E. McCORMICK, C. F. TATUM & J. L. CRECHIL. 1967. Occupational acroosteolysis. Report of 31 cases. *J. Am. Med. Assoc.* 201: 577.
161. WIRTSCHAFER, Z. T. & M. W. CROSNY. 1964. Free radical mechanism for solvent toxicity. *Arch. Environ. Health* 9: 185.
162. WITTMAN, I. 1966. *Peritoneoscopy. Vols. I & II.* Akadémiai Kiadó. Publishing House of the Hungarian Academy of Sciences. Budapest, Hungary.
163. ZAPP, J. A., JR. 1962. Toxic and health effects of plastics and resins. *Arch. Environ. Health* 4: 125.

DISCUSSION

DR. H. BUCHTER (*University of Cologne, Germany*): It would seem better for statistical investigations on the reported liver function tests if you could compare the results with similar tests done before the workers came into vinyl chloride manufacturing. Second, in one of the two plants in Germany with incidence of vinyl chloride disease, there is much alcohol intake in the workers. Until now no liver damage has been found.

Third, in regard to low lung-function tests, we have not confirmed a pathological diagnosis at all.

There should be more discussion about the cause of this disease, i.e., whether it is vinyl chloride alone or another agent which may get together with technical vinyl chloride, or vinyl acetate or additives. This may be important in the causation of angiosarcoma.

Last, it has been suggested that liver biopsy be done only after peritoneoscopy because of the risk of bleeding.

DR. M. L. NEWHOUSE: May I ask Dr. Ielbach how he selected his patients and how big was the exposed group he selected them from?

DR. W. K. IELBACH (*University of Bonn, Germany*): I will begin with the answer to the second part of the question. The whole group is, as far as we know, about 180 workers in this plant. We were alerted in December 1972 by Dr. Veltman's group that patients being examined for skin and bone disease also had signs of liver disease. We examined these patients.

Later we examined all patients we could bring to the clinic. It is, of course, a selected group and does not permit us to draw conclusions as to percentages.

B. Experimental

PRELIMINARY VIN

R. E.

Vinyl chloride and angiosarcoma and in workers exposed to low levels of VCM exposed via inhalation. Angiosarcomas, 2% rats exposed to 5 days/week for death.

No information at the present time of VCM and p to low levels of VCM exposed via inhalation conceived the idea pathway, and if a jugation of VCM pected. Therefore evaluating these p

Kinetic Studies of

Male Sprague-Dawley rats were exposed to 8 ppm (0.13-2.99 n depicts the closed, 4 rats were concu only the nares of t ber. Expired carb absorption on an was removed from manometer fitted dual syringe pump

The chamber s in-line Miran-I int

* These studies Association.

