

(b)

Acute Inhalation Toxicity of Vinyl Chloride to Laboratory Animals

E. MASTROMATTEO, M.D., A. M. FISHER, Ph.D.,
H. CHRISTIE, B.Sc., and H. DANZIGER, M.D.

*Division of Industrial Hygiene, Ontario Department of Health, Toronto, Ontario;
Department of Physiological Hygiene, School of Hygiene, University of Toronto,
Toronto, Ontario; and Welland County General Hospital,
Welland, Ontario*

Introduction

FIRST prepared in 1833, vinyl chloride has come into increased use in the last decade in the production of polyvinyl chloride resins. The fire and explosion hazard of the monomer and its narcotic properties on acute exposure are well known. As an industrial material, however, it has been regarded as of moderate toxicity. A Threshold Limit Value of 500 ppm was established by the American Conference of Governmental Industrial Hygienists.

No industrial fatalities associated with its use have previously been reported. It was therefore of special interest when two fatalities occurred in a plant handling vinyl chloride. This plant is engaged in the polymerization of the gaseous monomer with the aid of catalysts. One worker died while cleaning out a polymerization vessel and the other while working in a pit. Details of these two fatalities have been reported by one of the authors.¹

Published information on the acute toxicity of vinyl chloride to experimental animals proved scanty. Patty, Yant and Waite² in 1930 reported on the acute response of guinea pigs to vinyl chloride. They found that exposure to 20 to 40 per cent in air killed guinea pigs in a short time; exposure to ten per cent was endured for several hours. Congestion and edema of the lungs with hyperemia of the liver and kidneys were noted.

Peoples and Leake³ found the minimal anesthetic range of vinyl chloride in mice exposed for ten minutes was 8 to 12 per cent in air; 25 to 30 per cent was fatal in ten minutes. Dogs and rabbits were anesthetized within one minute when exposed to a concentration of about 18 per cent in air. Recovery was very rapid with no apparent untoward effect even after prolonged anesthesia.

Lehmann and Flury⁴ noted that vinyl chloride was highly narcotic but had a wide margin between its narcotic and lethal concentrations. Its local irritating effect and its toxicity were de-

scribed as very slight. Elimination from the body was very rapid. These authors quoted the work of Schaumann⁵ with experimental animals. Mice, rats and dogs were able to endure repeated narcosis without any pathological evidence of liver or kidney damage. According to Schaumann, concentrations of seven to ten per cent in air would be narcotic to humans in a short time and 12 per cent would be dangerous.

Vinyl chloride has been studied as a possible anesthetic agent. Oster and associates⁶ noted that incoordinated muscular activity of the extremities and cardiac arrhythmias of a serious nature were observed in all six dogs used in their experiment. Carr and others⁷ found that vinyl chloride produced sensitization of the myocardium in three of seven dogs in which it was administered as an anesthetic agent.

Two cases of vinyl chloride gassing occurred in a factory in Great Britain where polyvinyl chloride resin was being made. These were reported by the Chief Inspector of Factories.⁸ In one case a process worker was standing outside a polymerization vessel and washing it with a water stream. After ten minutes of this he suddenly collapsed across the open manhole. Artificial resuscitation was successfully applied. Subsequent symptoms experienced by this worker were tightness of the chest, nausea, abdominal pain and headache. The second case occurred in a maintenance worker who was overcome while repairing a vinyl chloride leak. Both men were hospitalized. Another worker whose hands were accidentally sprayed with vinyl chloride liquid under pressure developed a burn. This latter case was reported by Harris.⁹ Such burns may occur with other highly volatile materials when sprayed onto the hands as a liquid under pressure.

Filatova and Gronsberg¹⁰ in 1957 reported on hygienic conditions in a polyvinyl processing plant in the U.S.S.R. The plant was engaged in the polymerization of vinyl chloride by means of

Industrial

a catalyst. A concentration of from about 20 Threshold Limit Value (TLV) to 100 TLV was described in v. Because of above it was tion studies i cially availab for comparis information o cal changes in animal.

Experimental

Vinyl Chloride in a metal cylinder grade with th

Specific gravity a
Water content
Boiling range °C
Hydrogen chloride
Acetylene
Acetaldehyde
Iron
Purity (per cent)
Impurities (per c

Pure vinyl temperature; a sweetish od and its explo cent. It has t

Animals: A in the study. from the san standard com way. All were

Equipment. was 56.6 liter window and : leased in ga valve on the through cont brated Fisher by motor at a meter. The were combin glass Y-tube ing to the a continuing s Calculations duce the foll chloride in ai cent. All com port as per c

Chloride

i.D.,
J.

into, Ontario;
of Toronto,
il,

ination from the bo
hors quoted the wor
mental animals. Mic
endure repeated na
gical evidence of liv
ding to Schauman
> ten per cent in a
ns in a short time an
rous.

studied as a possib
associates⁴ noted th
tivity of the extrem
s of a serious natur
s used in their experi
nd that vinyl chloride
the myocardium in
h it was administered

de gassing occurred in
where polyvinyl chlor
these were reported by
torics.⁶ In one case a
g outside a polymeri
with a water stream
e suddenly collapsed
Artificial resuscitation
ubsequent symptoms
were tightness of the
in and headache. The
maintenance worker
siring a vinyl chloride
lized. Another worker
ntally sprayed with
pressure developed a
reported by Harris.
other highly volatile
the hands as a liquid

in 1957 reported on
polyvinyl processing
plant was engaged in
chloride by means of

a catalyst. Air sampling was done and the concentration of vinyl chloride was found to vary from about 20 ppm to about 315 ppm. (U.S.S.R. Threshold Limit Value: 1 mg/liter or about 400 ppm). A spastic type blood vessel disorder was described in workers from this plant.

Because of the recent fatalities mentioned above it was decided to undertake acute inhalation studies in laboratory animals with commercially available vinyl chloride. This was done both for comparison with earlier work and to provide information on the acute response and pathological changes in more than one species of laboratory animal.

Experimental Materials and Procedures

Vinyl Chloride: The vinyl chloride was supplied in a metal cylinder under pressure. A commercial grade with the following specifications was used:

Specific gravity at 20°C	0.992-0.995
Water content	None
Boiling range °C	-11 to -0.5
Hydrogen chloride content	None
Acetylene	10 ppm maximum
Acetaldehyde	20 ppm maximum
Iron	Essentially iron free
Purity (per cent)	99.5 maximum
Impurities (per cent)	0.5 maximum

Pure vinyl chloride is a colorless gas at room temperature; its boiling point is -13.9°C . It has a sweetish odor. Its flash point is given at -78°C and its explosive limits in air from 4 to 22 per cent. It has the molecular formula $\text{CH}_2=\text{CHCl}$.

Animals: Mice, rats and guinea pigs were used in the study. Test and control animals were taken from the same laboratory stock. All were fed a standard commercial diet and housed in the same way. All were of the same stage of development.

Equipment: The inhalation chamber capacity was 56.6 liters. It was equipped with a viewing window and an inlet tube. Vinyl chloride was released in gaseous form through an adjustable valve on the top of the containing cylinder, then through connecting rubber tubing and a recalibrated Fisher flow-meter. Fresh air was pumped by motor at an adjustable controlled rate through a meter. The streams of air and vinyl chloride were combined at appropriate rates of flow by a glass Y-tube leading through further rubber tubing to the animal chamber inlet, to deliver a continuing stream in the desired proportions. Calculations and adjustments were made to produce the following flow concentrations of vinyl chloride in air for delivery: 10, 20, 30 and 40 per cent. All concentrations are expressed in this report as per cent by volume in air. No determina-

tions of vinyl chloride concentrations were done in the test chamber during the experiments.

Experimental Procedures: Different groups of five mice, five rats and five guinea pigs were placed in the chamber and exposed for thirty minutes to concentrations of 10, 20 and 30 per cent vinyl chloride in air. Similar groups of control animals were maintained but not exposed to vinyl chloride. At the end of thirty minutes exposure the test animals were immediately removed to fresh air. An additional group of five guinea pigs was exposed to a concentration of 40 per cent vinyl chloride in air. In all 65 experimental animals were involved. Observations were recorded on the animals during and after exposure.

The animals which died either during the exposure or after a delay period were autopsied soon after death. Two weeks after the exposure surviving test animals and controls were sacrificed by a blow to the head with the exception of four control rats which were killed by exsanguination.

All animals were examined for gross pathological changes. Tissues were removed from all animals for microscopic examination. The lungs, liver, kidney and heart were removed in all cases. The brain, adrenal, spleen, trachea, lymph nodes, and the eye were removed from representative animals in each group for study. The tissues were preserved in formalin and sections made for staining with hematoxylin-eosin. Special stains were made where indicated.

Observations

Control Animals: No symptoms were exhibited by the 15 control animals and no deaths occurred.

Ten Per Cent Vinyl Chloride: The response of animals exposed to this concentration is recorded below in summary form:

Exposure Time (minutes)	Response
1	Slight irritation in mice and rats.
5	Increased motor activity first in mice, then rats and guinea pigs.
10	Increased motor activity in all species; twitching of extremities in mice.
15	Pronounced tremor, unsteady gait and muscular incoordination in all species.
20	Mice and rats in side position; muscular incoordination, tremors and twitching of extremities in guinea pigs.
25	Mice and rats unconscious, guinea pigs very unsteady but still standing.
30	Mice and rats in deep narcosis; guinea pigs in side position with tremors—also unconscious.

Exposure was stopped and the animals removed to fresh air. All recovered within five minutes.

Twenty Per Cent Vinyl Chloride: The response

20546002

of animals exposed to this concentration is recorded below in summary form:

Exposure Time (minutes)	Response
1	Immediate irritation in mice and rats.
2	Muscular incoordination in mice and rats.
5	Mice and rats on their sides with marked tremors and twitching of the extremities; unsteadiness and muscular incoordination in guinea pigs.
10	Mice and rats unconscious with rapid irregular breathing; guinea pigs unconscious but showing marked twitching.
15	All animals in deep narcosis; respirations irregular and rapid.
20	Deep narcosis; breathing slow and shallow.
25	Breathing ceased in one mouse; frothing at mouth and nostrils in mice and rats.
30	All animals in deep narcosis.

Exposure was stopped and the animals removed to fresh air. Mice and rats recovered faster than the guinea pigs and appeared normal within five minutes except for one mouse which was dead. The guinea pigs continued to show muscular incoordination, unsteadiness on their feet until 20 minutes after removal from exposure.

Thirty Per Cent Vinyl Chloride: The response of animals exposed to this concentration is recorded below in summary form:

Exposure Time (minutes)	Response
—	Mice showed irritation immediately and the rats showed irritation quickly thereafter.
1	Muscular incoordination in mice and rats.
2	Mice and rats in side position with marked tremors and twitching of the extremities; muscular incoordination in guinea pigs with developing paralysis of the extremities.
5	All animals unconscious; rapid irregular respirations in mice and rats with frothing about nose and mouth; twitching of extremities still occurred occasionally in guinea pigs.
10	Respirations stopped in mice; breathing slow and shallow in rats; guinea pigs still exhibited occasional twitching movements of the extremities.
15	Breathing stopped in rats; respirations slow and shallow in guinea pigs; twitching of extremities still present in guinea pigs.
30	Mice and rats still; guinea pigs in deep narcosis with slow shallow breathing.

Exposure was stopped and the animals removed to fresh air. The mice and rats were dead. The guinea pigs took 25 minutes to return to their normal appearance and activity. One guinea pig from this group died within 24 hours following exposure.

TABLE I
Number of Deaths in Different Groups of Mice, Rats and Guinea Pigs Exposed for Thirty Minutes to Varying Concentrations of Vinyl Chloride in Air

Vinyl chloride concentration (per cent by volume in air)	Laboratory animal			Total
	Mice	Rats	Guinea pigs	
10	0/5	0/5	0/5	0/15
20	1/5	0/5	0/5	1/15
30	5/5	5/5	1/5*	11/15
40	—	—	2/5*	2/5
Total	6/15	5/15	3/20	14/30

* A delayed death occurred within 24 hours following exposure.

Forty Per Cent Vinyl Chloride: Only five guinea pigs were exposed to this concentration. Signs of irritation were immediately apparent. Muscular incoordination appeared within seconds. After five minutes all the guinea pigs were unconscious with slow shallow breathing. At the end of the exposure period one guinea pig was dead and the remaining four were in deep narcosis. These survivors took 30 minutes to return to their normal appearance and activity, but one died within the following 24 hours.

The number of deaths occurring in different groups of five laboratory animals exposed for thirty minutes to varying concentrations of vinyl chloride in air is shown in Table I.

Pathological Findings

Gross pathological and histological studies were carried out. The findings are summarized below.

Control Animals: These animals showed no gross or microscopic evidence of damage.

Ten Per Cent Vinyl Chloride: All test animals survived this exposure. They were sacrificed two weeks later. On gross examination there was evidence in mice of slight hyperemia of the lungs. This was less marked in rats. Guinea pigs showed no difference from control animals. On histological examination the mice showed very slight engorgement of the pulmonary vessels. One mouse showed degenerative changes in the tubular epithelium of the kidney with hydropic swelling. The rats showed slight congestion of the capillaries in the lung. Lungs in the guinea pig were also slightly more hyperemic than those of the control animals.

Twenty Per Cent Vinyl Chloride: One mouse died as a result of 30 minutes exposure to this concentration. Pulmonary congestion was evident on gross examination. Microscopically, there was

engorgement of the blood vessels in the alveoli. The kidneys show changes in the epithelium.

The test animals sacrificed in two weeks showed congestion of the lungs; but it was more marked in the guinea pigs. Evidence of pulmonary edema. Some fatty infiltration of one rat. All other normal.

Thirty Per Cent Vinyl Chloride: The response of the animals to exposure revealed evidence of hemorrhagic areas. One guinea pig died. There was marked hemorrhages and the frangible. The microscopically which died included pulmonary blood vessel changes in the lungs. Superficial desquamation was also evident of the mice and rat which had the delay degeneration of the sections stained with

The four survivors two weeks later. There was present with hemorrhage. There was terminal infiltration. There was the appearance of the sections stained with

Forty Per Cent Vinyl Chloride: The four guinea pigs exposed during the exposure hours. The two with congestion of the lungs. This was the appearance of the sections stained with

The three survivors two weeks later. There was hemorrhage and microscopic evidence of the tracheal epithelium. Pathological studies of the brain, heart, spleen, the eyes showed and animals dying. Taken controls at

20546003

BFG03838

engorgement of the blood vessels in the lung with patchy areas of atelectasis and minimal edema. The kidneys showed minimal degenerative changes in the epithelium of the convoluted tubules.

The test animals surviving this exposure were sacrificed in two weeks. On gross examination, congestion of the lungs was present in all species, but it was more marked in the mice and rats than in the guinea pigs. Histologically, there was evidence of pulmonary congestion in exposed animals. Some fatty infiltration was present in the liver of one rat. All other tissues studied appeared normal.

Thirty Per Cent Vinyl Chloride: Gross examination of the animals which died as a result of this exposure revealed congestion of the lungs with hemorrhagic areas. The mice and rats in addition showed congestion of the liver and kidney. The one guinea pig death was delayed. In this animal there was marked congestion of the lungs with hemorrhages and the liver was distended and very friable. The microscopic changes in the animals which died included marked engorgement of the pulmonary blood vessels with edema and hemorrhages in the lungs. The trachea of one rat showed superficial desquamation of the epithelium. Congestion was also evident in the liver and kidney of the mice and rats. The liver of the guinea pig which had the delayed death showed severe fatty degeneration of the liver confirmed with frozen sections stained with Sudan III.

The four surviving guinea pigs were sacrificed two weeks later. Marked pulmonary congestion was present with hemorrhagic areas and edema. In one case there was evidence of secondary bacterial infiltration. The liver in these guinea pigs gave the appearance of fatty infiltration but no fat was demonstrated on frozen section.

Forty Per Cent Vinyl Chloride: One of five guinea pigs exposed to this concentration died during the exposure and another died within 24 hours. The two which died showed marked congestion of the lungs with hemorrhages on gross examination. This was also evident on microscopic examination. The liver of one of these gave the appearance of fatty infiltration, but no fat could be demonstrated on frozen section.

The three surviving guinea pigs were sacrificed two weeks later. Marked congestion of the lungs with hemorrhage was evident on both the gross and microscopic examination. In one guinea pig, the tracheal epithelium was completely absent.

Pathological studies which were made of the brain, heart, spleen, adrenals, lymph nodes and the eyes showed no difference between controls and animals dying as a result of exposure or between controls and the surviving animals sacri-

ficed two weeks after exposure. There was a tendency for the blood to remain unclotted in the animals dying during exposure. This feature was noted in the two human fatalities mentioned earlier but it is not a specific characteristic of vinyl chloride.

Discussion

The response of guinea pigs to inhalation of vinyl chloride was similar to that reported by Patty, Yant and Waite. The guinea pigs in our study, however, tolerated a greater exposure during the experimental period than either the mice or rats. The results in mice agreed closely with those reported by Peoples and Leake. Rats were similar to mice in their response.

The pathological changes in animals which died as a result of exposure were mainly those of vascular engorgement of the lungs with hemorrhages and edema. The severity of these changes varied with the severity of exposure. In the higher concentrations, pulmonary change was marked with severe damage to the tracheal epithelium. Congestion of the liver and kidneys also occurred in test animals. These changes are similar to those reported by Patty and associates.

Evidence of pulmonary congestion was still present in surviving animals sacrificed two weeks after exposure. Patty and associates reported that such changes had disappeared in about eight days.

Degenerative changes in the tubular epithelium of the kidney were noted in one mouse dying as a result of exposure and in one mouse sacrificed two weeks after being exposed. Such changes, however, were minimal and not shown by other animals in the group.

One of the guinea pigs with delayed death following exposure showed severe fatty infiltration of the liver. Changes suggestive of fatty infiltration were observed in other exposed animals but not confirmed by frozen sections stained with Sudan III.

Summary

Separate groups of laboratory animals comprising five mice, five rats and five guinea pigs were exposed in an inhalation chamber to concentrations of 10, 20 and 30 per cent vinyl chloride in air for 30 minutes. An additional five guinea pigs were exposed to a 40 per cent concentration for a similar period of time. The response by each of the three species to these concentrations is noted.

Inhalation of these relatively high concentrations produced narcosis and death. Mice were the most susceptible with guinea pigs considerably more resistant. Rats were similar to mice in their

20546004

response. Exposure to ten per cent vinyl chloride in air produced deep narcosis in mice and rats but no deaths; 30 per cent concentrations killed mice and rats. Exposure of guinea pigs to 20 per cent produced deep narcosis; but three of five guinea pigs survived exposure to 40 per cent concentration.

Animals dying as a result of exposure were autopsied shortly after death. Survivors and control animals were sacrificed two weeks later. Gross pathological and microscopic studies were done and these findings are described.

The principal pathological changes in animals dying from exposure were congestion of the lungs with pulmonary edema and hemorrhages in some, and congestion of the liver and kidneys. Failure of the blood to clot was also observed. One of the three guinea pigs which died following exposure showed severe fatty infiltration of the liver.

Survivors sacrificed two weeks after exposure showed little difference from the control animals. Pulmonary congestion was still evident but liver and kidney congestion was not.

References

1. DANKIGER, H.: Accidental Poisoning by Vinyl Chloride, part of Two Cases, *Canadian Med. Assoc. J.* 82: 525 (1960).
2. PATTY, F. A., W. P. YANT, AND C. P. WAITE: Acute Reaction of Guinea Pigs to Vapors of Some New Commercial Organic Compounds V. Vinyl Chloride, *Pub. Health Rept. U.S.* (August 1950).
3. PROPLES, A. E., AND C. D. LEAKE: The Anesthetic Action of Vinyl Chloride, *J. Pharmacology* 48: 284 (1932).
4. LEHMANN, K. B., & F. FLURY: *Toxicology and Hygiene of Industrial Solvents*, Translated by E. King and H. F. Schmitt, Jr., Williams & Wilkins, Baltimore (1943).
5. SCHNAUMANN, O.: Cited by Lehmann, K. B. and F. Flury.
6. OSTEN, R. H., C. J. CARR, J. C. KRANTZ, AND M. J. SALTWALD: Anesthesia XXVII. Narcosis with Vinyl Chloride, *Anesthesiology* 8: 389 (1947).
7. CARR, J., R. M. BURROWS, J. F. VITCHEL, AND J. C. KRANTZ, JR.: Anesthesia XXIV. Chemical Constitution of Hydrocarbons and Cardiac Automaticity, *J. Pharm. and Therap.* 97: 1 (1948).
8. *Annual Report of the Chief Inspector of Factories for the Year 1961*, London: H.M.S.O. Cmd. 8773 (March 1963).
9. HARRIS, D. K.: Health Problems in the Manufacture and Use of Plastics, *Brit. J. Ind. Med.* 10: 285 (1953).
10. FILATOVA, U. S., AND E. B. GROMOV: Hygienic Working Conditions in Polyvinyl Chloride Tar Plants, *Gigiena* 28-62 (January 1967). Abstracted in *Abst. of World Med.* 66 (July 1967).

BACK ISSUES OF AIHA QUARTERLY

FOR 18 YEARS THE AIHA QUARTERLY was the publication of AIHA, then in 1958 it was changed to be the AIHA Journal with six issues per year instead of four. Now limitation of storage space is forcing us to dispose of all back issues of the Quarterly. For individual issue quantities on hand vary from a low of 8 upward. All issues except March 1946 (Vol. 7, No. 1) and June 1956 (Vol. 17, No. 2) are available as we go to press. Some issues are reprinted without cover. The cost is \$1.50 per issue (\$1.00 each when more than three issues are ordered) from Volumes 7 through 18.

From 1940 to 1946 AIHA Quarterly appeared as a section in *Industrial Medicine and Surgery*, however these issues of these sections have been reprinted and are available as a unit. The complete reprint, Vols. 1-7, costs \$10.00 to members, and \$50.00 to companies or departments of government. Handsomely bound, complete sets of Vols. 1-18 with cumulative index are available for \$250.00 per set.

Address all orders to: George D. Clayton, Executive Secretary, AIHA, 14125 Prevost, Detroit 27, Michigan.

(Back issues of *AIHA Journal*, 1958 to present, may be obtained directly from our publisher, The Williams & Wilkins Company, 428 E. Preston, Baltimore 2, Maryland, at \$1.50 each issue.)

Acute

WILLI

Depart

Introduction

THE element thallium was discovered by Crookes in 1861 and isolated chemically by Lamy. There are no known natural sources of thallium and its principal uses of it to its rodenticidal properties used as a depilating agent also used in the production of optical glass and in the construction of counters.

An examination of the number of published reports of thallium toxicity following the isolation of thallium compounds containing thallium in the depilating action of laboratory animals attributed to thallium (Tl²⁰³) has been reported by Crookes.^{1, 2, 3, 4} Recently, Crookes⁵ have demonstrated that thallium sulfate act as acute thallium poisoning.

The work reported by Crookes⁶ on the acute toxicity of thallium oxide (Tl₂O) following the administration of thallium compounds to several laboratory animals. The ingested compounds was also investigated in three months.

Experimental Methods

The two thallium compounds obtained from the American Chemical Company, Denver, Colorado (thallium acetate, thallium nitrate). The purity of the compounds was relatively high (99.5-99.9%).

This paper is based on work done at the United States Atomic Energy Commission, Rochester Atomic Energy Laboratory.

20546005

BFG03840