


FILE NAME: DuPont (DUP)

DATE: 1967 July 5

DOC#: DUP052

DOCUMENT DESCRIPTION: Letter to Dr. Harriet L. Hardy, Asst. Medical Director in Charge of Occupational Medicine Service, MIT - from G.J. Stopps



July 5, 1967

C-00227

Dr. Harriet L. Hardy
Assistant Medical Director
in Charge of Occupational
Medicine Service
Room 20B-238
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dear Dr. Hardy:

In common with many persons in industries using substantial quantities of asbestos, we are frequently asked for guidance in setting industrial hygiene standards for this mineral. A recent conversation with Dr. Dixon of Pisons (England) suggested that there may be a real difference between the various forms of asbestos in regard to their ability to cause mesotheliomata. My first question would then be, is there such a difference? and does it apply to the other less specific tumors of the lung and gastrointestinal tract also associated with asbestos? Secondly, if there is such a difference in tumorigenicity between the various forms of asbestos, is it of a sufficient degree to be the basis for a recommendation to management on asbestos handling?

Yours sincerely,

ORIGINAL SIGNED BY
G. J. STOPPS

G. J. Stopps, M.B., B.S.
Chief, Physiology Section

GJS/dyb

JJC

6319