

COUNCIL ON PHARMACY AND CHEMISTRY
of the
AMERICAN MEDICAL ASSOCIATION

OFFICE OF THE SECRETARY,

535 N. DEARBORN STREET, CHICAGO 10, ILLINOIS.

November 4, 1947

Robert A. Kehoe, M.D.
Kettering Laboratory of Applied
Physiology,
College of Medicine - Eden Ave.,
Cincinnati 19, Ohio.

Dear Doctor Kehoe:

The Committee on Cosmetics of the American Medical Association is interested in the problem of percutaneous absorption of lead salts through the medium of a hair dye. Our attention has recently been called to a hair pomade containing 3% acetate of sugar lead to be applied two (2) to three (3) times a week.

Your article "Exposure to Lead" v.3 (Feb.) 1947, has prompted us to contact you. Certain conclusions have been reached as to safe levels of ingestion of lead daily. Could these levels be influenced to any marked degree by the continued application of a hair dye or pomade of 3% lead acetate content?

We would appreciate any assistance which you may give us with this problem of percutaneous lead absorption.

Sincerely yours,

Veronica L. Conley

Veronica L. Conley, A.B., M.N.
Corresponding Secretary

COMMITTEE ON COSMETICS

VLC:mts

KE 0015118