

Cardioselective β -Blockers in Patients with Reactive Airway Disease: A Meta-Analysis

Shelley R. Salpeter, MD; Thomas M. Ormiston, MD; and Edwin E. Salpeter, PhD

Objective: To assess the effect of cardioselective β -blockers on respiratory function of patients with reactive airway disease.

Data Sources: Comprehensive searches of the EMBASE, MEDLINE, and CINAHL databases from 1966 to May 2001 and scanning of references of the identified articles and related reviews.

Study Selection: Randomized, blinded, placebo-controlled trials that studied the effects of cardioselective β -blockers on FEV₁, symptoms, and the use of inhaled β_2 -agonists in patients with reactive airway disease were selected. Interventions studied were the administration of a cardioselective β -blocker and administration of β_2 -agonist after the study drug.

Data Extraction: Outcomes measured were the change in FEV₁ from baseline, the number of patients with respiratory symptoms, and the use of inhaled β_2 -agonists with active treatment compared with placebo.

Data Synthesis: Nineteen studies on single-dose treatment and 10 studies on continued treatment were included. Administration of a single dose of a cardioselective β -blocker was associated with a 7.46% (95% CI, 5.59% to 9.32%) decrease in FEV₁ and a 4.63% (CI, 2.47% to 6.78%) increase in FEV₁ response to β -ag-

onist compared with placebo, with no increase in symptoms. Trials lasting from 3 days to 4 weeks produced no significant change in FEV₁ (−0.42% [CI, −3.74% to 2.91%]), symptoms, or inhaler use compared with placebo but maintained an 8.74% (CI, 1.96% to 15.52%) increase in β -agonist response. No significant treatment effect in terms of FEV₁ was found in patients with concomitant chronic obstructive pulmonary disease, whether single doses (change in FEV₁, −5.28% [CI, −10.03% to −0.54%]) or continued treatment (change in FEV₁, 1.07% [CI, −3.3% to 5.44%]) was given.

Conclusions: Cardioselective β -blockers do not produce clinically significant adverse respiratory effects in patients with mild to moderate reactive airway disease. The results were similar for patients with concomitant chronic airways obstruction. Given their demonstrated benefit in such conditions as heart failure, cardiac arrhythmias, and hypertension, cardioselective β -blockers should not be withheld from patients with mild to moderate reactive airway disease.

Ann Intern Med. 2002;137:715-725.

www.annals.org

For author affiliations, see end of text.

See editorial comment on pp 766-767.

β -Adrenergic blocking agents, or β -blockers, are indicated in the management of angina pectoris, myocardial infarction, hypertension, congestive heart failure, cardiac arrhythmia, and thyrotoxicosis and are given to reduce perioperative complications (1–13). Despite clear evidence of the effectiveness and mortality benefit of these drugs, clinicians are often hesitant to administer them in patients with some common conditions for fear of adverse reactions (14–17).

Many patients with reactive airway disease, with or without a chronic obstructive component, have concomitant conditions such as hypertension or cardiac arrhythmias, which necessitate the use of β -blockers. However, review articles and practice guidelines usually list asthma and chronic obstructive pulmonary disease (COPD) as contraindications to β -blocker use, citing cases of acute bronchospasm during administration of noncardioselective β -blockers (6, 10, 18–22). Cardioselective β -blockers, or β_1 -blockers, have greater than 20 times more affinity for β_1 receptors than for β_2 receptors and in theory should pose much less risk for bronchoconstriction (23).

We used data from randomized, blinded, placebo-controlled trials to evaluate the effect of cardioselective β_1 -blockers on respiratory function in patients with reactive airway disease (defined as asthma or COPD with a reversible obstructive component). We also sought to evaluate the respiratory response to β_2 -agonists administered after β_1 -blockers or after placebo in the same participants. This

analysis has already been published as a review in the Cochrane Library (24).

METHODS

Patients

We chose to evaluate only patients with documented reactive airway disease because these patients are thought to be particularly susceptible to the adverse respiratory effects of β -blockers. Patients with COPD are generally at greater risk for ischemic heart disease than are patients with asthma and thus may benefit more from the use of β -blockers. This study evaluates a subgroup of patients with a documented chronic obstructive component of disease but was not designed to make recommendations about patients with COPD. A recent meta-analysis evaluated the use of cardioselective β -blockers in patients with COPD, given as a single dose or as continued treatment (25). Pooled data from 19 trials demonstrated no adverse effect on FEV₁ or respiratory symptoms for β_1 -blockers compared to placebo, even in patients with severe chronic airway obstruction.

Search Strategy

A search was performed to identify all relevant published clinical trials that addressed the effects of cardioselective β -blockers on airway function in patients with reactive airway disease. Two investigators jointly developed strategies with the help of an information service librarian

Context

Although β -blockers improve clinical outcomes in many patients with cardiovascular disease, clinicians sometimes avoid these agents in patients with concomitant lung disease because they fear precipitation of acute bronchospasm.

Contribution

This meta-analysis of 29 randomized trials shows that cardioselective β -blockers (β_1 -blockers), given for a few days to a few weeks, do not significantly worsen pulmonary function or respiratory symptoms and do not lead to increased use of inhalers in patients with mild to moderate reactive (reversible) airway disease.

Cautions

The studies in this meta-analysis were short, evaluated only cardioselective β -blockers, and did not include patients with severe or irreversible airway disease.

—The Editors

and the Cochrane Airways Group Trial Search Coordinator. The EMBASE, MEDLINE, and CINAHL databases were searched comprehensively to identify all relevant clinical trials in humans published between 1966 and May 2001. The search was performed by using the Cochrane Airways Group registry to identify randomized, blinded, placebo-controlled trials of reactive airways disease. Terms used in the search were *asthma**, *bronchial hyperreactivity**, *respiratory sounds**, *wheez**, *obstructive lung disease**, and *obstructive airway disease**. Trials of β -blockers were sought by using the terms *adrenergic antagonist**, *sympatholytic**, and *adrenergic receptor block**. Trials were not excluded on the basis of language. The search was further augmented by scanning references of identified articles, reviews, and abstracts at clinical symposia.

Study Selection

Two investigators independently evaluated studies for inclusion. In choosing articles, investigators were blinded to results but not to journal, author, or institution of studies. The observed interrater agreement for the assessment of inclusion was calculated as a percentage. For all clinical trials identified from the search, investigators determined whether the β -blocker used was cardioselective and whether it was considered to have intrinsic sympathomimetic activity (1, 26–36). Studies were evaluated if intravenous or oral cardioselective β -blockers were administered as a single dose or as continued treatment lasting 3 days or longer.

Single-dose trials were included if 1) FEV₁ at rest was reported, either as liters or as a percentage of the normal predicted value at baseline and at follow-up; 2) β_2 -agonists were withheld for at least 8 hours before initial FEV₁ measurement; 3) patients were not selected on the basis of

previous response to β -blockers; 4) the study was randomized, placebo-controlled, and single- or double-blinded; and 5) only patients with documented reactive airway disease were included. Reactive airway disease was demonstrated by a mean increase of at least 15% in FEV₁ in response to β_2 -agonist, response to methacholine challenge, or presence of asthma as defined by the American Thoracic Society (37). Crossover trials were included if different interventions were administered in random order.

We decided a priori that inclusion criteria 3, 4, and 5 would be applied to trials of continued treatment. Studies of continued treatment were included if they did not report FEV₁ but instead evaluated the amount of β_2 -agonist use and respiratory symptoms compared with placebo. Trials were also included if β_2 -agonists were not withheld during the trial.

Assessment of Validity

The methodologic quality of each trial was assessed according to the following factors: 1) Was the study randomized? If so, was the randomization procedure adequate, and was allocation concealed? 2) Were the patients and people administering the treatment blinded to the intervention? 3) Were withdrawals and dropouts described, and was the analysis performed on an intention-to-treat basis? On the basis of these criteria, studies were broadly subdivided as all quality criteria met (A), one or more quality criteria only partially met (B), or one or more criteria not met (C).

Clinical trials that did not meet criteria for inclusion but gave information on FEV₁ response to cardioselective β -blockers in patients with reactive airway disease were analyzed separately and used in a sensitivity analysis. These included studies that were not placebo-controlled; did not document asthma criteria; did not give baseline FEV₁ data; or, for single-dose studies, did not withhold β_2 -agonists for 8 hours before measurements.

Study Characteristics

The main intervention of interest was intravenous or oral cardioselective β -blockers versus placebo, given as a single dose or as continued treatment. Administration of a β_2 -agonist, intravenously or by inhalation, after the study medication or after placebo was also studied.

Each β_1 -blocker used was classified into one of two categories: β_1 -blockers without intrinsic sympathomimetic activity, and β_1 -blockers with intrinsic sympathomimetic activity.

Data Extraction

Two investigators independently extracted data on change in mean group FEV₁ in response to placebo or study drug; response of FEV₁ to β_2 -agonist administered after placebo or study drug; symptoms reported during the trial, such as wheezing, dyspnea, or exacerbation of asthma; and, for trials of continued treatment, weekly use of inhaled short-acting β_2 -agonists.

Data Synthesis

The ratio of the lowest group FEV₁ value after administration of study drug to baseline FEV₁ was measured for placebo and active treatment and was recorded as the percentage change from baseline. The placebo response was then subtracted from the treatment response to obtain the net treatment effect, reported as a percentage of the baseline FEV₁ value. For response to β_2 -agonists given after treatment or placebo, the new baseline value was the mean group FEV₁ value obtained after study drug but before β_2 -agonist administration. The net treatment effect was estimated by calculating the ratio of FEV₁ measured after agonist administration to the new baseline value for both placebo and active treatment and then subtracting the placebo-agonist response from the treatment-agonist response.

Whenever possible, the SD for the net treatment effect was calculated from individual-patient data or *P* values and was then used to derive the SDs for the analysis. Some trials provided SDs for treatment response and placebo response separately. For trials that reported no information on SDs, the average SD was obtained from trials that provided such data, calculated separately for placebo, treatment, and β -agonist responses. Sensitivity analyses were performed to evaluate the effect of including these trials by using the lowest and highest available SD in place of the pooled SD and also by excluding these trials from the analysis. The **Appendix Table** (available at www.annals.org) shows the method used to obtain SDs for each trial.

The mean treatment effects were pooled to obtain a weighted average of the study means using the fixed-effects model for continuous outcomes (38, 39). Confidence intervals with 95% significance were obtained for the pooled study means. The analysis was performed by using Meta View 4.1 (Cochrane Library software [Update Software, Oxford, United Kingdom]).

Results for respiratory symptoms were measured as a risk difference by subtracting the percentage of patients with respiratory events during treatment from the percentage of patients with respiratory events during placebo use. The risk differences were then pooled by using the fixed-effects model for dichotomous outcomes. The results for inhaler use during continued treatment trials were measured as the incidence of use during placebo minus the incidence of use during treatment. The weighted mean treatment effects were pooled by using the fixed-effects model for continuous outcomes.

To test for interstudy heterogeneity, the chi-square value was calculated for the assumption of homogeneity. In addition, the confidence intervals from the fixed-effects model were compared with those from the random-effects model (40). The fixed-effects model was chosen to report the results because minimal heterogeneity was seen in most of the analyses. When heterogeneity was noted, the results from both the random-effects model and fixed-effects model were reported.

A subgroup analysis was performed to compare the treatment effects of cardioselective β -blockers with and without intrinsic sympathomimetic activity. Another analysis was done to evaluate the response of patients with concomitant chronic airways obstruction, defined as a baseline FEV₁ of less than 80% of the normal predicted value or less than 1.8 L, or defined by using American Thoracic Society criteria (41). A third subgroup analysis evaluated the treatment response in participants known to have comorbid cardiovascular conditions, such as hypertension.

DATA SYNTHESIS

Search Results

The **Appendix Figure** (available at www.annals.org) shows the results of the search for articles. The database search identified 200 potentially relevant articles. After review of articles and bibliographies, 104 trials of β -blockers in patients with reactive airway disease were found. Of these trials, 29 met inclusion criteria: Nineteen gave information on single-dose studies (42–60) and 10 provided data on continued treatment of longer duration (28, 51, 61–68). One of the articles (51) gave data for both types of administration. Interrater agreement for study eligibility was 94%. Consensus was reached on the remaining trials. The **Appendix Table** (available at www.annals.org) shows the characteristics of the included studies.

Clinical trials that did not meet inclusion criteria but gave data on FEV₁ responses to cardioselective β -blockers in patients with reactive airway disease were analyzed separately and used in a sensitivity analysis (69–78).


Methodologic Quality of Included Studies

All studies were small crossover trials that received a quality score of B because the randomization process was not described in detail or the trial was single-blind instead of double-blind. Many of the trials were performed 20 or 30 years ago and did not provide adequate information with which to calculate SDs for the net treatment effect. Sensitivity analyses were performed to evaluate the effect of including trials that provided no information on SDs.

Quantitative Data Synthesis

Single-Dose Treatment

Nineteen studies of single-dose treatment included 240 patients, 79% of whom were men. Each study included an average of 12.6 patients, and the total dropout rate was 2.0%. From the available information, the age range of participants was 19.5 to 65.1 years (mean, 40.1 years). These baseline characteristics were the same for the placebo and treatment groups. The baseline FEV₁ was 2.41 ± 0.15 L in the treatment group and 2.42 ± 0.2 L in the placebo group. Cardioselective β -blockers without intrinsic sympathomimetic activity that were included in the study were atenolol, metoprolol, bisoprolol, and practolol. Cardioselective β -blockers with intrinsic sympathomimetic

Figure 1. Effects of treatment on FEV₁ for single-dose studies.

Diamonds represent the extent of the confidence intervals. For group 1, test for heterogeneity, $P > 0.2$; test for overall effect, $P < 0.001$. For group 2, test for heterogeneity, $P = 0.05$; test for overall effect, $P = 0.15$. For both groups, test for heterogeneity, $P = 0.09$; test for overall effect, $P < 0.001$.

activity that were studied were celiprolol, acebutolol, and xamoterol.

Compared with placebo, single doses of cardioselective β -blockers as a group were associated with a 7.46% (CI, 5.59% to 9.32%) reduction in FEV₁ but with a 4.63% (CI, 2.47% to 6.78%) increase in FEV₁ after β -agonist was given (Figures 1 and 2). The number of patients with respiratory symptoms did not increase significantly in any of the 19 studies (0.01% [CI, -0.02% to 0.03%]).

Continued Treatment

Data from 10 studies involving 141 participants (77% of whom were men) were evaluated for response to contin-

ued treatment ranging from 3 days to 4 weeks. Each study contained an average of 15.4 patients, and the dropout rate was 1.3%. From the available information, the average age of the participants was 51.3 years. The average baseline FEV₁ was 1.81 ± 0.13 L for the treatment group and 1.81 ± 0.15 L for the placebo group. Five of the studies (54 participants) did not provide data on FEV₁; these studies were included in analyses of symptoms and inhaler use.

In the continued treatment trials, cardioselective β -blockers as a group did not significantly differ from placebo in terms of FEV₁ response (-0.42% [CI, -3.74% to 2.91%]), number of patients with symptoms (0.01% [CI,

−0.02% to 0.04%)), or incidence of inhaler use (−0.11% [CI, −6.75% to 6.54%]). Cardioselective β -blockers produced an 8.74% increase (CI, 1.96% to 15.52%) in FEV₁ compared with placebo after β -agonist was given (Figures 3 and 4).

Interstudy Variance


No significant interstudy variance was found in FEV₁ treatment effect, symptoms, and long-term use of an inhaler. Heterogeneity was detected between studies in FEV₁ after β_2 -agonist use in both the single-dose and continued treatment studies. This heterogeneity was noted only for β -blockers without intrinsic sympathomimetic activity. When the random-effects model was compared with the fixed-effects model for the β_2 -agonist response in patients who received β -blockers without intrinsic sympathomi-

metic activity, a difference of less than 1 percentage point was found for single-dose studies (5.66% [CI, 1.81% to 9.51%] vs. 6.59% [CI, 4.18% to 9.01%]) and a difference of 1.7 percentage points was found for continued treatment studies (10.32% [CI, −6.38% to 27.01%] vs. 12.0% [CI, 4.12% to 19.89%]).


Subgroup Analysis

For single-dose trials, β_1 -blockers without intrinsic sympathomimetic activity were associated with a 6.5% reduction in FEV₁ (CI, 2.2% to 10.7%) compared to those with sympathomimetic activity. However, treatment with β_1 -blockers without intrinsic sympathomimetic activity was associated with a 9.7% increase in FEV₁ in response to β -agonist (CI, 5.6 to 13.7%) compared to those with sympathomimetic activity. In the continued treatment trials,

Figure 2. Effects of treatment after use of β_2 -agonists on FEV₁ for single-dose studies.



Diamonds represent the extent of the confidence intervals. For group 1, test for heterogeneity, $P < 0.001$; test for overall effect, $P < 0.001$. For group 2, test for heterogeneity, $P > 0.2$; test for overall effect, $P = 0.2$. For both groups, test for heterogeneity, $P < 0.001$; test for overall effect, $P < 0.001$.

Figure 3. Effects of treatment on FEV₁ for continued treatment studies.

Diamonds represent the extent of the confidence intervals. For group 1, test for heterogeneity, $P > 0.2$; test for overall effect, $P = 0.17$. For group 2, test for overall effect, $P > 0.2$. For both groups, test for heterogeneity, $P = 0.18$; test for overall effect, $P > 0.2$.

no significant difference in FEV₁ response was observed for β_1 -blockers without intrinsic sympathomimetic activity compared to those with sympathomimetic activity (5.94% [CI, -0.73% to 12.61%]). However, β -blockers without intrinsic sympathomimetic activity produced a 12.6% increase in FEV₁ (CI, 0.3% to 25.6%) after β -agonist administration compared to β -blockers with sympathomimetic activity.

To evaluate the treatment effect in patients with concomitant COPD, 10 trials that included only patients with documented chronic airway obstruction were analyzed separately (46, 52, 53, 59, 61, 63–66, 68). No significant difference in FEV₁ treatment effect was observed in single-dose trials (-5.28% [CI, -10.03% to -0.54%]) or continued treatment (1.07% [CI, -3.3% to 5.44%]), and no increase in symptoms occurred in any of the trials.

In eight of the trials, all participants had a comorbid condition, such as hypertension (46, 51, 52, 61, 63, 65, 67, 68). When only these trials were included in the analysis, the treatment effect for FEV₁ did not change significantly in single-dose trials (-6.83% [CI, -11.46% to -2.20%]) or with continued treatment (1.31% [CI, -2.62% to 5.24%]).

Sensitivity Analysis

A sensitivity analysis was performed to evaluate the effect of including studies that did not provide SDs (46, 55, 56, 60). When these trials were excluded from the analysis, the difference in all variables measured was less

than 0.5 percentage point. When the analysis was performed by replacing the pooled SD with the lowest and highest available SD, the difference in results between the highest and lowest SD was 2 percentage points or less.

A sensitivity analysis was also performed to evaluate the effect of excluding trials that did not meet the inclusion criteria set by the study but that provided information on FEV₁ and symptoms for cardioselective β -blocker use in patients with reactive airway disease (69–78). Data analysis of 10 excluded studies with 141 participants showed no significant difference in any variables compared with studies that met inclusion criteria.

DISCUSSION

Our results indicate that the first dose of a cardioselective β -blocker produces a small decrease in FEV₁ that is not associated with adverse respiratory effects compared to placebo. After continued treatment for a few days to weeks, FEV₁, symptoms, and inhaler use did not differ. Cardioselective β -blockers, given as a single dose or as continued treatment, were associated with an increase in response to β_2 -agonists compared with placebo. Of the 80 trials on cardioselective β -blockers that we identified, none demonstrated an increase in respiratory symptoms for β_1 -blockers compared with placebo or baseline values.

Subgroup analyses were performed to evaluate the effect of cardioselective β -blockers on patients with concom-

itant COPD or cardiovascular diseases, such as hypertension, because these patients are most often targeted for β -blocker treatment. No significant difference in the FEV₁ treatment effect or incidence of symptoms or inhaler use was observed.

Our meta-analysis has several limitations. Most of the participants were relatively young and had mild to moderate airway obstruction; persons with recent exacerbation of asthma were often excluded from study. In addition, because many of the studies were of short duration, we cannot comment on the effect of cardioselective β -blockers on the frequency or severity of acute asthma exacerbations after several months of treatment. Furthermore, this analysis was based only on published literature and therefore is subject to publication bias. However, funnel plots of effect size versus standard error for the trials in this analysis showed no evidence of bias. We believe that these pooled results provide valuable information on the safety of cardioselective β -blockers in patients with reactive airway disease, with or without concomitant COPD or cardiovascular disease.

The current standard of care is to consider reactive airway disease to be a contraindication to the use of all β -blockers (6, 10, 18–22). Because of the proven mortality benefit of β -blockers, many of the other relative or absolute contraindications traditionally listed for β -blockers, including impaired left ventricular function, peripheral vascular disease, diabetes mellitus, depression, and advanced age (7, 14, 79–87), have been questioned and disproved.


The original evidence of a potential adverse effect of

β -blockers in reactive airway disease was based on case reports of acute bronchospasm precipitated by high doses of noncardioselective blockers, presumably due to their blockade of β_2 receptors on bronchial smooth muscle (88–91). Pooled results of 16 trials that evaluated noncardioselective β -blockers showed that regular use of nonselective β -blockers compared with placebo caused a 13.5% decrease (CI, –23.0% to –4.0%) in FEV₁ and a 22.5% decrease (CI, –32.5% to –12.5%) in the FEV₁ response after β_2 -agonists were given (24, 32, 34, 43, 48, 52, 55–57, 59, 61, 68, 92–96). No significant increase in symptoms or inhaler use was found. However, the decrease in β -agonist response seen with nonselective β -blockers may increase the risk for a clinically significant adverse effect during an exacerbation of asthma.

Cardioselective β -blockers, such as atenolol, bisoprolol, and metoprolol, are at least 20 times more effective at blocking β_1 -receptors than β_2 -receptors; thus, at therapeutic doses, their β_2 -blocking effect is negligible (23). The doses of β_1 -blockers that we evaluated ranged from therapeutic to mildly suprathreshold. For example, single-dose studies using atenolol or metoprolol in doses ranging from 50 to 200 mg showed no clinically apparent effect on respiratory function. Linear regression analysis could not differentiate a treatment effect between low and high doses because there were few low-dose trials and no trials used doses high enough to diminish cardioselectivity.

Our results indicate that for cardioselective β -blockers without intrinsic sympathomimetic activity, the minimal decrease in FEV₁ noted with a single dose is attenuated over a few days to weeks. In addition, FEV₁ increases in

Figure 4. Effects of treatment after use of β_2 -agonists on FEV₁ for continued treatment studies.



Diamonds represent the extent of the confidence intervals. For group 1, test for heterogeneity, $P = 0.013$; test for overall effect, $P = 0.003$. For group 2, test for overall effect, $P > 0.2$. For both groups, test for heterogeneity, $P < 0.01$; test for overall effect, $P = 0.01$.

response to β_2 -agonist administration compared with placebo, and this increase is maintained with continued treatment. These results could be explained by upregulation or sensitization of β_2 receptors that is accompanied by an increased effect of endogenous or exogenous β_2 -agonist stimulation (97–99). Accumulating evidence indicates that continued use of inhaled β_2 -agonists in patients with reactive airway disease is associated with a tolerance to β_2 -agonist stimulation and an increase in asthma attacks (100–105). There is also evidence that treatment with β -blockers that have intrinsic β_2 sympathomimetic activity is associated with downregulation of β_2 receptors (106–109). This finding is consistent with data from our analysis showing that β -blockers with intrinsic sympathomimetic activity did not produce the increase in β_2 -agonist response that was seen with β -blockers without intrinsic sympathomimetic activity.

Only a small proportion of patients with heart disease who would benefit from β -blockers currently receive this treatment, mainly owing to unfounded fears about their adverse effects (110–113). A study of survivors of myocardial infarction included 46 000 patients with asthma and chronic obstructive lung disease and showed a significant reduction in total mortality rate among those treated with β -blockers compared with those who were not (14). Other studies of the use of β -blockers in patients with cardiac disease and concomitant chronic obstructive lung disease or asthma found that these medicines were well tolerated (114–116). Other trials evaluating the use of β -blockers in hypertensive patients, many of whom had reactive airway disease, did not demonstrate worsening of respiratory symptoms or FEV₁ in these patients (32, 92, 117, 118). A recent study showed that COPD and asthma were the comorbid conditions most commonly associated with β -blockers' being withheld in elderly patients after a myocardial infarction (119).

Patients with COPD are thought to be at greater risk than those with reactive airway disease for developing ischemic heart disease and other cardiovascular conditions requiring the use of β -blockers. However, the presenting features of COPD and reactive airway disease overlap substantially.

Another recent meta-analysis evaluated the effect of cardioselective β -blockers in patients with COPD and found no change in FEV₁ or respiratory symptoms for single doses or continued use of these agents compared with placebo (25). Subgroup analyses revealed no difference in results for patients with concomitant reactive airway disease and those with severe chronic airways obstruction, as demonstrated by a baseline FEV₁ less than 1.4 L or less than 50% the normal predicted value. Three of the trials from that meta-analysis are also included in our analysis (52, 61, 66). The cumulative evidence from these two meta-analyses indicates that cardioselective β -blockers should not be withheld in patients with reactive airway disease or COPD.

From Stanford University School of Medicine, Palo Alto, California; Santa Clara Valley Medical Center, San Jose, California; and Cornell University, Ithaca, New York.

Acknowledgments: The authors thank Steve Milan for guidance with the Cochrane review, Toby Lasserson for technical assistance, and Karen Blackhall for coordinating the trials search.

Requests for Single Reprints: Shelley Salpeter, MD, Department of Medicine, Santa Clara Valley Medical Center, 751 South Bascom Avenue, San Jose, CA 95128; e-mail, shelly.salpeter@hhs.co.santa-clara.ca.us.

Current author addresses and author contributions are available at www.annals.org.

References

1. Wadworth AN, Murdoch D, Brogden RN. Atenolol. A reappraisal of its pharmacological properties and therapeutic use in cardiovascular disorders. *Drugs*. 1991;42:468–510. [PMID: 1720383]
2. Heidenreich PA, McDonald KM, Hastie T, Fadel B, Hagan V, Lee BK, et al. Meta-analysis of trials comparing beta-blockers, calcium antagonists, and nitrates for stable angina. *JAMA*. 1999;281:1927–36. [PMID: 10349897]
3. Frishman WH, Furberg CD, Friedewald WT. Beta-adrenergic blockade for survivors of acute myocardial infarction. *N Engl J Med*. 1984;310:830–7. [PMID: 6142420]
4. Freemantle N, Cleland J, Young P, Mason J, Harrison J. Beta blockade after myocardial infarction: systematic review and meta regression analysis. *BMJ*. 1999;318:1730–7. [PMID: 10381708]
5. Cardiovascular risk and risk factors in a randomized trial of treatment based on the beta-blocker oxprenolol: the International Prospective Primary Prevention Study in Hypertension (IPPPSH). The IPPPSH Collaborative Group. *J Hypertens*. 1985;3:379–92. [PMID: 2864374]
6. The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med*. 1997;157:2413–46. [PMID: 9385294]
7. Lechat P, Packer M, Chalon S, Cucherat M, Arab T, Boissel JP. Clinical effects of beta-adrenergic blockade in chronic heart failure: a meta-analysis of double-blind, placebo-controlled, randomized trials. *Circulation*. 1998;98:1184–91. [PMID: 9743509]
8. Doughty RN, Rodgers A, Sharpe N, MacMahon S. Effects of beta-blocker therapy on mortality in patients with heart failure. A systematic overview of randomized controlled trials. *Eur Heart J*. 1997;18:560–5. [PMID: 9129883]
9. Steinbeck G, Andresen D, Bach P, Haberl R, Oeff M, Hoffmann E, et al. A comparison of electrophysiologically guided antiarrhythmic drug therapy with beta-blocker therapy in patients with symptomatic, sustained ventricular tachyarrhythmias. *N Engl J Med*. 1992;327:987–92. [PMID: 1355595]
10. Kendall MJ. Clinical relevance of pharmacokinetic differences between beta blockers. *Am J Cardiol*. 1997;80:15J–19J. [PMID: 9375944]
11. Klein I, Becker DV, Levey GS. Treatment of hyperthyroid disease. *Ann Intern Med*. 1994;121:281–8. [PMID: 7518659]
12. Jones MK, John R, Jones GR. The effect of oxprenolol, acebutolol and propranolol on thyroid hormones in hyperthyroid subjects. *Clin Endocrinol (Oxf)*. 1980;13:343–7. [PMID: 7438476]
13. Mangano DT, Layug EL, Wallace A, Tateo I. Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. Multicenter Study of Perioperative Ischemia Research Group. *N Engl J Med*. 1996;335:1713–20. [PMID: 8929262]
14. Gottlieb SS, McCarter RJ, Vogel RA. Effect of beta-blockade on mortality among high-risk and low-risk patients after myocardial infarction. *N Engl J Med*. 1998;339:489–97. [PMID: 9709041]
15. Kennedy HL, Rosenson RS. Physician use of beta-adrenergic blocking therapy: a changing perspective [Editorial]. *J Am Coll Cardiol*. 1995;26:547–52. [PMID: 7608462]

16. Viskin S, Barron HV. Beta blockers prevent cardiac death following a myocardial infarction: so why are so many infarct survivors discharged without beta blockers? [Editorial] *Am J Cardiol*. 1996;78:821-2. [PMID: 8857489]
17. Chafin CC, Soberman JE, Demirkan K, Self T. Beta-blockers after myocardial infarction: do benefits ever outweigh risks in asthma? *Cardiology*. 1999;92:99-105. [PMID: 10702651]
18. Craig T, Richerson HB, Moeckli J. Problem drugs for the patient with asthma. *Compr Ther*. 1996;22:339-44. [PMID: 8814381]
19. Belli G, Topol EJ. Adjunctive pharmacologic strategies for acute MI. *Contemp Intern Med*. 1995;7:51-9. [PMID: 10150611]
20. O'Malley K, Cox JP, O'Brien E. Choice of drug treatment for elderly hypertensive patients. *Am J Med*. 1991;90:27S-33S. [PMID: 2006657]
21. Tattersfield AE. Beta adrenoceptor antagonists and respiratory disease. *J Cardiovasc Pharmacol*. 1986;8 Suppl 4:S35-9. [PMID: 2427850]
22. Tattersfield AE. Respiratory function in the elderly and the effects of beta blockade. *Cardiovasc Drugs Ther*. 1991;4 Suppl 6:1229-32. [PMID: 1672601]
23. Wellstein A, Palm D, Belz GG, Butzer R, Polsak R, Pett B. Reduction of exercise tachycardia in man after propranolol, atenolol and bisoprolol in comparison to beta-adrenoceptor occupancy. *Eur Heart J*. 1987;8 Suppl M:3-8. [PMID: 2897299]
24. Salpeter S, Ormiston T, Salpeter E. Cardioselective beta-blocker use in patients with reversible airway disease (Cochrane Review). The Cochrane Library. Oxford: Update Software; 2001:2.
25. Salpeter S, Ormiston T, Salpeter E, Poole P, Cates C. Cardioselective beta-blocker use in patients with COPD. (Cochrane Review). The Cochrane Library. Oxford: Update Software; 2002:2.
26. Lois M, Honig E. Beta-blockade post-MI: safe for patients with asthma or COPD? *J Respir Dis*. 1997;18:568-91.
27. Fitzgerald JD. The applied pharmacology of beta-adrenoceptor antagonists (beta blockers) in relation to clinical outcomes. *Cardiovasc Drugs Ther*. 1991;5:561-76. [PMID: 1678960]
28. Nicolaescu V, Manicadide M, Stroescu V. β -Adrenergic blockade with practolol in acetylcholine-sensitive asthma patients. *Respiration*. 1972;29:139-54. [PMID: 4401715]
29. Palmer KN, Hamilton WF, Lge JS, Diamant ML. Effect of a selective beta-adrenergic blocker in preventing falls in arterial oxygen tension following isoprenaline in asthmatic subjects. *Lancet*. 1969;2:1092-4. [PMID: 4186991]
30. Löfdahl CG, Marlin GE, Svedmyr N. Pafenolol, a highly selective beta 1-adrenoceptor-antagonist, in asthmatic patients: interaction with terbutaline. *Clin Pharmacol Ther*. 1983;33:1-9. [PMID: 6129091]
31. Decalmer PB, Chatterjee SS, Cruickshank JM, Benson MK, Sterling GM. Beta-blockers and asthma. *Br Heart J*. 1978;40:184-9. [PMID: 25075]
32. Falliers CJ, Vrchota J, Blasucci DJ, Maloy JW, Medakovic M. The effects of treatments with labetalol and hydrochlorothiazide on ventilatory function of asthmatic hypertensive patients with demonstrated bronchiosensitivity to propranolol. *J Clin Hypertens*. 1985;1:70-9. [PMID: 3836297]
33. Singh BN, Whitlock RM, Comber RH, Williams FH, Harris EA. Effects of cardioselective beta adrenoceptor blockade on specific airways resistance in normal subjects and in patients with bronchial asthma. *Clin Pharmacol Ther*. 1976;19(5 Pt 1):493-501. [PMID: 6179]
34. Skinner C, Gaddie J, Palmer KN. Comparison of intravenous AH 5158 (ibidomide) and propranolol in asthma. *Br Med J*. 1975;2:59-61. [PMID: 236806]
35. Lois M, Roman J, Honig E. Using beta-blockers for CHF in patients with COPD. *J Respir Dis*. 1999;20:31-43.
36. Formgren H. Broncho- and cardioselective beta-receptor active drugs in the treatment of asthmatic patients. Clinical studies of effects and side effects of terbutaline, practolol and metoprolol. *Scand J Respir Dis Suppl*. 1977;97:1-37. [PMID: 266273]
37. Guidelines for the evaluation of impairment/disability in patients with asthma. American Thoracic Society. Medical Section of the American Lung Association. *Am Rev Respir Dis*. 1993;147:1056-61. [PMID: 8466106]
38. Mantel N, Haenszel WH. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst*. 1959;22:719-48.
39. Yusuf S, Peto R, Lewis J, Collins R, Sleight P. Beta blockade during and after myocardial infarction: an overview of the randomized trials. *Prog Cardiovasc Dis*. 1985;27:335-71. [PMID: 2858114]
40. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7:177-88. [PMID: 3802833]
41. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. American Thoracic Society. *Am J Respir Crit Care Med*. 1995;152:S77-121. [PMID: 7582322]
42. Tantucci C, Bruni B, Dottorini ML, Peccini F, Motolese M, Lecaillon JB, et al. Comparative evaluation of cardioselectivity of metoprolol OROS and atenolol: a double-blind, placebo-controlled crossover study. *Am Heart J*. 1990;120:467-72. [PMID: 2200257]
43. Chodosh S, Tuck J, Blasucci DJ. The effects of dilevalol, metoprolol, and placebo on ventilatory function in asthmatics. *J Cardiovasc Pharmacol*. 1988;11 Suppl 2:S18-24. [PMID: 2464094]
44. Lammers JW, Müller ME, Folgering HT, van Herwaarden CL. A comparative study on the ventilatory and haemodynamic effects of xamoterol and atenolol in asthmatic patients. *Br J Clin Pharmacol*. 1986;22:595-602. [PMID: 2878680]
45. Falliers CJ, Vincent ME, Medakovic M. Effect of single doses of labetalol, metoprolol, and placebo on ventilatory function in patients with bronchial asthma: interaction with isoproterenol. *J Asthma*. 1986;23:251-60. [PMID: 3771472]
46. Chatterjee SS. The cardioselective and hypotensive effects of bisoprolol in hypertensive asthmatics. *J Cardiovasc Pharmacol*. 1986;8 Suppl 11:S74-7. [PMID: 2439803]
47. Doshan HD, Brown R, Applin WJ, Kapoor M, Caruso FS. Effects of high doses of celiprolol in asthmatic patients. *J Cardiovasc Pharmacol*. 1986;8 Suppl 4:S109-11. [PMID: 2427837]
48. Doshan HD, Rosenthal RR, Brown R, Slutsky A, Applin WJ, Caruso FS. Celiprolol, atenolol and propranolol: a comparison of pulmonary effects in asthmatic patients. *J Cardiovasc Pharmacol*. 1986;8 Suppl 4:S105-8. [PMID: 2427836]
49. Lammers JW, Folgering HT, van Herwaarden CL. Ventilatory effects of beta 1-receptor-selective blockade with bisoprolol and metoprolol in asthmatic patients. *Eur J Clin Pharmacol*. 1984;27:141-5. [PMID: 6149939]
50. Greefhorst AP, van Herwaarden CL. Ventilatory and haemodynamic effects of beta 1-selective and beta 2-selective stimulation in asthmatic patients. *Eur J Respir Dis Suppl*. 1984;135:147-50. [PMID: 6329801]
51. Lawrence DS, Sahay JN, Chatterjee SS, Cruickshank JM. Asthma and beta-blockers. *Eur J Clin Pharmacol*. 1982;22:501-9. [PMID: 6127219]
52. Adam WR, Meagher EJ, Barter CE. Labetalol, beta blockers, and acute deterioration of chronic airway obstruction. *Clin Exp Hypertens A*. 1982;4:1419-28. [PMID: 6126285]
53. Löfdahl CG, Svedmyr N. Cardioselectivity of atenolol and metoprolol. A study in asthmatic patients. *Eur J Respir Dis*. 1981;62:396-404. [PMID: 6751838]
54. van den Bergh JH, van Herwaarden CL. Ventilatory effects of ordinary and slow-release tablets of metoprolol in asthmatic patients. *Eur J Respir Dis*. 1981;62:168-72. [PMID: 7308331]
55. Ellis ME, Sahay JN, Chatterjee SS, Cruickshank JM, Ellis SH. Cardioselectivity of atenolol in asthmatic patients. *Eur J Clin Pharmacol*. 1981;21:173-6. [PMID: 7032947]
56. Ruffin RE, Frith PA, Anderton RC, Kumana CR, Newhouse MT, Hargreave FE. Selectivity of beta adrenoreceptor antagonist drugs assessed by histamine bronchial provocation. *Clin Pharmacol Ther*. 1979;25:536-40. [PMID: 35296]
57. Benson MK, Berrill WT, Cruickshank JM, Sterling GS. A comparison of four beta-adrenoceptor antagonists in patients with asthma. *Br J Clin Pharmacol*. 1978;5:415-9. [PMID: 26371]
58. Skinner C, Palmer KN. Comparison of the effects of acebutolol (Sectral) and practolol (Eraldin) on airways obstruction in asthmatics. *Br J Clin Pharmacol*. 1975;2:417-22. [PMID: 786353]
59. Johnsson G, Svedmyr N, Thiringer G. Effects of intravenous propranolol and metoprolol and their interaction with isoprenaline on pulmonary function, heart rate and blood pressure in asthmatics. *Eur J Clin Pharmacol*. 1975;8:175-80. [PMID: 1233216]
60. Lammers JW, Müller ME, Folgering HT, van Herwaarden CL. Effects of terbutaline and atenolol on large and small airways in asthmatic patients. *Eur Respir J*. 1988;1:453-7. [PMID: 3169216]

61. Fogari R, Zoppi A, Tettamanti F, Poletti L, Rizzardi G, Fiocchi G. Comparative effects of celiprolol, propranolol, oxprenolol, and atenolol on respiratory function in hypertensive patients with chronic obstructive lung disease. *Cardiovasc Drugs Ther*. 1990;4:1145-9. [PMID: 1982059]
62. Nicolaescu V, Racoveanu C, Manicattide M. Effects of exercise on practolol-treated asthmatic patients. *Eur J Clin Pharmacol*. 1973;6:3-8. [PMID: 4587515]
63. Dorow P, Clauzel AM, Capone P, Mayol R, Mathieu M. A comparison of celiprolol and chlorthalidone in hypertensive patients with reversible bronchial obstruction. *J Cardiovasc Pharmacol*. 1986;8 Suppl 4:S102-4. [PMID: 2427835]
64. Butland RJ, Pang JA, Geddes DM. Effect of beta-adrenergic blockade on hyperventilation and exercise tolerance in emphysema. *J Appl Physiol*. 1983;54:1368-73. [PMID: 6863097]
65. Bauer K, Kaik G, Kaik B. Osmotic release oral drug delivery system of metoprolol in hypertensive asthmatic patients. Pharmacodynamic effects on beta 2-adrenergic receptors. *Hypertension*. 1994;24:339-46. [PMID: 7916004]
66. Fenster PE, Hasan FM, Abraham T, Woolfenden J. Effect of metoprolol on cardiac and pulmonary function in chronic obstructive pulmonary disease. *Clin Cardiol*. 1983;6:125-9. [PMID: 6851273]
67. van Zyl AI, Jennings AA, Bateman ED, Opie LH. Comparison of respiratory effects of two cardioselective beta-blockers, celiprolol and atenolol, in asthmatics with mild to moderate hypertension. *Chest*. 1989;95:209-13. [PMID: 2562825]
68. Lammers JW, Folgering HT, van Herwaarden CL. Ventilatory effects of long-term treatment with pindolol and metoprolol in hypertensive patients with chronic obstructive lung disease. *Br J Clin Pharmacol*. 1985;20:205-10. [PMID: 4041341]
69. Abraham TA, Hasan FM, Fenster PE, Marcus FI. Effect of intravenous metoprolol on reversible obstructive airways disease. *Clin Pharmacol Ther*. 1981;29:582-7. [PMID: 7214788]
70. Boye NP, Vale JR. Effect in bronchial asthma of a new beta-adrenergic blocking drug atenolol (ICI 66, 082). *Eur J Clin Pharmacol*. 1977;11:11-4. [PMID: 319006]
71. Schindl R, Würtz J, Hoffmann H. The effect of the cardioselective beta blocker celiprolol on pulmonary function in asthmatic patients. *J Cardiovasc Pharmacol*. 1986;8 Suppl 4:S99-101. [PMID: 2427862]
72. Larsson K. Influence of labetalol, propranolol and practolol in patients with asthma. *Eur J Respir Dis*. 1982;63:221-30. [PMID: 6124442]
73. Pujet JC, Dubreuil C, Fleury B, Provendier O, Abella ML. Effects of celiprolol, a cardioselective beta-blocker, on respiratory function in asthmatic patients. *Eur Respir J*. 1992;5:196-200. [PMID: 1348481]
74. Vilsvik JS, Schaanning J. Effect of atenolol on ventilatory and cardiac function in asthma. *Br Med J*. 1976;2:453-5. [PMID: 8188]
75. Beumer HM, Teirlinck C, Wiseman RA. Comparative investigation of the respiratory and cardiovascular effect of mepindolol, propranolol and pindolol in asthmatic patients. *Int J Clin Pharmacol Biopharm*. 1978;16:249-53. [PMID: 27463]
76. Mue S, Sasaki T, Shibahara S, Takahashi M, Ohmi T, Yamauchi K, et al. Influence of metoprolol on hemodynamics and respiratory function in asthmatic patients. *Int J Clin Pharmacol Biopharm*. 1979;17:346-50. [PMID: 489201]
77. Blaive B, Nahon F, Lemoigne F, Lewest G. [Long-term beta adrenergic antagonists in spastic bronchopathies]. *Allerg Immunol (Paris)*. 1988;20:161-2, 164-5. [PMID: 2900013]
78. Cazzola M, Noschese P, D'Amato M, D'Amato G. Comparison of the effects of single oral doses of nebivolol and celiprolol on airways in patients with mild asthma. *Chest*. 2000;118:1322-6. [PMID: 11083681]
79. Radack K, Deck C. Beta-adrenergic blocker therapy does not worsen intermittent claudication in subjects with peripheral arterial disease. A meta-analysis of randomized controlled trials. *Arch Intern Med*. 1991;151:1769-76. [PMID: 1679624]
80. Rosenson RS. The truth about beta-blocker adverse effects—depression, claudication, and lipids. *Journal of Ambulatory Monitoring*. 1993;6:163-71.
81. Jonas M, Reicher-Reiss H, Boyko V, Shotan A, Mandelzweig L, Goldbourt U, et al. Usefulness of beta-blocker therapy in patients with non-insulin-dependent diabetes mellitus and coronary artery disease. Bezafibrate Infarction Prevention (BIP) Study Group. *Am J Cardiol*. 1996;77:1273-7. [PMID: 8677865]
82. Kjekshus J, Gilpin E, Cali G, Blackey AR, Henning H, Ross J Jr. Diabetic patients and beta-blockers after acute myocardial infarction. *Eur Heart J*. 1990;11:43-50. [PMID: 1968386]
83. Wicklmayr M, Rett K, Dietze G, Mehnert H. Effects of beta-blocking agents on insulin secretion and glucose disposal. *Horm Metab Res Suppl*. 1990;22:29-33. [PMID: 1975244]
84. Bright RA, Everitt DE. Beta-blockers and depression. Evidence against an association. *JAMA*. 1992;267:1783-7. [PMID: 1347572]
85. Beto JA, Bansal VK. Quality of life in treatment of hypertension. A meta-analysis of clinical trials. *Am J Hypertens*. 1992;5:125-33. [PMID: 1349484]
86. Krumholz HM, Radford MJ, Wang Y, Chen J, Marciniak TA. Early beta-blocker therapy for acute myocardial infarction in elderly patients. *Ann Intern Med*. 1999;131:648-54. [PMID: 10577326]
87. Opie LH. Required beta blocker profile in the elderly. *Cardiovasc Drugs Ther*. 1991;4 Suppl 6:1273-80. [PMID: 1672603]
88. McNeill RS. Effect of a beta-adrenergic-blocking agent, propranolol, on asthmatics. *Lancet*. 1964;2:1101-2.
89. Zaid G, Beall GN. Bronchial response to beta-adrenergic blockade. *N Engl J Med*. 1966;275:580-4. [PMID: 5920412]
90. Anderson EG, Calcraft B, Jariwalla AG, Al-Zaibak M. Persistent asthma after treatment with beta-blocking agents. *Br J Dis Chest*. 1979;73:407-8. [PMID: 45308]
91. Raine JM, Palazzo MG, Kerr JH, Sleight P. Near-fatal bronchospasm after oral nadolol in a young asthmatic and response to ventilation with halothane. *Br Med J (Clin Res Ed)*. 1981;282:548-9. [PMID: 6109556]
92. George RB, Manocha K, Burford JG, Conrad SA, Kinasevitz GT. Effects of labetalol in hypertensive patients with chronic obstructive pulmonary disease. *Chest*. 1983;83:457-60. [PMID: 6337786]
93. Light RW, Chetty KG, Stansbury DW. Comparison of the effects of labetalol and hydrochlorothiazide on the ventilatory function of hypertensive patients with mild chronic obstructive pulmonary disease. *Am J Med*. 1983;75:109-14. [PMID: 6356897]
94. Giulekas D, Georgopoulos D, Papakosta D, Antoniadou H, Sotiropoulou E, Vamvalis C. Influence of pindolol on asthmatics and effect of bronchodilators. *Respiration*. 1986;50:158-66. [PMID: 2948256]
95. Devereux G, Fishwick K, Aiken TC, Bourke SJ, Hendrick DJ. Adverse effects of a single dose of (+)-sotalol in patients with mild stable asthma. *Br J Clin Pharmacol*. 1998;46:79-82. [PMID: 9690953]
96. Sue DY, Hansen JE, Wasserman K. Beta-adrenergic blockade with pindolol (LB-46) in mild to moderate asthma. *Chest*. 1981;80:537-42. [PMID: 7297143]
97. Motomura S, Deighton NM, Zerkowski HR, Doetsch N, Michel MC, Brodde OE. Chronic beta 1-adrenoceptor antagonist treatment sensitizes beta 2-adrenoceptors, but desensitizes M2-muscarinic receptors in the human right atrium. *Br J Pharmacol*. 1990;101:363-9. [PMID: 1979509]
98. Hall J, Kaumann A, Wells F, Brown M. Beta-2-adrenoceptor mediated inotropic responses of human atria: receptor subtype regulation by atenolol [Abstract]. *Br J Pharmacol Proc* 1983;Suppl 93:116P.
99. Hall J, Kaumann A, Brown M. Increased Beta-2 adrenoceptor coupling to inotropic effects in human atrial myocardium due to atenolol treatment [Abstract]. *Br J Pharmacol Proc* 1989;Suppl 87:391P.
100. Molinoff PB, Aarons RD. Effects of drugs on beta-adrenergic receptors on human lymphocytes. *J Cardiovasc Pharmacol*. 1983;5 Suppl 1:S63-7. [PMID: 6188922]
101. Kraan J, Koëter GH, vd Mark TW, Sluiter HJ, de Vries K. Changes in bronchial hyperreactivity induced by 4 weeks of treatment with antiasthmatic drugs in patients with allergic asthma: a comparison between budesonide and terbutaline. *J Allergy Clin Immunol*. 1985;76:628-36. [PMID: 4056250]
102. Borst SE, Hui KK, Conolly ME. Beta-adrenergic receptors on human lymphocytes: comparison of down-regulation in vivo and in vitro. *Pharmacology*. 1990;40:325-9. [PMID: 1978356]
103. Cockcroft DW, McParland CP, Britto SA, Swystun VA, Rutherford BC. Regular inhaled salbutamol and airway responsiveness to allergen. *Lancet*. 1993;342:833-7. [PMID: 8104272]
104. Taylor DR, Sears MR, Herbison GP, Flannery EM, Print CG, Lake DC, et al. Regular inhaled beta agonist in asthma: effects on exacerbations and lung function. *Thorax*. 1993;48:134-8. [PMID: 8493626]
105. Wahedna I, Wong CS, Wisniewski AF, Pavord ID, Tattersfield AE. Asthma control during and after cessation of regular beta 2-agonist treatment. *Am Rev Respir Dis*. 1993;148:707-12. [PMID: 8103655]

106. Sbirrazzuoli V, Drici M, Garraffo R, Candito M, Gibelin P, Morand P, et al. Changes in lymphocyte beta-adrenoceptor density after diltiazem oral treatment. *Drugs Exp Clin Res*. 1989;15:223-9. [PMID: 2551622]
107. Brodde OE, Daul A, Michel MC. Subtype-selective modulation of human beta 1- and beta 2-adrenoceptor function by beta-adrenoceptor agonists and antagonists. *Clin Physiol Biochem*. 1990;8 Suppl 2:11-7. [PMID: 1982757]
108. Basso A, Piantanelli L, Cognini G, Giunta S, Andreoni A, Paciaroni E. Acebutolol-induced decrease of mononuclear leukocyte beta-adrenoceptors in hypertension. *Pharmacology*. 1985;31:278-83. [PMID: 2999840]
109. Jakubetz J, Schmuck S, Poller U, Fuchs B, Gorf A, Radke J, et al. Cardiac effects of beta-adrenoceptor antagonists with intrinsic sympathomimetic activity in humans: beta1- and/or beta2-adrenoceptor mediated? *J Cardiovasc Pharmacol*. 1999;33:461-72. [PMID: 10069683]
110. Wang TJ, Stafford RS. National patterns and predictors of beta-blocker use in patients with coronary artery disease. *Arch Intern Med*. 1998;158:1901-6. [PMID: 9759686]
111. Krumholz HM, Radford MJ, Wang Y, Chen J, Heiat A, Marciniak TA. National use and effectiveness of beta-blockers for the treatment of elderly patients after acute myocardial infarction: National Cooperative Cardiovascular Project. *JAMA*. 1998;280:623-9. [PMID: 9718054]
112. Soumerai SB, McLaughlin TJ, Spiegelman D, Hertzmark E, Thibault G, Goldman L. Adverse outcomes of underuse of beta-blockers in elderly survivors of acute myocardial infarction. *JAMA*. 1997;277:115-21. [PMID: 8990335]
113. Sial SH, Malone M, Freeman JL, Battiola R, Nachodsky J, Goodwin JS. Beta blocker use in the treatment of community hospital patients discharged after myocardial infarction. *J Gen Intern Med*. 1994;9:599-605. [PMID: 7853068]
114. Mooss AN, Hilleman DE, Mohiuddin SM, Hunter CB. Safety of esmolol in patients with acute myocardial infarction treated with thrombolytic therapy who had relative contraindications to beta-blocker therapy. *Ann Pharmacother*. 1994;28:701-3. [PMID: 7919552]
115. Dorow P, Bethge H, Tönnemann U. Effects of single oral doses of bisoprolol and atenolol on airway function in nonasthmatic chronic obstructive lung disease and angina pectoris. *Eur J Clin Pharmacol*. 1986;31:143-7. [PMID: 2879733]
116. Quan SF, Fenster PE, Hanson CD, Coaker LA, Basista MP. Suppression of atrial ectopy with intravenous metoprolol in chronic obstructive pulmonary disease patients. *J Clin Pharmacol*. 1983;23:341-7. [PMID: 6630583]
117. Formgren H. The effect of metoprolol and practolol on lung function and blood pressure in hypertensive asthmatics. *Br J Clin Pharmacol* 1976;3:1007-14.
118. Krauss S, Spitz E, Krauss A, Grizotzki B, Clement S. Treatment of hypertension in mild asthmatic patients with atenolol. *Angiology*. 1984;35:773-8. [PMID: 6507943]
119. Heller DA, Ahern FM, Kozak M. Changes in rates of beta-blocker use between 1994 and 1997 among elderly survivors of acute myocardial infarction. *Am Heart J*. 2000;140:663-71. [PMID: 11011343]

Current Author Addresses: Dr. S. Salpeter: Department of Medicine, Santa Clara Valley Medical Center, 751 South Bascom Avenue, San Jose, CA 95128.

Dr. Ormiston: Santa Clara Valley Medical Center, 751 South Bascom Avenue, San Jose, CA 95128.

Dr. E. Salpeter: Center for Radiophysics and Space Research, Cornell University, 612 Space Sciences Building, Ithaca, NY 14853.

Author Contributions: Conception and design: S.R. Salpeter, T.M. Ormiston.

Analysis and interpretation of the data: S.R. Salpeter, T.M. Ormiston, E.E. Salpeter.

Drafting of the article: S.R. Salpeter, T.M. Ormiston.

Critical revision of the article for important intellectual content: S.R. Salpeter, T.M. Ormiston.


Final approval of the article: S.R. Salpeter, T.M. Ormiston, E.E. Salpeter.

Provision of study materials or patients: S.R. Salpeter, T.M. Ormiston.

Statistical expertise: E.E. Salpeter.

Collection and assembly of data: S.R. Salpeter, T.M. Ormiston.

Appendix Figure. Results of search for and selection of trials.



Appendix Table. Characteristics of Included Studies*

Study, Year (Reference)	Design	Inclusion and Exclusion Criteria†	Participants	Dropout Rate	Mean Age or Age Range
			<i>n</i>	%	<i>y</i>
Adam et al., 1982 (52)	Single dose, double-blind, crossover	Inclusion: hypertension and reversible airway disease	10	0	65.1
Bauer et al., 1994 (65)	Crossover (1 week treatment and 1 week placebo), double-blind	Inclusion: hypertension and stable asthma Exclusion: cardiac disease, renal disease, hepatic disease, diabetes, pregnancy, adverse reaction to β -blocker	18	0	48.6
Benson et al., 1978 (57)	Single dose, crossover, single-blind	Inclusion: reversible airway obstruction, stable	12	14 before start of study	32.2
Butland et al., 1983 (64)	Crossover (4 weeks treatment and 4 weeks placebo), double-blind	Inclusion: COPD, FEV ₁ < 1 L with >20% reversal; inhalers, steroids allowed Exclusion: other lung disease or other serious disease	12	0	61
Chatterjee, 1986 (46)	Single dose, crossover, double-blind	Inclusion: asthma and hypertension, FEV ₁ /vital capacity < 30% Exclusion: pregnancy, heart or renal failure, antihypertensive or inhaler treatment	12	0	60
Chodosh et al., 1988 (43)	Single dose, crossover, double-blind	Inclusion: normotensive patients with stable asthma, FEV ₁ 60%–90% predicted Exclusion: cromolyn use; change in steroid or inhaler use; recent asthma attack, upper respiratory tract infection, or status asthmaticus	16	11 before start of study	39
Dorow et al., 1986 (63)	Crossover (12 weeks treatment, 4 weeks placebo), double-blind	Inclusion: hypertension and reversible airway obstruction Exclusion: severe hypertension	34	0	Unclear
Doshan 1986a (47)	Single dose, crossover, double-blind	Inclusion: mild asthma	15	6	19–55
Doshan et al., 1986 (48)	Single dose, crossover, double-blind	Inclusion: normotensive and asthma, FEV ₁ > 50% predicted Exclusion: cromolyn or steroid use	34	0	18–57
Ellis et al., 1981 (55)	Single dose, crossover, double-blind	Inclusion: reversible airway disease	14	0	Unclear
Falliers et al., 1986 (45)	Single dose, crossover, double-blind	Inclusion: asthma, FEV ₁ < 80% predicted Exclusion: hypertension, hematologic or cardiovascular disease, recent asthma attack or respiratory infection, status asthmaticus, or cromolyn therapy	18	0	21–60
Fenster et al., 1983 (66)	Crossover (1 week treatment, 1 week placebo), single-blind	Inclusion: reversible airway disease	6	0	48.6
Fogari et al., 1990 (61)	Crossover (1 week treatment, 2 weeks placebo), double-blind	Inclusion: hypertension and reversible airway disease Exclusion: cardiovascular disease, renal insufficiency	10	0	57
Greeffhorst and van Herwaarden, 1984 (50)	Single dose, crossover, double-blind	Inclusion: intrinsic atopic asthma, stable Exclusion: cardiovascular disease	8	0	29
Johnsson et al., 1975 (59)	Single dose, crossover, single-blind	Inclusion: asthma for > 2 years Exclusion: acute exacerbation, heart disease	7	0	44
Lammers et al., 1984 (49)	Single dose, crossover, double-blind	Inclusion: asthma, some with chronic bronchitis Exclusion: recent respiratory tract infection or increase in bronchoconstriction	8	0	39
Lammers et al., 1985 (68)	Crossover (4 weeks treatment and placebo), double-blind	Inclusion: COPD and hypertension, with average FEV ₁ reversal > 15%; all stable, with no recent respiratory tract infection or event Exclusion: none listed	8	0	52.7
Lammers et al., 1986 (44)	Single dose, crossover, single- and double-blind	Inclusion: asthma Exclusion: heart disease	11	0	6.6
Lammers et al., 1988 (60)	Single dose, crossover, single-blind	Inclusion: asthma, FEV ₁ 40%–74% predicted, stable without steroids or theophylline Exclusion: none listed	11	0	22–60
Lawrence et al., 1982 (51)	Crossover (single dose, then 3 weeks treatment and placebo), single-blind	Inclusion: asthma and hypertension	14	0	55.7
Lofdahl et al., 1981 (53)	Single dose, crossover, double-blind	Inclusion: intrinsic asthma, stable Exclusion: none listed	8	0	52
Nicolaescu et al., 1973 (62)	Crossover (3 days treatment and 3 days placebo), double-blind	Inclusion: severe asthma for > 5 years; steroids continued, inhalers held before measurements Exclusions: recent asthma attack	10	0	46.6
Nicolescu et al., 1972 (28)	Crossover (3 days treatment and 3 days placebo)	Inclusion: mild asthma for > 5 y Exclusion: none listed	10	0	43.7
Ruffin et al., 1979 (56)	Single dose, crossover, double-blind	Inclusion: asthma with episodic dyspnea Exclusion: FEV ₁ < 70% predicted, or symptoms out of control	12	0	30.8
Skinner et al., 1975 (58)	Single dose, crossover, double-blind	Inclusion: asthma	10	0	36.8
Tantucci et al., 1990 (42)	Single dose, crossover, double-blind	Inclusion: reversible airway disease, stable Exclusion: history of atopy, recent respiratory tract infection, contraindications to β -blocker, pregnancy	12	0	37.3
van den Bergh and van Herwaarden, 1981 (54)	Single dose, crossover, single- and double-blind	Inclusion: asthma, mild to moderate, stable	8	0	24
van Zyl et al., 1989 (67)	Crossover (4 weeks treatment, 2 weeks placebo), single- and double-blind	Inclusion: asthma and hypertension, FEV ₁ < 85% predicted	12	16	45.9

* COPD = chronic obstructive pulmonary disease; OT = ordinary tablets; SR = sustained release.

† Some studies did not list exclusion criteria.

Appendix Table—Continued

Active Interventions	Comparison Interventions	Outcomes Measured	Comments
1: metoprolol, 100 mg; atenolol, 100 mg 2: inhaled salbutamol after treatment	1: placebo 2: inhaled salbutamol after placebo	FEV ₁ , symptoms	Nonselective agents studied were labetalol and propranolol; SDs derived from individual-patient data or <i>P</i> values
Osmotic-release metoprolol, 204 mg/d, atenolol, 100 mg/d	Placebo	Symptoms	
1: acebutolol, 300 mg; atenolol, 100 mg 2: inhaled isoprenaline after treatment	1: placebo 2: inhaled isoprenaline after placebo	FEV ₁ , symptoms	Nonselective agents studied were propranolol and pindolol; SDs derived from individual-patient data or <i>P</i> values
Atenolol, 100 mg/d; metoprolol, 100 mg/d	Placebo	Symptoms, exercise tolerance	FEV ₁ reported as percentage of predicted normal value
1: atenolol, 100 mg; bisoprolol, 10 mg; bisoprolol, 20 mg 2: inhaled salbutamol after treatment	1: placebo 2: inhaled isoprenaline after placebo Atenolol, 100 mg/d; metoprolol, 100 mg/d	FEV ₁ symptoms	No SDs provided 2
1: metoprolol 200 mg 2: inhaled isoproterenol after treatment	1: placebo 2: inhaled isoproterenol after placebo	FEV ₁ symptoms	Nonselective agent studied was dilevalol; SDs derived from individual patient data or <i>P</i> values
Celiprolol, 100–600 mg/d	Placebo	FEV ₁ , symptoms, weekly inhaler use	SDs were reported for treatment effect and placebo effect separately
1: celiprolol, 400 mg; celiprolol, 600 mg; atenolol, 100 mg 2: inhaled albuterol after treatment	1: placebo 2: inhaled albuterol after placebo	FEV ₁ , symptoms	SDs were reported for treatment effect and placebo effect separately
Celiprolol, 200 mg; celiprolol, 400 mg; atenolol, 100 mg	Placebo	FEV ₁ , symptoms	Nonselective agent studied was propranolol; SDs were reported for treatment effect and placebo effect separately
1: atenolol, 50 mg; atenolol, 100 mg; atenolol, 200 mg 2: inhaled isoprenaline after treatment	1: placebo 2: inhaled isoprenaline after placebo	FEV ₁ , symptoms	Nonselective agent studied was propranolol; no SD data provided
1: metoprolol, 100 mg; metoprolol, 200 mg 2: inhaled isoproterenol after treatment	1: placebo 2: inhaled isoproterenol after placebo	FEV ₁ , symptoms	Nonselective agent studied was labetalol; SDs were reported for treatment effect and placebo effect separately
Metoprolol, 200 mg/d	Placebo	Increase in symptoms	FEV ₁ reported as percentage of predicted value
1: atenolol, 100 mg/d; celiprolol 200 mg/d 2: inhaled salbutamol after treatment	1: placebo 2: inhaled salbutamol after placebo	FEV ₁ , symptoms	Nonselective agents studied were propranolol and oxprenolol; SDs were reported for treatment effect and placebo effect separately
1: metoprolol, 100 mg; acebutolol, 400 mg 2: intravenous terbutaline after treatment	1: placebo 2: intravenous terbutaline after placebo	FEV ₁ , symptoms	SDs were reported for treatment effect and placebo effect separately
Intravenous metoprolol, 0.12 mg/kg body weight	Intravenous placebo	FEV ₁ , symptoms	Isoprenaline given intravenously, but FEV ₁ not reported; nonselective agent studied was propranolol; SDs derived from individual-patient data or <i>P</i> values
1: bisoprolol, 10 mg; bisoprolol, 20 mg; metoprolol, 100 mg 2: inhaled terbutaline after treatment	1: placebo 2: inhaled terbutaline after placebo	FEV ₁ , symptoms	SDs were reported for treatment effect and placebo effect separately
Metoprolol, 100 mg twice daily	Placebo	Symptoms	Nonselective agent studied was pindolol
1: xamoterol, 200 mg; atenolol, 100 mg 2: inhaled terbutaline after treatment	1: placebo 2: inhaled terbutaline after placebo	FEV ₁ , symptoms	Placebo given single-blind, treatments given double-blind; SDs were reported for treatment effect and placebo effect separately
1: atenolol, 50 mg 2: inhaled terbutaline after treatment	1: placebo 2: inhaled terbutaline after placebo	FEV ₁ , symptoms	No SDs provided
1: atenolol, 100 mg; metoprolol, 100 mg 2: inhaled salbutamol after treatment 3: atenolol, 100 mg/d; metoprolol, 100 mg twice daily	1: placebo 2: inhaled salbutamol after placebo 3: placebo	FEV ₁ , symptoms, asthma attacks, weekly inhaler use	SDs derived from individual-patient data or <i>P</i> values
1: atenolol, 100 mg; metoprolol 100 mg 2: terbutaline, intravenous then inhaled, after treatment	1: placebo 2: terbutaline, intravenous then inhaled, after placebo	FEV ₁ , symptoms	SDs derived from individual-patient data or <i>P</i> values
1: practolol, 200 mg/d 2: inhaled orciprenaline after treatment	1: placebo 2: inhaled orciprenaline after placebo	FEV ₁ , symptoms	SDs derived from individual-patient data or <i>P</i> values
1: practolol, 50 mg four times daily 2: inhaled orciprenaline after treatment	1: placebo 2: inhaled orciprenaline after placebo	FEV ₁ , symptoms	SDs derived from individual-patient data or <i>P</i> values
Metoprolol, 100 mg	Placebo	FEV ₁ , symptoms	Nonselective agents studied were propranolol and timolol; no SDs provided
1: acebutolol, 300 mg; practolol, 300 mg 2: inhaled isoprenaline after treatment	1: placebo 2: inhaled isoprenaline after placebo	FEV ₁ , symptoms	SDs were reported for treatment effect and placebo effect separately
Osmotic-release metoprolol, 204 mg; atenolol, 100 mg	Placebo	FEV ₁ , symptoms	SDs derived from individual-patient data or <i>P</i> values
1: metoprolol OT, 100 mg; metoprolol OT, 200 mg; metoprolol SR, 200 mg 2: intravenous terbutaline after treatment	1: placebo 2: intravenous terbutaline after placebo	FEV ₁ , symptoms	Single-blind; metoprolol, 100 mg, and placebo; double-blind: metoprolol, 200 mg, 2 doses; SDs derived from individual-patient data or <i>P</i> values
Atenolol, 100 mg/d; celiprolol, 400/d	Placebo	Symptoms, weekly inhaler use	Placebo run-in, single-blind; treatment double-blind