



DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
NATIONAL INSTITUTES OF HEALTH
BETHESDA, MARYLAND 20014

April 5, 1976

Daniel S. Dixler, Ph.D.
Keller and Heckman
1150 17th Street, N.W.
Suite 1000
Washington, D.C. 20036

Dear Dr. Dixler:

Thank you very much for your letter of March 9. I appreciate your kind words regarding the talk I gave down at Research Triangle Park on March 2, and I am enclosing a copy of the talk in the form in which it will be published in Environmental Health Perspectives.

After receiving your letter, I called NIOSH regarding fabrication plant angiosarcoma cases, and was informed that the NIOSH data presented as my Table 4 have not been withdrawn. You might wish to communicate further regarding this matter with Dr. J. William Lloyd of NIOSH.

Although I did not discuss "community" cases of angiosarcoma in my talk, I am grateful for the information on community cases you enclosed in your letter of March 9.

Again, my thanks to you and Mr. Heckman for your help.

Yours truly,

Myra Karstadt, Ph.D.
Consultant

ASI 00023042

PVC:HEALTH IMPLICATIONS AND PRODUCTION TRENDS

MYRA KARSTADT

MARCH 2, 1976

ASI 00023043

In January 1974, B. F. Goodrich Chemical Company announced that three reactor workers in a polyvinyl chloride (PVC) polymerization plant had developed a rare cancer, angiosarcoma of the liver. Exposure to vinyl chloride was linked to development of angiosarcoma of the liver by subsequent epidemiological studies. Animal studies, several initiated before the B.F. Goodrich announcement, also link exposure to vinyl chloride (VCM) with development of cancer.

Regulations promulgated in the United States of America (USA) since early 1974 have resulted in the lowering of VCM concentrations in the atmosphere of PVC manufacturing plants, and decreased concentrations of VCM in PVC resin. This has reduced the cancer risk of PVC production workers, workers in PVC fabrication plants and the ultimate consumers of fabricated PVC products.

We are now at a crucial stage as regards studies of the effects of plastics and plastics components on human health. Human exposure to massive doses of VCM, whether due to uncontrolled venting of PVC polymerization vessels or the voluntary venting in a home bathroom of an aerosolized VCM hair spray, is essentially over. The time has come for study of possible health hazards posed by the complex plastic system called PVC.

Why be concerned about PVC? A publication(1) states that:

"Total world employment in the VCM and PVC producing

...likely to be well over 70,000 workers. Those employed in industries which use PVC as a basic element are likely to total in the millions. Those who come in contact with PVC every day in some form or other probably make up about at least one-third of the human race."

What are the human health effects of PVC? There are several components of PVC which, on their own, could affect human health adversely: VCM, PVC dust considered as a dust, additives of various sorts ("additives" for our purposes includes stabilizers, colorants, flame retardants, plasticizers, fillers etc.)

But one cannot consider the components of PVC solely as independently-acting potentially toxic agents. PVC is a complex chemical system with little-understood interactions, both physical and chemical, among its components.

Leaving aside the multi-component interaction problem for the time being, consider the individual components of PVC which may pose risk of harm to humans.

VCM is already identified as a cancer-causing agent. Even given the reductions of VCM in PVC resin, which have been achieved by changes in both production and compounding procedures, some RVCN(residual vinyl chloride monomer) is still present in PVC.

In 1974 Rowe(2) estimated the average monomer concentration of PVC resin to be approximately 1000 ppm. As shown in Table 1, RVCN concentrations of 100 ppm or less for suspension resins before compounding were not unusual by April 1975. An advertisement(3) run/by a major manufacturer of PVC stated that general purpose PVC homopolymers and blending resins were being produced with a maximum of 10 ppm, and usually less than 5 ppm, RVCN.

Tables 1, 2 and 3, taken from the United States Environmental Protection Agency(EPA) publication "Standard Support and Environmental Impact Statement:Emission Standard for Vinyl Chloride"(4) give some idea of the present RVCN in several PVC resins. In particular, note company E in table 1. 86.8% of suspension resin production is said to be at 0-50 ppm RVCN(with a footnoted qualification that "some values are speculative until improved stripping is installed".) Note also that for company A, 60% of whose resin had a RVCN concentration of 2000-4000 ppm in April 1975, with the remaining 40% at 4000-6000 ppm,"all suspension resins (were) to be reduced to less than 400 ppm RVCN by 7/75".

Health problems which may be associated with RVCN in finished PVC resin are of importance to workers and consumers. The residual VCM in the old high-residue PVC evidently posed problems for workers in fabrication plants. Data collected by the United States National Institute for Occupational Safety and Health(NIOSH) indicate that several angiosarcoma cases were identified in fabrication plant workers in this country and abroad(Table 4)(5).

NIOSH data(Table 5)(6) presented in late 1974 indicated that breathing zone samples in fabrication plants turned up relatively low concentrations of VCM-even at a time when RVCN concentrations were high compared to today's values. Even at that time, as Wagoner put it:"60% of the samples were less than 1 ppm".(7). 1 ppm is the maximum exposure level (8 hour time weighted average) currently permitted by the United States Occupational Safety and Health

ASI 00023046

Administration(OSHA) standard./Again quoting Wagener(8):

"I think it(the occurrence of angiosarcoma of the liver in fabrication plant workers-author's note) points to the possibility of low level exposure being obviously very hazardous." (8).

A recent contract study(9) prepared for NIOSH reported on fabricating plant VCM concentrations in 1975. The VCM concentrations, as shown in Table 6, are, in a number of cases, very low(less than 0.01 ppm, the limit of detection). However, certain work stations had values which exceeded the 1 ppm 8 hour time-weighted average currently permitted by OSHA.

How low must you go before RVCM ceases to represent a hazard? A "no-effect" level for VCM carcinogenesis has not been found to date;cancer has been found in all groups of animals tested with non-zero concentrations of inhaled VCM. Just how low are the VCM concentrations in fabrication plants now? Publication of monitoring data would be useful for analysis of possible risks of harm to health.

A fabricated PVC product category with wide consumer exposure is PVC automobile upholstery, rooftops and interior fittings. The EPA has a contract study in progress entitled "Sampling of Automobile Interiors for Vinyl Chloride Monomer". (10).

According to Hedley(11), who is working on the EPA contract study, of 7 cars in which 30-60 minute grab samples were taken in 1975 under varying heat and sunlight conditions, one reading was 1.2 ppm VCM, one 0.4 ppm and the remaining five less than 0.05 ppm(the detection limit). The car with the highest

reading was sampled one month after assembly-under the following temperature conditions: ambient 56⁰C., interior air of car 60⁰C., seat surface 66⁰C.

No attempt was made to follow any decrease in off-gassing of RVCM over time.

Given the number of individuals exposed to "new car atmospheres" and cars or other motor vehicles generally, some further data on the RVCM content of PVC auto fittings might be useful.

Even if risk to consumers from RVCM in PVC seat covers is low, what is the risk of harm to persons working in automobile factories?

A recent study by Infante et al(12) is especially interesting. The data collected by Infante and his coworkers have been interpreted to suggest that VCM may exert a mutagenic effect on sperm cells of PVC production workers, leading to an increase in fetal deaths among pregnancies of workers' wives. Infante's findings are not surprising in light of the demonstration by ~~Ducatron et al.~~ ~~Selikoff~~ (13) of chromosome aberrations in PVC production workers and the finding by Ames(14) that VCM is mutagenic in a Salmonella test system.

The setting of vinyl automobile upholstery and other vinyl fabric automobile fittings is, evidently, in large part "woman's work". Even given the multi-component nature of the PVC in auto fittings, it might be interesting to determine the incidence of cancer, stillbirths and birth defects in offspring among women vinyl auto upholstery workers.

What happens to PVC on pyrolysis? There seems to be some disagreement as to whether VCM is liberated(15) or whether the principal health threat from burning PVC is caused by evolution of HCl(16). Has there been a definitive demonstration that VCM is not evolved during pyrolysis of PVC? What about depolymerization on aging? in sunlight? in heat?

PVC exposure has been linked both to dermatitis and Raynaud's syndrome/occupational acroosteolysis./Dermatitis (16a). could be attributable to several of the additives in PVC, or possibly to VCM itself. A catalogue of resin components which have caused skin sensitization reactions would be useful.

One of the important us consumer uses of PVC is in food packaging. Regulations(17) recently proposed by the United States Food and Drug Administration(FDA) deal with PVC food packaging. Although industry data(18) have been presented^{to FDA}/which indicate migration of VCM from PVC at below 50 ppb(the "lowest detectable" level established by the FDA proposal), challenges(19) have been raised to permitting use of PVC food wrap if there is

any RVCM capable of migrating.

The debate about PVC food wrap points up what may be the salient characteristic of PVC considered as a source of residual VCM. RVCM contained in PVC has now joined the ranks of other carcinogenic substances whose presence in trace amounts--the amounts limited only by the means of detection--in the US food supply is forbidden by the Delaney Clause, an amendment to the United States Food, Drug and Cosmetic Act. That clause reads in pertinent part:

"...no additive shall be deemed to be safe if it is found to induce cancer when ingested by man or animal, or if it is found, after tests which are appropriate for the evaluation of the safety of food additives, to induce cancer in man or animal..." (20).

FDA's attempt to set a non-zero tolerance for VCM in food wrap has been challenged as illegal due to conflict with the Delaney Clause(21). It may be that final promulgation of a proposal setting a non-zero tolerance for RVCM in food wrap will be challenged in the courts.

Efforts to reduce VCM residual in PVC have come a long way in just over two years. The industry has demonstrated its ability to reduce VCM levels in PVC production plants from hundreds or thousands of ppm to a level close to or at the OSHA & hoo. TWA of 1 ppm.

• ASI 00023050

Since chemical reactions rarely go to completion and are usually reversible, will it ever be possible to produce a PVC resin completely free of residual VCM and incapable of breaking down to release VCM? The stumbling block for the continued health and growth of the PVC industry may well prove

to be not the threat of massive doses of VCM and high potential risk of cancer for the few, but the threat of low doses of VCM and a low but non-zero risk of cancer for the many.

The presence of BVCM in PVC represents a potential health hazard for the large number of humans exposed to resin or, more commonly, finished plastic goods. The dusty nature of PVC resin may represent a health hazard for persons with an occupational exposure to PVC.

Where could PVC dust represent a significant hazard? In a polymerization plant, work stations which could be dusty include drying and bagging areas, storage rooms or silos where PVC resin is held for shipment, mixing areas in compounding and/or fabrication facilities, and baghouses or cyclones in production or fabrication facilities. If the dusty operations are not completely isolated from the remainder of the production or fabrication facility, dust hazards could be present throughout the plant.

How dusty do PVC plants get? Two photographs (Figure 1 and Figure 2) (22) taken in 1975 in a large PVC polymerization facility in the US may be of interest. The first of the two pictures gives a general idea of the powdery coating on the equipment in the bagging area. The second is a picture of the floor of the plant-note the depth of the dust as indicated by the footprints.

PVC dust concentration data are not readily available; at least, the author had difficulty obtaining such data from labor, government, academic and industry sources. It

would be very helpful to those interested in problems of dust-caused disease for industry, labor or government to take measurements of PVC dust in the workplace and make them public.

Particle sizes within the respirable particle range have been noted in a US PVC polymerization plant(23). (This was not the same plant as the one in which the photographs displayed as Figures 1 and 2 were taken.)

A number of clinical studies have demonstrated pulmonary abnormalities in PVC production and fabrication workers.(24). Abnormalities have ranged from decreased pulmonary function, as measured by FEV, to pulmonary fibrosis. Given the data currently available, work is needed to separate the effects on the lung and respiratory system of VCM from those of PVC dust.

Experimental studies associate pulmonary fibrosis with exposure of animals to PVC dust. In a recent paper from Italy, Frongia et al(25) performed a series of studies with a simple basic plan-place guinea pigs or rats in cages in the bagging area of a PVC polymerization plant and observe the effects on the animals' lungs. These animal experiments were inspired by a clinical report of PVC-induced pneumoconiosis.(26). The Frongia group, after microscopic examination of tissues of the animals under study, concluded that they had demonstrated a pneumoconiotic reaction brought on by PVC dust. Experiments

reported as "in progress" as of late 1974 were designed to determine whether the pulmonary fibrosis was reversible on cessation of exposure to PVC. Data on reversibility of pulmonary and respiratory effects of PVC exposure should also be obtained for human populations.

Work is needed to determine whether VCM exposure heightens pulmonary response to PVC dust or whether, conversely, PVC dust may convey low levels of VCM or additives to the lungs. (26a) Possible enhancement of dust effects by other chemicals in the work atmosphere and/or the additives in the PVC resin should also be investigated.

PVC dust could also be swallowed. Volkheimer(27) has found that PVC particles can be "persorbed" from the intestine into the lymph or blood systems. The effect of locally high concentrations of PVC particles carrying minute amounts of potentially leachable RVCM or additives merits investigation, especially as concerns contact with lung or liver tissue.

The possibility of PVC dust serving as a skin irritant should be explored. PVC dust could serve as an efficient means for bringing a sensitizing compound into contact with the skin.

The suggestive nature of the clinical and experimental data available at this time supports reduction of PVC dust levels in polymerization, compounding and fabrication facilities. The OSHA standard(28) does not include PVC dust protection or reduction regulations, although such regulations were recommended by NIOSH(29).

ASI 00023053

Finally, the question of additives.

The final table (Table 6) lists several major categories of PVC additives-colorants, fibrous reinforcements, flame retardants, plasticizers and stabilizers.(30). Based on experimental and/or clinical data, questions are in order as to the safety of representatives of each of these chemical classes.

*
For instance, Ames(31) has recently reported that a certain flame retardant compound used in children's sleepwear is highly mutagenic in a Salmonella test system. Compounds exhibiting mutagenic activity in this bacterial test system are likely to be carcinogens as well. What about the toxicity of flame retardants used in PVC plastics?

In September 1972, NIEHS held a conference on phthalic acid esters.(32). What is the current thinking of regulators and researchers regarding the safety of these chemicals?

There is considerable documentation of toxic effects of certain organotin compounds.(34). The biocidal activity of organotin compounds, several of which are being used as pesticides at present, should cause concern about possible effects of organotin plastics additives on human health.

Possibility of complex interactions among PVC plastics components and between the components and living systems should be emphasized. Most probably, very little is known about these interactions.

ASI 00023054

Since last Fall, the author has been working, as a consultant to NIEHS, on a project designed to identify chemicals which should be studied for possible adverse health effects

for humans. Because of its wide population exposure due to enormous production volume and broad spectrum of uses, and health problems already associated with VCM and PVC dust and certain additives, PVC seems a likely candidate for further investigation.

Given the current state of scientific knowledge-at least as regards data available for public scrutiny-producers and fabricators of PVC cannot assure the public that PVC plastics are safe for human use.

Therefore, interested scientists the world over, whether working for industry, government or labor unions, or engaged in research at universities, should make a concerted effort to collect and analyze such data as are available at present on possible risks of harm to human health attributable to PVC. Scientists should identify research which should be carried out to determine whether or not the family of PVC plastics is safe for general use, and seek the necessary funding to carry out whatever studies are needed.

TABLE 1

Table 4-6 APRIL 1975 STATUS OF SUSPENSION RESIN STRIPPING^{1/}

Company	Number of Suspension Resin Grades	% of Suspension Resin Production	R VCM (ppm)
A ^{2/}	-	60	2000-4000
		40	4000-6000
B ^{2/}	-	100	2000-5000
C	5 Grades	100	<400
D	-	80	400
		20	500-700
E	13 Grades	86.8	0-50 ^{3/}
		5.4	100
		3.8	500
		3.2	1000
		.8	4000
F	15 Grades	100	<400
G	4 Homopolymer	6	1500
		14	2500
		80	100
	4 Copolymer	16	300
		84	400
H	4 Grades	100	2000-10,000
I	-	38	400
		9	600
		25	800
		5	1800
		15	3600
		8	4000

^{1/}Based on data from responses to 3/31/75 section 114 request.

^{2/}All suspension resins to be reduced to <400 ppm R VCM by 7/75.

^{3/}Some values are speculative until improved stripping is installed.

TABLE 2

Suspension Resins

ASTM ^{1/} Cell Class	% Prod.	PPM VCM into stripper	Avg. ppm VCM out of stripper (dry basis)	Avg. ppm out of dryer	Emission factor kg VCM/ 100 kg PVC
GP1-26250	5	(40-60) x 10 ³	3,050	308	0.27
GP2-16340	5	(40-60) x 10 ³	2,050	162	0.18
GP3-15340	13	(50-70) x 10 ³	1,200	104	0.09
GP4-15340	35	(60-90) x 10 ³	1,050	95	0.09
GP6-15343	24	(90-120) x 10 ³	950	82	0.08
GP1-14443	0	(80-100) x 10 ³	600 ^{2/}	25	0.06
GP2-14443	1	(80-100) x 10 ³	550	16	0.06
GP3-15433	1	(80-100) x 10 ³	500	21	0.05
GP5-15433	5	(90-110) x 10 ³	450	15	0.05
GP6-15433	9	(100-140) x 10 ³	450	18	0.05
<u>Copolymers</u>					
C11-8500	1	(40-60) x 10 ³	2,100	280	0.18
C11-3500	1	(90-110) x 10 ³	1,000	95	0.09

^{1/}ASTM cell class is standard definition of resin. ASTM-D-1755

^{2/}Based on limited data (less than three samples).

This company plans to install secondary strippers to further reduce the residual vinyl chloride content shown above by increasing the stripping temperature to as high as 85°C (185°F) and the residence time to one hour.

4.12.1.2 Company L^{2/}

The values below were obtained by conventional steam stripping at the conditions shown. Values are for 1973 and 1974.

Table 4-5. APRIL 1975 STATUS OF DISPERSION RESIN STRIPPING WITH PROJECTION OF FUTURE CAPABILITIES^{1/}

Plant	Current RVCM ^{2/} After Stripper	(ppm) In Product	Projected RVCM (ppm) After Stripper	Time to Implement Controls Needed to Meet Projected Levels
A	30,000	1-3	20,000	6 months
B	50,000	5	10,000	6 months
C	20,000	5-15	4000-6000	48 months
D	1500-4000	-	-	-
E	15,000	1-10	6000	12-18 months
F	13,000-18,000	10	4500	30 months
G	2000-10,000	25	500-1000	42 months
			2000	30 months
			400	48 months
H	14,000-18,000	5-30	1200	30 months
I	2000	<1	-	-

^{1/} From responses to March 31, 1975, section 114 request

^{2/} RVCM = Residual Vinyl Chloride Monomer

TABLE 4

NIOSH
September 12, 1975

TABLE II

REPORTED CASES OF ANGIOSARCOMA OF THE LIVER AMONG
NON-POLYMERIZATION WORKERS EXPOSED TO VINYL CHLORIDE

COUNTRY	CASE #	BIRTH DATE	1st VC or PVC EXPOSURE	DX ANGIO-SARCOMA	AGE AT DX	YRS 1st EXP TO DX	TOT YRS EXP	DATE OF DEATH	
Fed. Rep. Germany	03*	07-16-30	06-09-52	03-00-73	43	21	14	10-10-73	Note 6
Fed. Rep. Germany	07*	05-09-36	00-00-60	00-00-74	38	14	03	12-16-74	Note 9
Great Britain	02*	09-08-14	00-00-46	02-00-70	55	24	11	12-00-70	Note 1
Italy	01	06-15-34	00-00-65	04-19-71	36	6	3	04-16-71	Note 2 & 7
Sweden	02*	11-27-11	00-00-45	05-15-72	61	27	23	08-16-72	Note 3
United States	14	00-00-13	08-18-38	06-00-73	60	36	00	07-03-73	Note 4 & 8
United States	15*	00-00-25	00-00-00	07-00-72	47	00	00	02-15-73	Note 5

Note 1 Pouring PVC oil mixture onto fabric bases

Note 2 Production of PVC sacks

Note 3 Production of Vinyl Chloride

Note 4 Machine operator covering electrical wire with PVC plastic insulation

Note 5 Accountant at plant making PVC fabric

Note 6 Loading pesticide cans with VC propellant

Note 7 Angiosarcoma involving liver, lung, and pericardium. Although difficult to determine, primary site seems to be pericardium.

Note 8 Diagnosis: Sarcoma (possibly "angiosarcoma"), liver. Possibility of generalized neoplasm of the reticuloendothelial cell system cannot be ruled out.

Note 9 Assistant Factory Chemist


* Indicates microscopically confirmed angiosarcoma of the liver.

"00" Indicates unknown data.

SUMMARY OF VCM CONCENTRATIONS IN PVC FABRICATION FACILITIES

Type of plant	Concentrations			Fraction > 1 parts per million
	Maximum	Minimum	Mean	
Film and mold (2 plants):				
Wash data:				
(a) Dressing zone samples.....	12	<.3	3	2.5
(b) Area samples.....	8	<.3	1.2	1.11
(c) Source samples.....	340	<.3	18	10.21
Pipe and extrud (2 plants):				
Industry data:				
(a) Dressing zone samples.....	<10	<.1	<1	4.21
(b) Area samples.....	35	<.1	6	16.20
(c) Source samples.....	540	<.1	142	10.17

FIGURE 1



ASI 00023061

ASI 00023062

FIGURE 2



ADDITIVES

colorants

fibrous reinforcements

asbestos

flame retardants

plasticizers

phthalic acid esters

stabilizers

organotin compounds

References

1. Levinson, C.: Vinyl Chloride: a case study of the new occupational health hazard, International Chemical Federation, Geneva, Switzerland, 1974, 15.
2. Rowe, V. K.: comments in discussion, 246 Ann. N.Y. Acad. Sci. 317,(1975).
3. Tenneco PVC resins have VCM down for a 10 count...down to 10 parts per million(advertisement), The Wall Street Journal, July 9, 1975, 4.

The advertisement states(in part):
"At Tenneco, we've developed a unique new production technology that allows us to produce our general purpose PVC homopolymer, blending and dispersion resins with no more than 10 parts per million of VCM(vinyl chloride monomer). In fact, the typical VCM content of these homopolymer and blending resins is less than 5 parts per million and under 3 parts for our dispersion resins. And, we are producing our flexible compounds at non-detectable(less than 1 ppm) levels.
"This is a 98% to 99% reduction from what was once the typical practice of the industry for residual VCM."
4. EPA-450/2-75-009, U.S. Environmental Protection Agency, Office of Air and Waste Management, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina (October 1975).
5. The U.S. NIOSH has been maintaining a register of VCM-related angio-sarcoma cases; this is one of the continuing series of tabulations published by the agency.
6. Vinyl Chloride, Hearing before the Subcommittee on Environment of the Committee on Commerce, United States Senate, August 21, 1974, Serial No. 93-110, U.S. Government Printing Office, Washington, D.C. 1974, 51.
7. 6, above, 50.
- 7a. Standard for Exposure to Vinyl Chloride, Occupational Safety and Health Administration, Department of Labor, 39 Fed. Reg. 35890 (October 4, 1974).
8. 6, ibid.
9. Barnhart, W. L., Toney, C.R., and J. B. Devlin: Environmental/Industrial Hygiene Surveys of Vinyl Chloride Monomer Manufacturing Operations and Operations where Polyvinyl Chloride and Copolymers of Polyvinyl Chloride are Processed, Contract No. CDC-99-74-50, U.S. Department of Health, Education and Welfare, Public Health Service, Center for Disease Control, National Institute for Occupational Safety and Health (August 1975).
10. Fearheller, W. R., Cheng, J. T., and R. J. McCormick: Sampling of Automobile Interiors for Vinyl Chloride Monomer, EPA Contract No. 68-02-1404, Task 1, Cost and Performance Report, 1 June 1975 to 30 June 1975, Environmental Protection Agency, Research Triangle Park, North Carolina.
11. Personal communication, February 23, 1976.

12. Infante, P.F., Wagoner, J.K., McMichael, A.J., Waxweiler, R.J. and H. Falk: Genetic Risks of Vinyl Chloride, Lancet (in press).
13. Ducatman, A., Hirschorn, K. and I. J. Selikoff: Vinyl Chloride exposure and human chromosome aberrations, 31 Mutation Res. 163 (1975).
14. McCann, J., Choi, E., Yamasaki, E. and B. H. Ames: Detection of carcinogens as mutagens in the *Salmonella*/microsome test: Assay of 300 chemicals, 72 Proc. Nat. Acad. Sci. USA 5135 (1975).
15. Boettner, E.A., Ball, G.L. and B. Weiss, Combustion Products from the Incineration of Plastics, EPA Grant No. EP-00386, Program Element No. 1D2063, EPA-670/2-73-049, 25, 26 (July 1973).
16. See, for instance, Dyer, R.F. and V.H. Esch, Polyvinyl Chloride Toxicity in Fires, 235 J.A.M.A. 393 (1976); Hornblower, M.; Plastic Fumes Insidious, The Washington Post, December 11, 1975, A8.
- 16a. See, generally, 1 above; 246 Ann. N.Y. Acad. Sci. (Proceedings of International Workshop on Vinyl Chloride, May 1974), (1975).
17. Department of Health, Education and Welfare, Food and Drug Administration: Vinyl Chloride polymers in contact with food, Notice of Proposed Rule-making, 40 Fed. Reg. 40529 (September 3, 1975).
18. Heckman, J., counsel (Washington, D.C.), Society of the Plastics Industry, Inc.: Submission for the Record in FDA Docket No. 75-N-0190; Vinyl Chloride Polymers in Contact with Food; Notice of Proposed Rule-making, 40 Fed. Reg. 40529, September 3, 1975 (December 19, 1975).
19. Johnson, A. and S. Wolfe, Health Research Group, Washington, D.C.: Submission for the record in proposed FDA regulation Vinyl Chloride Polymers in contact with food (December 19, 1975).
20. 21 USC § 348(c)(3)(a).
21. 19, above.
22. Photographs, taken in unidentified PVC polymerization plant, provided by Mr. Steven Wodka, Oil, Chemical and Atomic Workers International Union, Citizen-Legislative Department, Washington, D.C.
23. Personal communication, William J. Nicholson, Mt. Sinai School of Medicine, February 18, 1976.
24. See, for instance, Lilis, R., Anderson, H., Nicholson, W.J., Daum, S., Fischbein, A.S. and I. J. Selikoff: Prevalence of Disease among Vinyl Chloride and Polyvinyl Chloride Workers, 246 Ann. N.Y. Acad. Sci. 22 (1975); Miller, A., Teirstein, A.S., Chuang, M. and I. J. Selikoff; Changes in Pulmonary Function in Workers Exposed to Vinyl Chloride and Polyvinyl Chloride, 246 Ann. N.Y. Acad. Sci. 42(1975); Wegman, D.;

Comments in discussion, 246 Ann. N.Y. Acad. Sci. 20(1975).

25. Frongia, R., Spinazzola, A. and A. Bucarelli; Lesioni pulmonari sperimentali da inalazione prolungata di polveri di PVC in ambiente di lavoro, 65 Med. Labor 321(1974) (Title translation provided by authors): Experimental lung damage from prolonged inhalation of airborne PVC dust.
26. Szende, B., Lapis, K., Nemes, A. and A. Pinter: Pneumoconiosis caused by the inhalation of PVC dust, 61 Med. Labor 433(1970).
- 26a. According to Miller et al., (24, above, at 51): "...If exposure to vinyl chloride and/or PVC is causally related to air flow impairment, we do not know which substance is more pathogenic. The fine dust of the polymer might be more likely to concentrate in and damage the small Airways. This dust, and other small respirable particles, perhaps from tobacco smoke, may also serve as carriers for molecules of a monomer gas to settle in these airways."
27. Volkheimer, G.: Hematogenous dissemination of ingested polyvinyl chloride particles, 246 Ann. N.Y. Acad. Sci. 164(1975).
28. U.S. Department of Labor, Occupational Safety and Health Administration, Standard for Exposure to Vinyl Chloride, 39 Fed. Reg. 35890 (October 4, 1974).
29. For the NIOSH recommendations, see 1, above.

To date, neither the US OSHA nor the US EPA has taken any specific steps to reduce PVC dust considered as dust. Dust reduction and protection was recommended to the OSHA by NIOSH, EPA on p. 49 of its standard support document (4, above) states:

"With regard to the potential problem of polyvinyl chloride particulate as a possible cause of pneumoconiosis, NIOSH is currently involved in experimental studies on the effects of the particulate in animals. The extent of public exposure (as opposed to occupational exposure) to ambient concentrations of the particulate is unknown at this time. Ambient measurements of polyvinyl chloride particulate have not been made by EPA in the vicinity of industrial sources because no technology is currently available for separating polyvinyl chloride particulate from total suspended particulate. As data become available from NIOSH and other sources on the health effects of polyvinyl chloride particulate, EPA may find that it is necessary to reevaluate the need to propose standards for polyvinyl chloride particulate."

30. See, generally, Schoengood, A.A.: Additives, 28 Soc. Plastics Engineering J. 21 (1972).

31. Environmental Defense Fund, Petition pursuant to 15 U.S.C. § 2059 to the Consumer Product Safety Commission to commence a proceeding for the issuance of a consumer product safety rule, 7 (March 24, 1976).

32. Proceedings of the meeting are printed in 3 Environmental Health Perspectives (1973).

33. (citation deleted)

34. Piver, W.T.: Organotin Compounds: Industrial applications and biological investigation, 4 Environmental Health Perspectives 61 (1973).