

Katherine E. Reed, Ph.D.
Staff Vice President

3M Environmental Technology
and Safety Services

900 Bush Avenue
Building 42-2E-26
PO Box 33331
St Paul, MN 55133-3331
651 778 4331

ME #
64661

AR 226 - 1153

RECEIVED
OFFICE

2002 DEC 13 AM 10:08

8EHQ - 1202 - 00373



CERTIFIED MAIL

December 4, 2002

Document Processing Center (7407M)
EPA East - Room 6428 Attn: Section 8(e)
Office of Pollution Prevention and Toxics
US EPA
1200 Pennsylvania Ave., NW
Washington, DC 20460-0001

COMPANY SANITIZED

2002 DEC 31 AM 10:08

TSCA 8(E) SUBSTANTIAL RISK SUPPLEMENTAL NOTICE ON: Perfluorooctane sulfonate (PFOS), perfluorohexanesulfonate (PFHS), N-methyl-N-acetyl-perfluorooctanesulfonamide (M570)
Reference 8EHQ -1180-373, 374

Dear Sir:

3M has received the draft report for a repeated dermal contact absorption/toxicology study in rats conducted with four mill-applied and aftermarket carpet/fabric/upholstery protector products that 3M ceased manufacturing as part of its perfluorooctanyl chemistries phase-out. The attachment hereto provides the composition for each of these products. The results indicate that repeated daily dermal contact over a 28-day period with these formulations, in the liquid state or after application and drying on gauze, resulted in increases above background controls in the concentration of perfluorooctanesulfonate (PFOS) in liver and serum (all but one product), and of perfluorohexanesulfonate (PFHS) and N-methyl-N-acetyl-perfluorooctanesulfonamide (M570) in serum (not measured in liver) for certain products. M570 is an intermediary metabolite of a manufacturing residual (N-methyl-perfluorooctanesulfonamido ethanol (N-MeFOSE)) present in certain products, which can be further metabolized to PFOS. No toxicity attributable to any sample was evident in this study, either on the skin, or systemically.

The study was conducted by the Fraunhofer Institute of Toxicology, Hannover, Germany. Cotton gauze patches (~1 square inch) to which the liquid test material had previously been applied and allowed to dry, or liquid samples (0.1 ml) applied to gauze without drying, were applied to the shaved dorsal skin of male and female rats in a semi-occluded fashion for six hours each day for up to 28 days. The preparation of the samples dried on cotton gauze involved saturating the gauze with liquid product, wringing the saturated gauze through a roller press at 100 pounds pressure and allowing the wrung gauze to air dry at room temperature. As such, the application of product to gauze did not precisely

8EHQ-60-373

0008114527

SAFETY

Office of Pollution Prevention and Toxics
Page No. 2
December 4, 2002

mimic normal application on carpet, fabrics and upholstery but was expected to represent the higher end of normal product application. The liquid samples were adjusted to provide approximately the same amount of product solids as present in the dried samples applied to fresh gauze in a volume of 0.1. Fresh treated patches were applied each day. Pairs of rats, 2/sex, were sacrificed on days 4, 14, 28 and 42 for pathology and determination of PFOS level in liver and PFOS, PFHS, and M570 levels in serum (selected time points). Rats sacrificed on day 42 were treated for the first 28 days followed by a 14-day period without treatment.

Repeated dermal contact under semi-occlusive conditions with all but one of the test samples resulted in consistently quantifiable serum and liver concentrations of PFOS that exceeded levels in control rats exposed to gauze patches moistened only with saline. The exception was a mill-applied carpet protector. With that product, only one male exposed to the liquid product for 28 days, and one female exposed to previously treated and dried gauze patches for 28 days showed liver PFOS levels at close to the level of quantitation (LOQ) of 10 ppb. All other samples were below the LOQ. M570 was detected at consistently quantifiable levels in serum from exposure to only one of the product samples. This sample also yielded trace levels (about 1 ppb) of PFHS in serum of rats exposed daily for 28 days. PFHS was also detected at higher levels (up to about 1 ppm) in the serum of animals exposed to the product that also yielded the highest liver (up to 7-10 ppm) and serum (up to 1 ppm) levels of PFOS.

3M will submit the final report to EPA when available. Because of differences in the nature of the treated fabric (cotton gauze v. carpet or upholstery) and conditions of dermal contact, these data can not be used quantitatively to determine human exposures to PFOS, PFHS and N-MeFOSE from contact with treated materials. These data do, however, indicate that dermal absorption of PFOS, PFHS and N-MeFOSE from contact with these materials is possible. As indicated above, 3M discontinued the manufacture of these products for commercial sale as part of 3M's phase out of its perfluorooctanyl chemistries.

Please contact Dr. John L. Butenhoff (651-733-1962) if you have any questions or if we can provide additional information.

Sincerely,

Katherine E. Reed
Staff Vice President, Environmental Technology and Safety Services

Enclosure

AR226-1154

SUMMARY

December 4, 2002

Compositions of Tested Products, Dermal Contact Absorption/Toxicology Study (PFOS, PFHS, M570)

T-7086.2(Liquid), T-7336(dried)	Component Name	CAS#	Min %W	Max %W	Trade Secret
	WATER	7732-18-5	85.33	85.33	
	ISOBUTANE	75-28-5	7.0	7.0	
	1-(2-BUTOXYPROPOXY)-2-PROPANOL	29911-28-2	2.52	2.52	
	MIDE				YES
	1-METHOXY-2-PROPANOL	107-98-2	0.86	0.86	
	ETHYL ALCOHOL	64-17-5	0.837	0.837	
	MIDE				YES
	FLUOROCHEMICAL ALCOHOL	24448-09-7	0.0	0.0	YES
	METHYL ISOBUTYL KETONE	108-10-1	0.0	1.0	YES
	SODIUM NITRITE	7632-00-0	0.186	0.186	
	MORPHOLINE	110-91-8	0.186	0.186	
	AMMONIUM HYDROXIDE	1336-21-6	0.0384	0.0384	
	METHYL ALCOHOL	67-55-1	0.0326	0.0326	
	2-IMIDAZOLINE-1-ETHANOL, 2-UNDECYL-	136-99-2	0.015	0.015	
	2-methoxy-1-propanol	1589-47-5	0.014	0.014	
	SOLVENT NAPHTHA (PETROLEUM), LIGHT ALIPHATIC	64742-89-8	0.0086	0.0086	
	HEPTANE	142-82-5	0.0	0.0093	
	ETHYL ACETATE	141-78-6	0.0	0.0837	
	Fluorochemical Alcohol	24448-09-7	0.0	0.0375	
	RESIDUAL ORGANIC FLUORO-CHEMICALS		0.0	0.625	
	METHYL ISOBUTYL KETONE	108-10-1	0.0	0.0093	

BEST COPY AVAILABLE

T-7340(Liquid), T-7341(dried)	Component Name	CAS#	Min	Max	Trade Secret
	WATER	7732-18-5	74.0	74.0	
					YES
					YES
					YES
	1,2-DIBROMO-2,4-DICYANO BUTANE	35691-65-7	0.05	0.05	
	N-METHYL PERFLUOROOCTANESULFONAMIDE	31506-32-8	0.0	0.03	
	LINEAR N-ETHYL PERFLUOROOCTANESULFONAMIDE	4151-50-2	0.0	0.02	
	Fluorochemical Alcohol	24448-09-7	0.0	0.2	
	METHYL ISOBUTYL KETONE	108-10-1	0.0	1.0	
	RESIDUAL ORGANIC FLUORO CHEMICALS		0.0	1.0	

T-7080.2(Liquid), T-7089(dried)	Component Name	CAS#	Min	Max	Trade Secret
	ISOPROPYL ALCOHOL	67-63-0	57.25	57.25	
	BUTANE	106-97-8	35.31	35.31	
	PROPANE	74-98-6	5.05	5.05	
	METHYL ISOBUTYL KETONE	108-10-1	1.46	1.46	
					YES
	DI-N-OCTYL ADIPATE	123-79-5	0.1	0.5	

BEST COPY AVAILABLE

T-7342(Liquid), T-7343(dried)	Component Name	CAS#	Min	Max	Trade Secret
	WATER	7732-18-5	79.0	79.0	
					YES
	BORIC ACID	10043-35-3	0.62	0.62	
	TRISODIUM N-(HYDROXYETHYL)ETHYLENEDIAMINETRIACETATE	139-89-9	0.5	0.5	
	SODIUM MONO-C10-16-ALKYL SULFATES	68585-47-7	0.32	0.32	
	RESIDUAL ORGANIC FLUORO-CHEMICALS		0.2	0.2	
	FRAGRANCE		0.07	0.07	
	DISODIUM HYDROXYETHYLETHYLENEDIAMINE DIACETATE	62099-15-4	0.026	0.026	
	SODIUM GLYCOLATE	2836-32-0	0.026	0.026	
	Alcohols, C10-16	67762-41-8	0.014	0.014	
	TRISODIUM NITRILOTRIACETATE	5064-31-3	0.013	0.013	
	SODIUM HYDROXIDE	1310-73-2	0.013	0.013	
	SODIUM SULFATE	7757-82-6	0.011	0.011	
	SODIUM CHLORIDE	7647-14-5	0.0055	0.0055	
	2-BUTOXYETHANOL	111-76-2	11.64	13.55	
	ISOPROPYL ALCOHOL	67-63-0	1.71	2.62	