

AROCLORS

Acute Oral (Rat) and Skin (Female Rabbit) Toxicity

Compound	Oral LD ₅₀ Rats (mg/kg)	Survival time (days)	Skin MLD Rabbits (F) (mg/kg)	Survival time (days)
<u>1221</u> (undil.)	3980 (3300 - 4775)	1 - 3	2000 - 3160	5 - 7
<u>1232</u> (undil.)	4470 (3490 - 5720)	1 - 4	1250 - 2000	4-9
<u>1242</u> 1953 - (undil.)	5730 (5445 - 6000)	1 - 6		
1962 - (undil.)	8650 (7610 - 9775)	1- 3	794 - 1260	4 - 10
<u>1248</u> (undil.)	12,500 (11,000 - 14,250)	1 - 4	794 - 1260	5 - 8
<u>1254</u> (50% corn oil)	10,000 (8,600 - 11,600)	2 - 6	1260 - 2000	4 - 8
<u>1260</u> (50% corn oil)	10,000 (8,600 - 11,600)	2 - 5	1260 - 2000	7 - 8
<u>262</u> (50% corn oil)	11,300 (9,940 - 12,800)	1 - 6	1260 - 3160	5 - 8
<u>1268</u> (50% corn oil)	10,900 (9,590 - 12,320)	2 - 5	2510	
<u>4465</u> (50% corn oil)	16,000 (13,760 - 18,560)	2 - 6	2000 - 3160	9 - 21
<u>5442</u> (50% corn oil)	10,600 (9,110 - 12,300)	2 - 4	1260 - 2000	11 - 19
<u>5460</u> (33 1/3% in corn oil)	ca 20,000	-	7940	-

MUNS 073405