

Company work history records were used to classify employee exposures by work area assignment (Bond et al, 1985; 1986) as well as by potential for exposure to 11 chemical agents chosen because they had been shown to produce cancer of the liver or biliary passages under experimental conditions in animals (Chu et al, 1984; Haseman et al, 1984; NTP, 1985; Wilbourn et al, 1986). Visits to the Medical Department by the cases were reviewed by a nurse co-investigator (EAM) to ascertain a history of excessive alcohol use or hepatitis, as they are established risk factors for primary liver cancer (Falk, 1982). This exercise was of limited usefulness, however, owing to: (1) the unavailability of medical records for those subjects who terminated their employment prior to 1947 (N = 24) and (2) a lack of information concerning any post-employment illnesses among the workers.

Risk ratios were computed adjusted for birthyear using the Mantel-Haenszel (1959) procedure. Analyses which additionally controlled for period of hire showed similar results to those which controlled only for birthyear and are not presented. Likewise, analyses of the primary liver cancers separate from the gall bladder/bile duct cancers did not reveal additional information and are not shown. Ninety five per cent test-based confidence intervals were calculated about the risk ratio estimates (Miettinen, 1976).

RESULTS

Analyses by work area assignments (Table I) showed the cases were significantly less likely to have held assignments in electrochemicals or organic chemical

production, but were more likely to have held assignments in office maintenance (RR = 3.6; 95% CI = 1.4-9.1). A positive association with assignment to polyvinyl-polyvinylidene chloride production did not reach statistical significance (RR = 2.0; 95% CI = 0.9-4.6).

There were no statistically significant trends apparent when analyses were made of duration of assignment within work areas; however, the odds ratio for an assignment of five or more years to polyvinyl-polyvinylidene chloride production was 3.7 (95% CI = 0.9-14.6) based on the contribution of two cases.

Assignment to polyvinyl-polyvinylidene chloride production areas was explored further in Figure 1 via a plot of the "attributable fraction" (Goldsmith et al, 1980) for each year during the period 1940-1980. The peak "attributable fraction" occurred in 1952, a year in which four of the seven cases assigned to these areas were all employed, no two in the same plant, however.

None of the chemical agent comparisons revealed statistically significant differences (Table II). The highest risk ratio (RR = 2.2; 95% CI = 0.9-5.7) was for presumed exposure to vinyl chloride based on the contribution of five cases. There were no statistically significant trends for analysis by duration of exposure to any of the 11 agents, although in many instances there were too few exposed cases to permit meaningful conclusions. Likewise, there were no statistically significant trends apparent when analyses were made by estimated dose of exposure.

DISCUSSION

The present investigation revealed a statistically significant positive association with assignment to office maintenance, and some marginal associations with polyvinyl-polyvinylidene chloride production. The association with an assignment in office maintenance is curious, since this type of work is not generally considered hazardous. In fact, it was a common practice to assign to that department any workers who were not considered medically fit to work in more strenuous jobs. Company medical records were available to review for four of the five cases assigned to office maintenance, and revealed that two of the men had been hospitalized for chronic alcoholism suggesting that this finding may have resulted from confounding.

High levels of vinyl chloride exposure have produced a specific hazard of angiosarcoma of the liver in some work forces (Doll, 1988). In the present study, five of the seven cases of liver and biliary tract cancer who held assignments in the polyvinyl-polyvinylidene work area were considered to have had the opportunity for exposure to vinyl chloride monomer. There was no evidence that any of the five cases experienced angiosarcoma, however, pathology reports and/or medical records were available to confirm the diagnosis for only two of them. Table III presents some descriptive information concerning these cases. By virtue of the time period in which they worked and the nature of their jobs, each of the men was considered to have had the opportunity for high levels of exposure to vinyl chloride monomer, with peak exposures probably exceeding 1000 ppm. Although two of the

cases had spent more than 20 years each in areas of potential vinyl chloride monomer exposure, the other three had very brief exposures on the order of one to several months.

Of interest, none of the other chemical agents suspected a priori as potential risk factors was shown to be statistically significantly associated with liver or biliary tract cancer. It may be that some of the agents are weak carcinogens, but this study could not explore that hypothesis. An examination of the upper limits of the 95% confidence intervals about the risk ratios presented in Table II shows that the study apparently had sufficient power to rule-out modest increases in risk for several chemicals (i.e., carbon tetrachloride and ethylene dichloride), but only relatively large increases in risk could be excluded for others.

Some misclassification of exposure may have occurred; however, the availability of industrial hygiene data from the distant past for many of the plants at this production site should have reduced this possibility considerably. If misclassification did occur, and was non-differential, the tendency would be to diminish the strength of the associations.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge technical assistance from Tom Bradley, Wendy Dahar, Mike Dupuis, Stan Gorgacz, Harold Hoyle, Bruce Houtman, Carl Mueller, Gordon Roush, Dan Schmidt, Belinda Scortichini, Tricia Stirrett, and Mike Strieter. They also thank Diana Diamond for help with preparation of the manuscript.

R&S 100016

REFERENCES

- Bond GG, Flores GG, Shellenberger RJ, Cartmill JB, Fishbeck WA, Cook RR (1986):
Nested case-control study of lung cancer among chemical workers. *American Journal of Epidemiol*, 124; 1:53-66.
- Bond GG, McLaren EA, Cartmill JB, Wymer KT, Sobel W, Lipps TE, Cook RR (1987):
Cause-specific mortality among male chemical workers. *Am J of Industr Med*
12:353-383.
- Bond GG, Shellenberger KJ, Flores GH, et al (1985): A case-control study of renal
cancer mortality at a Texas chemical plant. *Am J Industr Med* 7:123-139.
- Chu KC, Cueto C, Ward JM (1984): Factors in the evaluation of 200 National Cancer
Institute Bioassays. *J Toxicol Environ Hlth* 14:621-639.
- Doll SR (1988): Effects of exposure to vinyl chloride - An assessment of the
evidence. *Scand J Work Environ Health* 14:61-78.
- Falk H (1982): Liver. In Schottenfeld D and Fraumeni JF (editors). *Cancer
Epidemiology and Prevention*. Philadelphia: W. B. Saunders Company, pp.
668-682.
- Goldsmith OF, Smith AH, McMichael AJ (1980): A case-control study of prostate
cancer within a cohort of rubber and tire workers. *J Occup Med* 22:533-541.

Kupper LL, McMichael AJ, Spirtas R (1975): A hybrid epidemiologic study design useful in estimating relative risk. Journal of the American Statistical Association 70; 351:524-528,.

Haseman JK, Crawford DD, Huff JE, Boorman GA, McConnell (1984): Results from 86 two-year carcinogenicity studies conducted by the National Toxicology Program. Journal of Tox and Environ Health 14:621-639.

Miettinen OS (1976): Estimability and estimation in case referent studies. Am J Epidemiol 103:226-235.

National Toxicology Program (NTP) (1985): NTP Technical Report on the toxicology and carcinogenesis studies of dichloromethane in F-344/N rats and B6C3F1 mice (Inhalation studies) NTP-TR-306 (board draft).

Wilbourn J, Haroun L, Haseltine E, et al (1986): Response of experimental animals to human carcinogens: An analysis based upon the IARC Monographs programme. Carcinogenesis 7:1853-1863.

TABLE I

AGE-ADJUSTED RISK RATIOS FOR ANALYSES BY MAJOR AND
MINOR WORK AREAS; STUDY OF LIVER AND BILIARY TRACT
CANCER AMONG CHEMICAL WORKERS, MIDLAND, MICHIGAN,
1940-82

Major/Minor Work Areas	<u>Exposure Frequencies (%)</u>			
	Cases (N = 44)	Referents (N = 1888)	RR	95% CI
Administration	0.0	5.8	-	0.0- 1.5
Manufacturing	68.2	75.7	0.7	0.4- 1.3
Agricultural Chemicals	9.1	26.2	0.4	0.1- 1.0
Herbicides	6.8	17.1	0.5	0.2- 1.6
Insecticides	4.6	9.7	0.6	0.1- 2.4
Inorganic Chemicals	47.7	40.0	1.1	0.6- 2.0
Ammonia	0.0	1.0	-	0.0- 8.4
Brine	6.8	4.6	0.9	0.3- 3.0
Brominated Products	4.6	5.9	0.9	0.2- 3.7
Calcium/Magnesium	9.1	12.5	0.7	0.2- 1.8
Caustic	4.6	1.9	2.1	0.5- 9.5
Cellulose Products	2.3	7.6	0.4	0.1- 2.5
Electrochemicals	0.0	10.2	-	0.0- 0.8
Metal Fabrication	22.7	16.8	1.1	0.6- 2.2
Organic Chemicals	36.4	56.8	0.5	0.3- 0.9
Acrylamide	0.0	2.1	-	0.0- 4.0
Alkyl Benzenes	0.0	3.9	-	0.0- 2.2
Chlorinated Benzenes	0.0	3.6	-	0.0- 2.3
Dyes	6.8	6.1	0.7	0.2- 2.3
Fabricated Plastics	6.8	16.5	0.6	0.2- 1.9
Glycols	0.0	1.8	-	0.0- 4.7
Halogenated	4.6	7.7	0.4	0.1- 1.6
Hydrocarbons				
Ion Exchange Resins	2.3	2.0	1.2	0.2- 8.7
Latex	2.3	2.7	0.8	0.1- 6.2
Olefins	4.6	2.5	1.5	0.3- 6.7
Phenol Derivatives	6.8	13.0	0.4	0.1- 1.4
Polystyrene	6.8	15.0	0.5	0.1- 2.0
Polyvinyl-	15.9	10.1	2.0	0.9- 4.6
Polyvinylidene Chloride				
Unknown Manufacturing	0.0	1.0	-	0.0- 8.4
Research	15.9	22.8	0.8	0.3- 1.8
Service	65.9	59.8	1.1	0.6- 2.0
Animal Handling	0.0	0.6	-	0.0-14.0
Boiler Shop	4.6	3.4	1.0	0.2- 4.1

R&S 100019

Continued on next page

TABLE I CONTINUED

AGE-ADJUSTED RISK RATIOS FOR ANALYSES BY MAJOR AND
MINOR WORK AREAS; STUDY OF LIVER AND BILIARY TRACT
CANCER AMONG CHEMICAL WORKERS, MIDLAND, MICHIGAN,
1940-82

Major/Minor Work Areas	<u>Exposure Frequencies (%)</u>			
	Cases (N = 44)	Referents (N = 1888)	RR	95% CI
Service (continued)				
Construction	13.6	13.6	0.7	0.3- 1.7
Electric Shop	2.3	2.8	0.9	0.1- 6.4
Engineering	2.3	1.6	1.7	0.2-12.5
Garage	0.0	2.5	-	0.0- 3.4
Lead Shop	0.0	0.3	-	0.0-28.0
Lubrication	0.0	0.4	-	0.0-21.0
Machine Shop	4.6	4.2	0.9	0.2- 3.5
Maintenance	4.6	8.6	0.4	0.1- 1.6
Material Handling	11.4	8.1	1.4	0.5- 3.7
Office Maintenance	11.4	3.3	3.6	1.4- 9.1
Paint Shop	4.6	7.9	0.6	0.1- 2.5
Pipe Shop	9.1	10.3	0.8	0.3- 2.3
Plant Protection	9.1	5.6	1.1	0.4- 3.2
Power	2.3	7.1	0.2	0.0- 1.3
Road and Yard Mtc.	4.6	2.0	2.2	0.5- 9.6
Shops-Steel & Instrument	6.9	2.3	2.2	0.7- 7.5
Waste Management	0.0	1.2	-	0.0- 7.0
Well Services	0.0	0.1	-	0.0-83.8

Workers may have been assigned to more than one work area during
their employment and thus may be counted more than once in the
Table.

R&S 100020

TABLE II

AGE-ADJUSTED RISK RATIOS FOR ANALYSES BY SELECTED
CHEMICAL AGENT EXPOSURES; STUDY OF LIVER AND BILIARY
TRACT CANCER AMONG CHEMICAL WORKERS, MIDLAND,
MICHIGAN, 1940-82

Agent	<u>Exposure Frequencies (%)</u>			
	Cases (N = 44)	Referents (N = 1888)	RR	95% CI
Carbon Tetrachloride	11.4	20.0	0.5	0.2- 1.4
Chlorodibenzo-(p)-dioxins	2.3	9.1	0.2	0.4- 1.4
Chloroform	4.6	6.7	0.6	0.2- 2.7
Ethylene Dibromide	2.3	7.3	0.4	0.1- 3.0
Ethylene Dichloride	6.8	14.6	0.4	0.1- 1.3
Methylene Chloride	4.6	8.9	0.8	0.2- 3.6
Perchloroethylene	13.6	11.3	1.8	0.8- 4.3
Polychlorinated Biphenyls	0.0	1.2	-	0.0- 7.0
1,1,2-Trichloroethane	0.0	0.5	-	0.0-16.8
2,4,6-Trichlorophenol	0.0	0.9	-	0.0- 9.3
Vinyl Chloride	11.4	7.8	2.2	0.9- 5.8

Workers may have been exposed to more than one agent during their employment and thus may be counted more than once in the Table.

R&S 100021

TABLE III

DESCRIPTIVE DATA ON FIVE CASES PRESUMED TO HAVE BEEN EXPOSED TO
VINYL CHLORIDE MONOMER; STUDY OF LIVER AND BILIARY TRACT CANCER
AMONG CHEMICAL WORKERS, MIDLAND, MICHIGAN, 1940-82

Case No.	Year First Exposed	Year Last Exposed	Duration of Exposure (yrs; mo)	Year of Death	Job Classification	Best Diagnosis (Source)
1	1943	1943	0; <1	1960	Assistant Operator	Adenocarcinoma of common bile duct (DC)
2	1943	1943	0; 3	1964	Janitor	Carcinoma of liver (DC)
3	1950	1970	20; 0	1976	Operator	Hepatocellular carcinoma (PR)
4	1946	1967	20; 8	1980	Operator	Intrahepatic bile duct cancer (MR and PR)
5	1952	1952	0; 2	1972	Handyman/Packager	Cancer of liver (DC)

DC = death certificate


MR = medical record

PR = pathology report

Figure 1. STUDY OF LIVER AND BILIARY TRACT CANCER

Attributable Fraction for Assignment to
Polyvinyl-Polyvinylidene Chloride Areas
in each Year During 1940-1980

R&S 100023



R&S 100024