

FILE NAME: Ferro Engineering (FER)

DATE: 1966 Aug 10

DOC#: FER010

DOCUMENT DESCRIPTION: Newspaper Article - Asbestos Recognized as Cancer Risk

SUNDAY TIMES 10th Aug 1966

Asbestos recognised —as cancer risk—

By Dr Alfred Byrne

REGULATIONS which acknowledge that occupational exposure to asbestos is causing the most important industrial cancer of our time will be introduced next week by Miss Margaret Herbyson, Minister of Social Security.

From August 22 the tumour known as mesothelioma will be scheduled as a "prescribed disease" so that asbestos workers who contract it as a result of their occupation after a certain date will be eligible for industrial benefit. Mesothelioma is an unusual tumour of the pleura—covering the lung—and of similar tissue lining the abdominal cavity, and is invariably fatal. Several steps are being taken towards discovering the full effects of this recently recognised disease so that measures to prevent it can be devised.

A register of all past and present cases occurring in Britain has been set up by the Medical Research Council's Pneumoconiosis Research Unit, directed by Dr J. C. Gibson, in Cardiff. They have added 70 new cases to their original nucleus of 120 since the Sunday Times broke the news of this epidemic on October 21, 1965. One hospital in London and another in Belfast are each admitting an average of one new case a month.

An average of 40 years usually elapses between exposure to asbestos and the appearance of the growth. Exposure for only

two or three months may cause the cancer. Living near an asbestos works predisposes to mesothelioma, and relatives of asbestos workers can also get it through inhaling asbestos fibres given off the clothing of their menfolk.

Such patients were not working with asbestos, so they will not be eligible for compensation under the Industrial Injuries Act of 1947. Incidentally, perhaps, there are also other mesothelioma victims who are not eligible under the new regulation because they worked for a short time in the industry only before the new Act came into force.

There are three types of asbestos used commercially—chrysotile, crocidolite and amosite—whose possible hazards are now being assessed in the laboratory and the field. With Dr Gibson as chairman, the International Union against Cancer is arranging mesothelioma surveys in areas where only one type of fibre is prevalent. Studies in South Africa have already shown that the tumour is more common in the neighbourhood of mines producing crocidolite than amosite. To complete the picture the committee are anxious to see a similar study performed in Canada in relation to mines producing chrysotile.

Asbestos is now used to make gutters, roofing, tiles, brake linings and numerous other things not envisaged when the risk of asbestos was first realised.

Copy to
H. J. Sheets
10/17/66