

velocities (particularly localized drafts) should be avoided since differential cooling of one area of the body may produce surprisingly unpleasant reactions in quite different parts of the body. In a recent experiment¹⁵ it was shown that the application of an ice pack to an area of 60 sq cm on the back of the neck for 15 min caused a drop of 17 F in the skin temperature of the fingers and that this low temperature of the fingers persisted for one hour after the ice pack was removed.

ADAPTATION TO HOT CONDITIONS

It will be observed from Fig. 1 that a zone ranging from about 70 to 80 F, at 45 per cent relative humidity (or from 66 to 74 deg ET) the body is adequately able to maintain equilibrium through control of radiation and convection losses combined, and evaporative loss. This corresponds to the zone over which the largest percentage of persons find optimum comfort. For higher temperatures, up to an upper limit in the neighbor-

TABLE 2. PHYSIOLOGICAL RESPONSES TO HEAT OF MEN AT REST AND AT WORK^a

EFFECTIVE TEMP	ACTUAL CLOTHED TEMP (Deg Fahr)	MEN AT REST			MEN AT WORK 90,000 FT-LB OF WORK PER HOUR			
		Rise in Rectal Temp (Deg Fahr per Hr)	Increase in Pulse Rate (Beats per Min per Hr)	Approximate Loss in Body Weight by Perspiration (Lb per Hr)	Total Work Accomplished (Ft-Lb)	Rise in Body Temp (Deg Fahr per Hr)	Increase in Pulse Rate (Beats per Min per Hr)	Approximate Loss in Body Wt by Perspiration (Lb per Hr)
60	-----	0.0	0	0.2	225,000	0.0	6	0.5
70	-----	0.0	0	0.3	225,000	0.1	7	0.6
80	96.1	0.0	0	0.3	209,000	0.3	11	0.8
85	96.6	0.1	1	0.4	190,000	0.6	17	1.1
90	97.0	0.3	4	0.5	153,000	1.2	31	1.5
95	97.6	0.9	15	0.9	102,000	2.3	61	2.0
100	99.6	2.2	40	1.7	67,000	4.0	103 ^b	2.7
105	104.7	4.0	83	2.7	49,000	6.0 ^b	158 ^b	3.5 ^b
110	-----	5.9 ^b	137 ^b	4.0 ^b	37,000	8.5 ^b	237 ^b	4.4 ^b

^aData by A.S.H.V.E. Research Laboratory.

^bComputed value from exposures lasting less than one hour.

hood of 100 F, at 45 per cent relative humidity (or approximately 87 deg ET) control is had through availability of perspiration on the body surface for evaporation. While a fair degree of temperature equilibrium is maintained over this range it is nevertheless had with considerable discomfort.

Studies at the *John B. Pierce Laboratory of Hygiene*¹⁶ have indicated the relation between discomfort and the degree of wetting of the body surface by perspiration for lightly clothed subjects in a semi-reclining position, and the investigators there have designated this as the *zone of evaporative regulation*. Work of the A.S.H.V.E. Research Laboratory¹⁷

¹⁵The Relative Influence of Radiation and Convection Upon the Temperature Regulation of the Clothed Body, by C. E. A. Winslow, L. P. Herrington and A. P. Gage (*American Journal of Physiology*, Vol. 124, October, 1938, p. 51).

¹⁶Relations Between Atmospheric Conditions, Physiological Reactions, and Sensations of Pleasantness, by C. E. A. Winslow, L. P. Herrington and A. P. Gage (*American Journal of Hygiene*, Vol. 26, July, 1937, p. 102). The Reactions of the Clothed Human Body to Variations in Atmospheric Humidity, by C. E. A. Winslow, L. P. Herrington and A. P. Gage (*American Journal of Physiology*, Vol. 124, December, 1938, p. 692).

¹⁷Loc. Cit. Note 11.

over the past two decades has made available data on the relation between sensible perspiration and the atmospheric environment for normal persons at rest and at work.

For colder conditions below 70 F, with 45 per cent relative humidity (or about 66 deg ET) thermal equilibrium is maintained, first, by the amount of clothing worn; second, and to a smaller extent, by limiting availability of heat at the surface by decreasing peripheral blood circulation, which results in a drop in skin temperature; and third, by an increase in the metabolic rate. Here again, while thermal equilibrium is fairly well maintained, any drop in the skin temperature is accompanied by a certain degree of discomfort.

Studies at the A.S.H.V.E. Research Laboratory¹⁸ and elsewhere¹⁹ during the past two decades have made available a mass of information dealing with the physiological effects of hot atmospheres on workers and means to alleviate the distress and hazards associated therewith. This interest has been termed *air conditioning in industry*, or *the effects of hot atmospheres in industrial hygiene*, and is a growing factor in air conditioning applications. Table 2 gives some of the physiological responses of men at rest and at work, to hot environments. Recent physiological studies²⁰ indicate that frequent and continued exposure of workers to hot environments results in not only violent but subtle physiological derangement, affecting the leucocyte count of the blood and other factors dealing with man's mechanism of defense against infection.

Another of the deleterious effects of high temperatures is that the blood is diverted from the internal organs to the surface capillaries, in order to serve in the process of cooling. This affects the stomach, heart, lungs and other vital organs, and it is suggested that the feeling of lassitude and discomfort experienced is due in part to the anaemic condition of the brain. The stomach loses some of its power to act upon the food, owing to a diminished secretion of gastric juice, and there is a corresponding loss in the antiseptic and antifermentative action which favors the growth of bacteria in the intestinal tract²¹. These are considered to be the potent

¹⁸A.S.H.V.E. RESEARCH REPORT No. 654—Some Physiological Reactions to High Temperatures and Humidities, by W. J. McConnell and F. C. Houghten (A.S.H.V.E. TRANSACTIONS, Vol. 29, 1923, p. 129). A.S.H.V.E. RESEARCH REPORT No. 672—Further Study of Physiological Reactions, by W. J. McConnell, F. C. Houghten and F. M. Phillips (A.S.H.V.E. TRANSACTIONS, Vol. 29, 1923, p. 353). A.S.H.V.E. RESEARCH REPORT No. 690—Air Motion, High Temperatures and Various Humidities—Reactions on Human Beings, by W. J. McConnell, F. C. Houghten and C. P. Yaglou (A.S.H.V.E. TRANSACTIONS, Vol. 30, 1924, p. 167). A.S.H.V.E. RESEARCH REPORT No. 718—Work Tests Conducted in Atmospheres of High Temperatures and Various Humidities in Still and Moving Air, by W. J. McConnell and C. P. Yaglou (A.S.H.V.E. TRANSACTIONS, Vol. 31, 1925, p. 101). A.S.H.V.E. RESEARCH REPORT No. 719—Basal Metabolism Before and After Exposure to High Temperatures and Various Humidities, by W. J. McConnell, C. P. Yaglou and W. B. Fulton (A.S.H.V.E. TRANSACTIONS, Vol. 31, 1925, p. 123). A.S.H.V.E. RESEARCH REPORT No. 908—Heat and Moisture Losses from Men at Work and Application to Air Conditioning Problems, by F. C. Houghten, W. W. Teague, W. E. Miller and W. P. Yant (A.S.H.V.E. TRANSACTIONS, Vol. 37, 1931, p. 541). A.S.H.V.E. RESEARCH REPORT No. 1106—Air Conditioning in Industry—Physiological Reactions of Individual Workers to High Effective Temperatures, by W. L. Fleisher, A. E. Stacey, Jr., F. C. Houghten and M. B. Ferderber (A.S.H.V.E. TRANSACTIONS, Vol. 45, 1939, p. 59). A.S.H.V.E. RESEARCH REPORT No. 1153—Seasonal Variation in Reactions to Hot Atmospheres, by F. C. Houghten, A. A. Rosenberg and M. B. Ferderber (A.S.H.V.E. TRANSACTIONS, Vol. 46, 1940). Physiologic Effects of Hot Atmospheres, by F. C. Houghten, M. B. Ferderber and A. A. Rosenberg (*Industrial Medicine*, January, 1940, p. 7).

¹⁹A.S.H.V.E. RESEARCH REPORT No. 1151—The Peripheral Type of Circulatory Failure in Experimental Heat Exhaustion, by R. W. Keeton, F. K. Hick, Nathaniel Glickman and M. M. Montgomery (A.S.H.V.E. TRANSACTIONS, Vol. 46, 1940).

²⁰A.S.H.V.E. RESEARCH REPORT No. 1153—Seasonal Variation in Reactions to Hot Atmospheres, by F. C. Houghten, A. A. Rosenberg and M. B. Ferderber (A.S.H.V.E. TRANSACTIONS, Vol. 46, 1940).

²¹Influence of Effective Temperature upon Bactericidal Action of Gastro-Intestinal Tract, by Arnold and Brody (*Proceedings Society Exp. Biol. Med.*, Vol. 24, 1927, p. 832).