

AR226-0314

Acute Ocular Irritation Test

with T-3494

in Albino Rabbits

Experiment No.:

0884EB0011

Conducted At:

Safety Evaluation Laboratory  
Riker Laboratories, Inc.  
St. Paul, Minnesota

Dates Conducted:

January 3, 1984 to January 10, 1984

Conducted By:

D. M. Markoe, Jr. 1/24/84  
D. M. Markoe, Jr., BS Date  
Toxicologist  
Study Director

Reviewed By:

K. D. O'Malley 1/24/84  
K. D. O'Malley, BS Date  
Senior Toxicologist  
Acute Toxicology

dc: M. T. Case  
K. L. Ebbens  
F. D. Griffith  
W. C. McCormick

005529

### Summary

The results of the acute ocular irritation test conducted from January 3, 1984 to January 10, 1984 at Riker Laboratories, Inc., St. Paul, Minnesota indicate that T-3494 is minimally irritating (3.3/110.0) to the eye of the female albino rabbit. Neither corneal opacity nor iritis were produced during the study. Minimal conjunctivitis was noted at the one hour evaluation and subsided by the two day evaluation.

### Introduction

The objective of this study was to determine the acute ocular irritation properties of T-3494 when instilled into the eye of female albino rabbits. This study is not regulated by the Food and Drug Administration's Good Laboratory Practice Regulation of 1978, although the standard operating procedures of this laboratory adhere to the general principles of this regulation. The raw data generated by the Study Director and the final report are stored in the conducting laboratory's archives.

005530

### Method and Results

Young albino rabbits of the New Zealand breed<sup>a</sup> were used to evaluate the ocular irritating properties of the test article. The test method was modeled after that of Draize et al<sup>b</sup>.

The test article was instilled into the conjunctival sac of the right eye of each rabbit according to the treatment procedure presented in Table 1 with the left eye of each animal serving as a control. At each scoring interval, the cornea, iris and palpebral conjunctiva were examined and graded for irritation and injury according to a standard scoring system<sup>b</sup>. The maximum possible score at any one examination and scoring period 110 points, which indicates maximal irritation and damage to all three ocular tissues (cornea, iris, conjunctiva) while a score of zero indicates no irritation (Table 2). In this scoring system, special emphasis is placed upon irritation or damage to the cornea, while less emphasis is placed upon damage to the iris and conjunctiva.

After completion of the test, the scores were analyzed, and a descriptive eye irritation rating was assigned to the test article. The criteria used for assignment of the descriptive rating were the frequency, the extent and the persistence of irritation or damage which occurred to the three ocular tissues (Table 3). The individual results are presented in Table 4.

<sup>a</sup> Hazleton Dutchland, Inc., Denver, PA

<sup>b</sup> Draize: Appraisal of the Safety of Chemicals in Foods, Drugs and Cosmetics (1965).

The rating is arrived at by selecting the maximum mean irritation score at one hour, one, two or three days after instillation. If the rate of dissipation of injury does not meet the requirements defined for the descriptive rating appropriate for a particular numerical score, the descriptive rating is raised by one or more levels. The rating system is presented in Table 3. The protocol, principal personnel involved in the study, composition characteristics and Quality Assurance statement are contained in Appendices I - IV.

005532

Table 1

## Eye Irritation Test - Albino Rabbits

## Treatment Procedure

| Test Article | Number of Animals Evaluated | Form Administered | Quantity of Test Article Administered | Contact Period (seconds) | Volume of Wash (tap water) | Evaluation Time Post Dose Administration |
|--------------|-----------------------------|-------------------|---------------------------------------|--------------------------|----------------------------|------------------------------------------|
| T-3494       | 6                           | solid             | 0.1 gm                                | unlimited                | none                       | 1 Hour, 1, 2, 3 and 7 Days               |

005533

Table 2

## Eye Irritation Test - Albino Rabbits

Scale of Weighted Scores for  
Grading the Severity of Ocular Lesions

| Ocular<br>Tissues | Description                                                                                                                                                                          | Draize<br>Grade    |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|
| Conjunctiva       | <u>Redness (A)</u>                                                                                                                                                                   | 1                  |
|                   | Redness (refers to palpebral conjunctiva only).                                                                                                                                      |                    |
|                   | Vessels definitely injected above normal.                                                                                                                                            | 2                  |
|                   | More diffuse, deeper crimson red, individual vessels not easily discernible.                                                                                                         | 3                  |
|                   | Diffuse beefy red.                                                                                                                                                                   |                    |
|                   | <u>Chemosis (B)</u>                                                                                                                                                                  | 1                  |
|                   | Any swelling above normal (included nictitating membrane).                                                                                                                           | 2                  |
|                   | Obvious swelling with partial eversion of the lids.                                                                                                                                  | 3                  |
|                   | Swelling with lids about half-closed.                                                                                                                                                | 4                  |
|                   | Swelling with lids about half-closed to completely closed.                                                                                                                           |                    |
|                   | <u>Discharge (C)</u>                                                                                                                                                                 | 1                  |
|                   | Any amount different from normal (Does not include small amount observed in inner canthus of normal animals).                                                                        | 2                  |
|                   | Discharge with moistening of the lids and hairs just adjacent to the lids.                                                                                                           | 3                  |
|                   | Discharge with moistening of the lids and hairs and considerable area around eye.                                                                                                    |                    |
|                   | Score (A + B + C) x 2                                                                                                                                                                | Total maximum = 20 |
| Cornea            | <u>Opacity (A)</u>                                                                                                                                                                   |                    |
|                   | Opacity - Degree of density (area which is most dense is taken for reading).                                                                                                         | 1                  |
|                   | Scattered or diffuse area, details of iris clearly visible.                                                                                                                          | 2                  |
|                   | Easily discernible translucent areas, details of iris slightly obscured.                                                                                                             | 3                  |
|                   | Opalescent areas, no details of iris visible, size of pupil barely discernible.                                                                                                      | 4                  |
|                   | Opaque, iris invisible.                                                                                                                                                              |                    |
|                   | <u>Area of Cornea Involved (B)</u>                                                                                                                                                   | 1                  |
|                   | One quarter (or less) but not zero.                                                                                                                                                  | 2                  |
|                   | Greater than one-quarter, but less than one-half.                                                                                                                                    | 3                  |
|                   | Greater than one-half, but less than three-quarters.                                                                                                                                 | 4                  |
|                   | Greater than three-quarters, up to whole area.                                                                                                                                       |                    |
|                   | Score equals A x B x 5                                                                                                                                                               | Total maximum = 80 |
| Iris              | <u>Values (A)</u>                                                                                                                                                                    | 1                  |
|                   | Folds above normal, congestion, swelling, circumcorneal injection (any or all of these or combination of any thereof), iris still reacting to light (sluggish reaction is positive). |                    |
|                   | No reaction to light, hemorrhage, gross destruction (any or all of these).                                                                                                           | 2                  |
|                   | Score equals A x 5                                                                                                                                                                   | Total maximum = 10 |

Note: The maximum total score is the sum of all scores obtained for the  
conjunctiva.

005534

Table 3

## Eye Irritation Test - Albino Rabbits

Classification of Test Materials  
Based on Eye Irritation Properties

| Rating                     | Range         | Definition                                                                                                                                                                                                                                                                                                                                                                       |
|----------------------------|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Non-Irritating             | 0.0 - 0.5     | To maintain this rating, all scores by the one day reading must be zero; otherwise, increase rating one level.                                                                                                                                                                                                                                                                   |
| Practically Non-Irritating | >0.5 - 2.5    | To maintain this rating, all scores by the one day reading must be zero; otherwise, increase rating one level.                                                                                                                                                                                                                                                                   |
| Minimally Irritating       | >2.5 - 15.0   | To maintain this rating, all scores by the three day reading must be zero; otherwise, increase rating one level.                                                                                                                                                                                                                                                                 |
| Mildly Irritating          | >15.0 - 25.0  | To maintain this rating, all scores by the seven day reading must be zero; otherwise, increase rating one level.                                                                                                                                                                                                                                                                 |
| Moderately Irritating      | >25.0 - 50.0  | To maintain this rating, scores by seven days must be $\leq 10$ for 60% or more of the animals. Also, mean seven day score must be $\leq 20$ . If seven day mean score $\leq 20$ but $< 60\%$ of animals show scores $< 10$ , then no animal among those showing scores $> 10$ can exceed a score of 30 if rating is to be maintained; otherwise raise rating one level.         |
| Severely Irritating        | >50.0 - 80.0  | To maintain this rating, scores by seven days must be $\leq 30$ for 60% or more of the animals. Also, mean seven day score must be $\leq 40$ . If seven day mean score $\leq 40$ but $< 60\%$ of the animals show scores $\leq 30$ , then no animal among those showing scores $> 30$ can exceed a score of 60 if rating is to be maintained; otherwise, raise rating one level. |
| Extremely Irritating       | >80.0 - 110.0 |                                                                                                                                                                                                                                                                                                                                                                                  |

BEST COPY AVAILABLE

TABLE 4

## EYE IRRITATION TEST - ALBINO RABBITS

with T-3494

| Tissue       | Examination Period | RESULTS<br>ANIMAL NUMBERS |           |           |           |           |           | Means |
|--------------|--------------------|---------------------------|-----------|-----------|-----------|-----------|-----------|-------|
|              |                    | 3B2037                    | 3B2036    | 3B2035    | 3B2040    | 3B2039    | 3B2038    |       |
| Cornea (D-A) | 1 Hour             | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Iris         |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Conjunctiva  |                    | 4 (2-0-0)                 | 4 (2-0-0) | 4 (2-0-0) | 2 (1-0-0) | 4 (2-0-0) | 2 (1-0-0) | 3.3   |
| (RSD)        |                    | 4                         | 4         | 4         | 2         | 4         | 2         | 3.3   |
| Cornea (D-A) | 1 Day              | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Iris         |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Conjunctiva  |                    | 0                         | 2 (1-0-0) | 2 (1-0-0) | 0         | 0         | 0         | 0.7   |
| (RSD)        |                    | 0                         | 2         | 2         | 0         | 0         | 0         | 0.7   |
| Cornea (D-A) | 2 Days             | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Iris         |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Conjunctiva  |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| (RSD)        |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Cornea (D-A) | 3 Days             | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Iris         |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Conjunctiva  |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| (RSD)        |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Cornea (D-A) | 7 Days             | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Iris         |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| Conjunctiva  |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |
| (RSD)        |                    | 0                         | 0         | 0         | 0         | 0         | 0         | 0.0   |

005536

5536

**BEST COPY AVAILABLE**

Riker Experiment No.: \_\_\_\_\_

8

APPENDIX I  
PROTOCOL

TEST: Acute Ocular Irritation Test

SPONSOR: 3M Commercial Chemicals

CONDUCTED BY: Safety Evaluation Laboratory, Riker Laboratories, Inc., St. Paul, Minnesota

TEST ARTICLE: T-3494

CONTROL ARTICLE: None

PROPOSED STARTING/COMPLETION DATE OF TEST: 1/24 - 4/74

TEST SYSTEM: Female New Zealand White Albino Rabbits

SOURCE: Dr. H. J. Auer, Riker Laboratories

OBJECTIVE: The objective of this test will be to determine the irritation potential of the test article to the ocular tissues (cornea, iris and conjunctiva) of 1 albino rabbits. Rabbits were selected as the test system for their sensitivity to irritants, historical use, ease of handling and general availability.

METHOD: The animals will be housed in standard wire-mesh cages in temperature and humidity controlled rooms with food<sup>a</sup> and water offered *ad libitum*. Each animal will be assigned a numbered ear tag which will correspond to a card affixed to the outside of the cage. The test article will be instilled into the conjunctival sac of the right eye at a dose of 0.1 g with the contralateral eye of each animal serving as a control. At 1 hours and 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 (additional scoring intervals may be added to further characterize the ocular reactions), the tissues will be examined and graded for irritation and injury according to a standard scoring system of 1 to 4 et al.<sup>b</sup>. After completion of the test, the scores will be analyzed, and a descriptive eye irritation report assigned to the test article. Eye examinations may be carried out with the aid of sodium fluorescein. If deemed necessary by the study director, washed eye procedures entailing a 5 and 30 second contact period with a 5 ml wash over a 5 minute period will be conducted on 3 animals per procedure. All raw data generated by the study director and the final report will be stored in the Riker Laboratories' Archive, St. Paul, Minnesota.

<sup>a</sup> Purina Rabbit Chow, Ralston-Purina, St. Louis, Missouri

<sup>b</sup> Draize: Appraisal of the Safety of Chemicals in Foods, Drugs, and Cosmetics (1965)  
Published by the Editorial Committee of The Association of Food and Drug Officials  
of the United States.

Sponsor

Date

Study Director

005537

APPENDIX IIPrincipal Participating Personnel Involved in the Study

| <u>Name</u>           | <u>Function</u>                         |
|-----------------------|-----------------------------------------|
| D. M. Markoe, Jr., BS | Toxicologist<br>Study Director          |
| K. L. Ebbens, BS      | Supervisor<br>Toxicology Testing        |
| K. D. O'Malley, BS    | Senior Toxicologist<br>Acute Toxicology |
| G. C. Pecore          | Supervisor<br>Animal Laboratory         |

005538

APPENDIX IIIComposition Characteristics

This study is not regulated by the Good Laboratory Practice Act of 1978 and therefore information pertaining to composition characteristics is not applicable for inclusion in this study.

005539

APPENDIX IVQuality Assurance Statement

This study is not officially regulated by the Good Laboratory Practice Regulation of 1978, and therefore a statement signed and prepared by the Compliance Audit department is not applicable.

The standard operating procedures of this laboratory does adhere to the general principles of this regulation. The Compliance Audit department does inspect different significant phases for studies underway in the Acute Toxicology Laboratory on a recurring cycle, and the facilities are examined on a three month schedule. In addition a select number of Research & Development studies are routinely picked at random from the Archives by the Compliance Audit department for review.

005540