

Urban air and tobacco smoke in benzene exposure in a cohort of traffic policemen

Roberto Bono^{a,*}, Deborah Traversi^a, Luciano Maestri^b, Tiziana Schilirò^a,
Sergio Ghittori^b, Claudio Baiocchi^c, Giorgio Gilli^a

^a Department of Public Health and Microbiology, University of Torino, Italy

^b Salvatore Maugeri Foundation, Laboratory of Environmental Hygiene and Industrial, Toxicology, Pavia, Italy

^c Department of Analytical Chemistry, University of Torino, Italy

Available online 12 April 2005

Abstract

Benzene (B) is a typical micro-pollutant present in air, especially urban air. In this study a possible correlation between personal benzene exposure and S-phenylmercapturic acid (S-PMA) as a biomarker of internal dose was evaluated in a cohort of traffic policemen. The results confirm that S-PMA is significantly correlated to benzene measured in personal air. B and S-PMA were analyzed considering seasonality, work quarters, time spent indoors, outdoors, and directing traffic, but no significant differences were recorded.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Benzene; Traffic policemen; Biomarkers; Urinary S-phenylmercapturic acid

1. Introduction

Benzene (B) is an ubiquitous micro-pollutant present primarily in air (99%). In particular, active smoke represents the most important source of B exposure for smokers, while, for nonsmokers, B exposure is principally due to outdoor ambient air, where B pollution is caused by auto exhaust or gasoline vapor emissions [1] and to indoor air where the presence of B is largely due to environmental tobacco smoke. B air pollution is a very important topic for environmental health due to its carcinogenic properties by all routes of exposure.

The aim of this study is to search for a possible correlation between personal passive exposure to B and levels of urinary S-phenylmercapturic acid (S-PMA) in a cohort of traffic policemen that, for their specific working functions, spend their time indoors and outdoors. Due to the presence of B in tobacco smoke, urinary cotinine as a confounding factor was also investigated in all the subjects.

2. Materials and methods

The present study enrolled 206 volunteers among traffic policemen working in Turin, a city with 900,000 inhabitants located in north-western Italy. The enrolled

* Corresponding author.

E-mail address: roberto.bono@unito.it (R. Bono).

subjects were also administered a questionnaire with questions including their smoking habits and the time spent during work in three kinds of environment: offices (indoor), in the streets (outdoor) and directing traffic (outdoor, close to high traffic volume).

The sampling period lasted from March 2002 to April 2003. Each Thursday, six policemen wore a personal diffusive sampler able to measure air B exposures for a whole working shift. At the end of the working shift a questionnaire was administered and a urine sample was collected to measure S-PMA and cotinine as biomarkers of benzene and of tobacco smoke respectively.

B was analyzed using a gas chromatograph (GC) Carlo Erba 5300 Mega Series equipped with a flame ionization detector (FID) and capillary column DB-624 30 m $\times$ 0.318 mm i.d., film 1.8 μ m. Each sample was eluted using ultrapure carbon disulfide (CS2 99.9% low benzene content, Aldrich 34.227-0) (3 ml for air samples, 2 ml for personal monitor samples). A calibration curve was prepared from known concentrations of benzene, toluene, and xylenes. The GC thermal program was 45 °C for 4 min, increase of 10 °C/min, from 45 to 145 °C, then 145 °C for 2 min [2,3].

Urinary S-PMA was measured by a method already described elsewhere [4] and slightly modified in order to improve sensitivity. Briefly, samples were purified by a two-step protocol with reversed-phase (C18-EC) and with strong anionic exchange (SAX) solid-phase extraction cartridges. Then they were deacetylated by means of porcine acylase. After centrifugal ultrafiltration with disposable units (cut-off 10,000), the de-proteinized samples were derivatized for 5 min with the *o*-phthaldialdehyde/2-mercaptoethanol (OPA-MCE) reagent and the fluorescent derivatives were separated by high-performance liquid chromatography (HPLC) on a reversed phase column. The limit of detection of the method was 0.2 μ g/l of S-PMA in urine.

Urine samples were stored at -70 °C until cotinine analysis was performed, within 6 months. In a centrifuge tube, 1 ml of NaOH 5 M, 8.5 g NaCl and 5 ml of chloroform were added to 25 ml of urine. The methodology for cotinine analysis and quality control was published previously [5]. Instrumental analysis was performed using a gas chromatograph (GC) equipped with a flame ionization-nitrogen-selective detector (NPD) using an oven programmed temperature rising

from 100 to 250 °C. The external quality control for the GC technique consisted in a re-analysis for 20 times of a cotinine standard solution added to a pool of nonsmokers' urine samples [6]. Results show a coefficient of variation (CV) of 4.3%, a detection limit below 1 ng/ml, and recovery around 100%. Pyridine was used as an internal standard for quality control.

3. Results

The sites of work of the 206 policemen covered the entire municipal area of Turin involving both residential and traffic-congested areas. Subjects declared through a questionnaire their age (years 38 ± 7), gender (male 60%), length of service (years 8.7 ± 8). A general Pearson correlation analysis between number of cigarettes smoked per day and urinary cotinine showed that urinary cotinine is a sensitive and specific internal dose marker of tobacco smoke exposure (Fig. 1). Furthermore, the cotinine level in smokers and nonsmokers were 671 and 25 ng/ml, respectively ($p < 0.0001$).

A general statistical distribution of B and S-PMA values are shown in Table 1. Linear regression and Pearson's coefficient r of the two parameters are reported in Fig. 2.




Fig. 1. Regression analysis between number of cigarettes smoked per day and urinary cotinine for all the 206 subjects.

Table 1
General statistical distribution of B and S-PMA values

	N	Min	Max	Mean	S.D.
Benzene (μ g/m ³)	206	0.1	65.0	8.8	8.1
S-PMA (μ g/g of creatinine)	206	0.3	8.1	1.5	1.0




Fig. 2. Regression analysis between personal benzene and S-PMA for all the 206 subjects.

Table 2
Statistical comparison of B and S-PMA values sub-grouped by smoking habit

	<i>N</i>	Mean	S.D.	<i>t</i> -Test
Benzene (μg/m ³)				
Nonsmokers	154	7.6	7.5	0.0001
Smokers	52	12.7	8.6	
S-PMA (μg/g of creatinine)				
Nonsmokers	154	1.2	0.6	0.0001
Smokers	52	2.3	1.6	

Taking smoking habits into account, *t*-test underlines a significantly higher level of B and S-PMA for smokers (Table 2). By means of ANOVA multivariate analysis, mean values of B and S-PMA were analyzed considering seasonality, work quarters, time spent indoors, outdoors, and directing traffic, but no statistically significant differences were recorded.

4. Discussion and conclusions

The present study confirms that the level of S-PMA is significantly correlated to benzene in personal air. The traffic policemen are a very useful human model able to represent different environments where people live and work. In particular, no differences were recorded considering either indoor or outdoor air exposure, or the direct exposure of traffic policemen to B pollution originating from vehicle exhaust. These data demonstrate, on one hand, the very low level of B contamination in outdoor air and, on the other hand, the not marginal level of B in indoor air (the offices). Fur-

Table 3
Comparison between S-PMA ($\mu\text{g}/\text{g}$ of creatinine) detected in other studies and in this study

	Boogaard and Van Sitter [7]	Melikian et al. [8]	Ghittori et al. [9]	This study
Smokers	3.6	9.1	7.8	2.3
Nonsmokers	1.9	4.8	1.0	1.2

thermore, the important role of tobacco smoke in the synthesis of S-PMA is, at this point, clear.

Several measures were adopted by the European Community and, locally, by Turin municipality to prevent outdoor B exposure, in particular that arising from automobiles. In other studies, urinary S-PMA measured in smokers was higher, probably due to higher B exposure (Table 3). The data obtained in this study show that the subjects analyzed are exposed to low levels of B, as demonstrated by the low excretion of S-PMA in nonsmokers, independently from the working conditions. This study demonstrates also that, for traffic policemen, tobacco smoke is the main source of B exposure, as evidenced by the doubling of S-PMA excretion in smokers.

These data have important implications for both prevention policy at work and against the tobacco smoking habit.

Acknowledgement

Thanks are given to all the traffic policemen of Turin city and in particular to the Commander-in-Chief Dr. Mauro Famigli.

References

- [1] L. Wallace, Environmental exposure to benzene: an update, *Environ. Health Perspect* 104 (1996) 1129–1136.
- [2] R. Bono, E. Scursatone, T. Schilirò, G. Gilli, Ambient air levels and occupational exposure to benzene, toluene and xylenes in Northwestern Italy, *J. Toxicol. Environ. Health* 66 (2003) 519–531.
- [3] R. Bono, E.H. Bugliosi, T. Schilirò, G. Gilli, The Lagrange Street story: the prevention of aromatics air pollution during the last nine years in a European city, *Atmos. Environ.* 35 (2001) S107–S113.
- [4] L. Maestri, S. Ghittori, M. Imbriani, Determination of specific mercapturic acids as an index of exposure to environmental benzene, toluene, and styrene, *Ind. Health* 35 (1997) 489–501.

- [5] R. Bono, R. Russo, W. Arossa, E. Scursatone, G. Gilli, Involuntary exposure to tobacco smoke in adolescents: Urinary cotinine and environmental factors, *Arch. Environ. Health* 51 (1996) 127–131.
- [6] R. Bono, M. Vincenti, T. Schilirò, D. Traversi, C. Pignata, E. Scursatone, G. Dotti, G. Gilli, Cotinine and *N*-(2-hydroxyethyl)valine as markers of passive exposure to tobacco smoke in children, *J. Exposure Anal. Environ. Epidemiol.* 15 (1) (2005) 66–73.
- [7] P.J. Boogaard, N.J. Van Sitter, Biological monitoring of exposure to benzene: a comparison between S-phenylmercapturic acid, trans, trans muconic acid, and phenol, *Occup. Environ. Med.* (1995) 611–620.
- [8] A.A. Melikian, R. O'Connor, A.K. Prahalad, P. Hu, H. Li, M. Kagan, S. Thompson, Determination of the urinary benzene metabolites S-phenylmercapturic acid and trans-, trans-muconic acid by liquid chromatography tandem mass spectrometry, *Carcinogenesis* 4 (1999) 719–726.
- [9] S. Ghittori, M. Imbriani, L. Maestri, E. Capodaglio, A. Cavaleri, Determination of S-phenylmercapturic acid in urine as an indicator of exposure to benzene, *Toxicol. Lett.* 108 (1999) 329–334.