

FILE NAME: Brakes (BRK)

DATE: 2012

DOC#: BRK222

DOCUMENT DESCRIPTION: Journal Article - Asbestos-Related Diseases in Automobile Mechanics

Asbestos-Related Diseases in Automobile Mechanics

JACQUES AMEILLE^{1*}, NICOLE ROSENBERG², MIREILLE MATRAT^{3,4},
ALEXIS DESCATHA¹, DOMINIQUE MOMPOINT⁵, LOUNIS HAMZI⁶,
CATHERINE ATASSI⁷, MANUELA VASILE⁷, ROBERT GARNIER² and
JEAN-CLAUDE PAIRON^{3,4}

¹AP-HP, Unité de pathologie professionnelle, Hôpital Raymond Poincaré, 92380 Garches, France;

²AP-HP, Consultation de pathologie professionnelle, Hôpital Fernand Widal, 75010 Paris, France;

³Centre Hospitalier Intercommunal, Unité de pathologie professionnelle, 94000 Créteil, France;

⁴INSERM, Unité 955, 94000 Créteil, France; ⁵AP-HP, Service de radiologie, Hôpital Raymond Poincaré, 92380 Garches, France; ⁶AP-HP, Service de radiologie, Hôpital Fernand Widal, 75010 Paris, France; ⁷Service de radiologie, Centre Hospitalier Intercommunal, 94000 Créteil, France

Received 24 March 2011; in final form 23 June 2011; published online 28 September 2011

Purpose: Automobile mechanics have been exposed to asbestos in the past, mainly due to the presence of chrysotile asbestos in brakes and clutches. Despite the large number of automobile mechanics, little is known about the non-malignant respiratory diseases observed in this population. The aim of this retrospective multicenter study was to analyse the frequency of pleural and parenchymal abnormalities on high-resolution computed tomography (HRCT) in a population of automobile mechanics.

Methods: The study population consisted of 103 automobile mechanics with no other source of occupational exposure to asbestos, referred to three occupational health departments in the Paris area for systematic screening of asbestos-related diseases. All subjects were examined by HRCT and all images were reviewed separately by two independent readers; who in the case of disagreement discussed until they reached agreement. Multiple logistic regression models were constructed to investigate factors associated with pleural plaques.

Results: Pleural plaques were observed in five cases (4.9%) and interstitial abnormalities consistent with asbestosis were observed in one case. After adjustment for age, smoking status, and a history of non-asbestos-related respiratory diseases, multiple logistic regression models showed a significant association between the duration of exposure to asbestos and pleural plaques.

Conclusions: The asbestos exposure experienced by automobile mechanics may lead to pleural plaques. The low prevalence of non-malignant asbestos-related diseases, using a very sensitive diagnostic tool, is in favor of a low cumulative exposure to asbestos in this population of workers.

Keywords: asbestos; automobile mechanics; brakes; high-resolution computed tomography; pleural plaques

INTRODUCTION

The presence in the past of chrysotile asbestos in automobile brakes or clutches has given rise to concerns about increased risks of asbestos-related

diseases among automobile mechanics (Lemen, 2004; Paustenbach et al., 2004; Welch, 2007).

A large number of automobile mechanics have been potentially exposed to asbestos. According to the data of the 1999 census, the estimated population of automobile mechanics, aged 16–60 years, in France was 242 360 (Imbernon et al., 2005).

Partly due to the different methodologies employed, available data concerning atmospheric measurements

*Author to whom correspondence should be addressed.
Tel: +33147107754; fax: +33147107768;
e-mail: jacques.ameille@rpc.aphp.fr

of asbestos fibers during brake or clutch repair show disagreement (Rohl *et al.*, 1976; Rodelsperger *et al.*, 1986; Flato *et al.*, 1995; Paustentuch *et al.*, 2003; Finley *et al.*, 2007; Blake *et al.*, 2008; Cohen and Van Orden, 2008; Jiang *et al.*, 2008), contributing to uncertainties about the level of risk of asbestos-related diseases in automobile mechanics.

Cases of malignant pleural mesothelioma have been reported in automobile mechanics (Paur *et al.*, 1985; Rodelsperger *et al.*, 1986; Bittor *et al.*, 2003). However, published meta-analysis of the epidemiological literature that examined the risk of pleural mesothelioma (Wong, 2001; Goodman *et al.*, 2004) or lung cancer (Goodman *et al.*, 2004) did not show any elevated meta-relative risks.

Very few data concerning non-malignant asbestos-related diseases or impairment of lung function in automobile mechanics have been published (Boillat and Loh, 1973; Marcus *et al.*, 1987; Dahlqvist *et al.*, 1992; Flato *et al.*, 1995). Only one study, using full-size chest radiographs, investigated the prevalence of pleural plaques (Marcus *et al.*, 1987). To our knowledge, no study using chest computed tomography (CT) scan for the detection of asbestos-related diseases in automobile mechanics has been published.

The aim of the present retrospective multicenter study was to analyze the frequency of pleural and parenchymal abnormalities on chest high-resolution computed tomography (HRCT), in a population of automobile mechanics, referred by occupational physicians to occupational health departments in the Paris area, for systematic screening of asbestos-related diseases.

STUDY POPULATION AND METHODS

Study population

The eligible population consisted of all automobile mechanics, referred to occupational health departments in university hospitals of Garches, Créteil, and Paris-Fernand Widal, between 1 January 2002 and 30 June 2009, for HRCT detection of asbestos-related diseases. These subjects were systematically referred by occupational physicians because of known exposure to asbestos. As social welfare benefits, such as early retirement allowance or pension, are provided in France for workers with asbestos-related diseases, including pleural plaques, when confirmed by chest CT scan examinations, a consensus conference, held in Paris in January 1999, recommended the use of chest CT scan, after a sufficient latency period, for the surveillance of workers occupationally exposed to asbestos

(Société française de médecine du travail, Société de pneumologie de langue française 1999). Subjects with sources of occupational exposure to asbestos other than their work as automobile mechanics were excluded from the analysis.

METHODS

For each subject, information was collected on gender, age, previous chest diseases, and smoking status. Subjects were classified into three categories according to tobacco consumption: smokers, ex-smokers, and non-smokers. Ex-smokers were defined as those who had quit smoking for at least 1 year.

For each subject, the year of starting work as an automobile mechanic was recorded, together with the type of work (light vehicles, trucks, mixed activity), duration of exposure, and the time interval between start of asbestos exposure and the HRCT (latency). For calculation of the duration of exposure to asbestos, only work prior to 1997 (year in which asbestos was banned in France) was taken into account. Information on the use of compressed air during brake cleaning was also collected.

High-resolution computed tomography

Spiral HRCT acquisition parameters were defined in accordance with the French Thoracic Imaging Society guidelines: full inspiration, slice thickness of 1.5–5 mm, pitch of 1.5–2.0, 120 KV, 60–150 mAs. All CT scans were performed without intravenous contrast agent.

Localized acquisition in the prone position was required in the case of suspected gravitational lung abnormalities.

All images were reviewed separately by two independent readers in each occupational health department (a radiologist and an occupational diseases specialist), who if they disagreed discussed further until they reached agreement concerning the presence of pleural or parenchymal abnormalities consistent with an asbestos-related disease.

A structured form, elaborated from a previous study (Ameille *et al.*, 2007), was filled out, describing the features of pulmonary abnormalities consistent with asbestosis (thickened interstitial short lines: septal lines, non-septal lines, honeycombing, curvilinear subpleural lines, ground glass opacities) and pleural abnormalities.

When associated with rounded atelectasis or parenchymal bands, pleural thickenings were considered to correspond to diffuse pleural thickening (fibrosis of visceral pleura) (Gevenois *et al.*, 1998; Beigelman-Aubry *et al.*, 2007).

Pleural plaques were considered to be present in the case of circumscribed quadrangular pleural elevations, with sharp borders and tissue density, sometimes calcified, or in the case of multiple, bilateral, less typical images, with a typical topography (Beigelman-Aubry et al., 2007).

Pleural abnormalities not meeting the criteria for pleural plaques or diffuse pleural thickening were considered to be nonspecific pleural thickening.

Statistical analysis

Values for continuous variables were expressed as the mean (standard deviation, SD). Bivariate analysis was performed to study factors associated with pleural plaques, using Fisher's exact test for qualitative variables or Wilcoxon's test for continuous variables. A *P*-value <0.05 was considered statistically significant. Multiple logistic regression models were constructed, including age, previous respiratory diseases, smoking status, latency, and duration of exposure to asbestos. Due to the strong collinearity between latency and duration of exposure, separate models were constructed. A *P*-value <0.05 was considered statistically significant.

Statistical analysis was performed using SAS software version 9.1 (SAS Institute Inc, Cary, NC, USA).

Informed consent was obtained from all subjects. The study was approved by the French national committee for data protection (Commission Nationale de l'Informatique et des Libertés: CNIL).

Independent ethics committee approval was not necessary, as chest CT scans are recommended in France for screening purposes in asbestos-exposed workers (Société Française de Médecine du Travail, 1999).

RESULTS

During the study period, 242 automobile mechanics were referred to the occupational health departments participating in the study. Eighty-nine automobile mechanics with other possible sources of asbestos exposure, and 50 automobile mechanics with no available HRCT were excluded. The study population therefore consisted of 103 subjects.

General characteristics of the study population are given in Table 1. All subjects were male with a mean age of 54.4 (7.2) years; 69 (68%) were smokers or ex-smokers. Mean latency was 36.8 (7.0) years (range: 19–57 years). Mean duration of asbestos exposure was 22.6 (8.1) years (range: 4–45 years). Most of the study population (83.5%) used compressed air for brake cleaning.

Table 1. Characteristics of the study population (*n* = 103).

	<i>n</i>	%	Mean (SD)
Gender: male	103	100	
Age (years)			54.4 (7.2)
<50	23	22.3	
50–59	62	60.2	
>60	18	17.5	
Tobacco consumption			
Non-smokers	32	31.7	
Ex-smokers	21	40.6	
Smokers	48	27.7	
Latency ^a (years)			36.8 (7.0)
<20	1	1.0	
20–30	10	9.7	
>30	92	89.3	
Duration of exposure ^b (years)			22.3 (8.1)
<20	31	30.1	
20–30	50	48.5	
>30	22	21.4	
Brake repair of trucks	42	41.8	
Use of compressed air	86	83.5	

^aTime interval between start of asbestos exposure and HRCT.

^bDuration of asbestos exposure before 1997.

Interstitial abnormalities consistent with asbestosis were observed in one subject. This subject had worked for 24 years as an automobile mechanic, ensuring brake repairs on trucks for >8 h a week. Interstitial abnormalities consisted of bilateral honeycombing, ground-glass opacities, and non-septal lines in the postero-inferior zones of the lungs. Five automobile mechanics had pleural plaques, four had diffuse pleural thickening, and nine had nonspecific pleural thickening. No case of lung cancer or pleural mesothelioma was detected. Pleural plaques were diagnosed in each occupational health department participating in the study (one in two departments, two in one department). The five subjects with pleural plaques had no previous chest diseases likely to explain pleural thickening.

Characteristics of the subjects according to HRCT abnormalities are presented in Table 2. Subjects with nonspecific pleural thickening had similar asbestos exposure to that of the subjects without HRCT abnormalities, in terms of latency or duration of exposure. Subjects with pleural plaques and subjects with diffuse pleural thickening had longer latency and duration of asbestos exposure than subjects without HRCT abnormalities (Table 2).

In comparison with subjects without pleural plaques, subjects with pleural plaques had significantly

Table 2. Characteristics of the subjects according to HRCT abnormalities.

	No abnormalities <i>n</i> = 84	Nonspecific pleural thickening <i>n</i> = 9	Pleural plaques <i>n</i> = 5	Diffuse pleural thickening <i>n</i> = 4	Asbestosis ^a <i>n</i> = 1
Age: m in years (SD)	53.9 (7.4)	54.1 (4.0)	59.0 (6.9)	59.5 (7.1)	63
Latency ^b : m in years (SD)	36.2 (7.1)	36.0 (4.8)	42.2 (5.6)	42.5 (5.9)	47
Duration of exposure ^c : m in years (SD)	21.8 (8.0)	23.1 (5.8)	31.0 (3.9)	27.8 (12.9)	24
Brake repair of trucks: <i>n</i> (%)	34 (40.5)	5 (55.6)	3 (60.0)	0 (0)	1 (100)
Use of compressed air: <i>n</i> (%)	68 (81.0)	8 (88.9)	5 (100)	4 (100)	1 (100)
Previous respiratory diseases: <i>n</i> (%)	10 (11.9)	3 (33.3)	1 (20.0)	1 (25.0)	0

m, mean.

^aInterstitial abnormalities consistent with asbestosis.^bTime interval between start of asbestos exposure and HRCT.^cDuration of asbestos exposure before 1997.

Table 3. Characteristics of the subjects according to the presence of pleural plaques.

	Pleural plaques		<i>P</i> ^a
	yes, <i>n</i> = 5	no, <i>n</i> = 98	
Age: m (SD)	59.0 (6.9)	54.2 (7.1)	0.12
Latency ^b : m (SD)	42.2 (5.6)	36.6 (7.0)	0.047
Duration of exposure ^c : m (SD)	31.0 (3.9)	22.2 (8.0)	0.01
Brake repair of trucks: <i>n</i> (%)	3 (60.0)	40 (40.6)	
Use of compressed air: <i>n</i> (%)	5 (100.0)	81 (81.8)	

m, mean.

^aWilcoxon test.^bTime interval between start of asbestos exposure and HRCT.^cDuration of asbestos exposure before 1997.

longer latency and duration of asbestos exposure (Table 3).

After adjustment for age, smoking status, and history of previous asbestos-unrelated respiratory diseases, multiple logistic regression models showed a statistically significant association between the duration of asbestos exposure and the presence of pleural plaques (Table 4).

DISCUSSION

In this population of automobile mechanics referred for systematic detection of asbestos-related diseases by HRCT, only one case of interstitial abnormalities consistent with asbestosis, five cases of pleural plaques (4.9%) and four cases (3.9%) of diffuse pleural thickening were observed. The five sub-

Table 4. Pleural plaques: study of determinants by multivariate logistic regression analysis.

	aOR ^a (95% CI)	<i>P</i>
Model 1		
Age (years)	1.0 (0.8–1.2)	NS ^b
Previous respiratory diseases (yes versus no)	1.2 (0.1–13.7)	NS
Smokers or ex-smokers (yes versus no)	2.0 (0.2–20.7)	NS
Latency ^c (years)	1.1 (0.9–1.4)	NS
Model 2		
Age (years)	1.0 (0.9–1.2)	NS
Previous respiratory diseases (yes versus no)	1.1 (0.1–17.8)	NS
Smokers or ex-smokers (yes versus no)	3.7 (0.3–46.3)	NS
Duration of exposure ^d (years)	1.2 (1.0–1.4)	0.048

^aAdjusted odds ratio.^bNon significant.^cTime interval between start of exposure and HRCT.^dDuration of exposure before 1997.

jects with pleural plaques had no known history of chest disease likely to explain pleural abnormalities.

The prevalence of pleural plaques in this study is similar to that observed in the only previous published study that investigated the frequency of pleural plaques in car mechanics (Marcus *et al.*, 1987). In that study, based on chest radiographs, pleural plaques, defined by pleural thickening at least 5-mm thick on the chest wall or 3-mm thick on the diaphragm, were observed in 41 of 925 car mechanics (4.4%). However, our results must be compared cautiously with those reported by (Marcus *et al.*, 1987) due to the selection factors and the use of different

tools for the radiological detection of asbestos-related abnormalities.

The prevalence of pleural plaques observed in this study was similar to that observed on HRCT examinations in subjects exposed to indoor asbestos pollution (De Raeve et al., 2001) or in urban transportation workers with low cumulative exposure to asbestos (Ameille et al., 2007).

The main limitations of this study are the relatively small number of subjects, the absence of an unexposed control group, and the lack of objective assessment of the level of occupational exposure to asbestos. The representativity of the study sample is also questionable. However, inclusion criteria had nothing to do with symptoms or abnormalities on chest radiographs.

One of the strengths of this study is the use of HRCT for detection of asbestos-related diseases. HRCT has been shown to be much more sensitive and specific than chest radiographs for the detection of early pleural and parenchymal changes (Aberle et al., 1988a; 1988b; Friedman et al., 1988; Staples et al., 1989; Neri et al., 1994; Gevenois et al., 1994).

Another strength is the standardized reading, with systematic double interpretation followed by consensus reading in the case of disagreement.

Furthermore, only automobile mechanics with no other known occupational exposure to asbestos and with a sufficiently long latency (>20 years in 99% of cases, >30 years in 89% of cases) to allow evaluation of asbestos-related diseases, especially pleural plaques, were included in the study (Schwartz, 1991; Paris et al., 2009).

In the absence of a control group, demonstration of a significant relationship between pleural plaques and duration of exposure (used as a surrogate of cumulative exposure to asbestos) is an argument in favor of the causal role of asbestos (Paris et al., 2009).

The low prevalence of pleural plaques and interstitial abnormalities in this study is an indirect argument in favor of low cumulative exposure to asbestos for most automobile mechanics. These data are consistent with the meta-analysis of epidemiologic studies that investigated the incidence of lung cancer and pleural mesothelioma in automobile mechanics and failed to demonstrate increased meta-relative risks (Wong, 2001; Goodman et al., 2004). Our data are also consistent with the results of a recently published population-based case-control study (Rohland et al., 2010) that reported a slight but not significant increased risk of pleural mesothelioma in male car mechanics (odds ratio = 1.50; 95% confidence interval: 0.76–2.95).

Many measurements in car repair workshops have shown that occupational exposure of car mechanics to asbestos is generally fairly low. Robl et al., 1976 reported airborne chrysotile concentrations of up to 30 f ml^{-1} . However, these were peak exposure concentrations measured during brake cleaning with the use of compressed air. According to Pausenbach et al., 2003, analysis of the available published data indicates that 90% of 8 h time-weighted average (TWA) concentrations, experienced by mechanics servicing brakes in the 1970s, was $<0.1 \text{ f ml}^{-1}$ (mean 0.04 f ml^{-1}). In Sweden, Plato et al., 1995, using a model based on data from the international literature and quantitative asbestos measurements performed in 1976–1978 in Swedish car repair workshops, estimated the mean cumulative exposure to be $2.6 \text{ f ml}^{-1} \times \text{years}$, in a group of 103 car and bus mechanics, with >15 years of asbestos exposure. Car mechanics had a TWA exposure range of $0.11\text{--}0.41 \text{ f ml}^{-1}$ (mean 0.21 f ml^{-1}) in 1965 and $0.003\text{--}0.08 \text{ f ml}^{-1}$ in 1985 (mean 0.021 f ml^{-1}). In Germany, based on dust measurements in 76 service stations and the occupational histories of 210 vehicle mechanics with an average duration of employment of 21 ± 10 years, a mean cumulative exposure to asbestos was estimated to be $0.54 \pm 1.1 \text{ f ml}^{-1} \times \text{years}$ for car mechanics (Rodelsperger et al., 1986). Finley et al., 2007 recently developed estimates of cumulative chrysotile exposures experienced by automobile mechanics working with brakes and manual clutches. They reported that the 95th percentile and 99th percentile cumulative exposures for vehicle mechanics in the 1970s were 2.0 and $5.7 \text{ f ml}^{-1} \times \text{years}$, respectively.

Another explanation for the low prevalence of asbestos-related diseases in automobile mechanics might be that decomposition of asbestos occurs during normal use of brakes due to thermal decomposition into forsterite, a non-fibrous material (Lemen, 2004).

CONCLUSION

The asbestos exposure experienced by French car mechanics may lead to pleural plaques. The low prevalence of non-malignant asbestos-related diseases, despite the use of HRCT, which is a very sensitive diagnostic tool, is in favor of low cumulative exposure to asbestos in the study population.

FUNDING

The Caisse nationale d'assurance maladie des travailleurs salariés (Département des risques professionnels); Caisse régionale d'assurance maladie d'Ile de France.

Acknowledgements—The authors declare that they have no conflict of interest.

REFERENCES

- Aberle DR, Gamsu G, Ray CS. (1988a) High-resolution CT of benign asbestos-related diseases: clinical and radiographic correlation. *AJR Am J Roentgenol*; 151: 883–91.
- Aberle DR, Gamsu G, Ray CS *et al.* (1988b) Asbestos-related pleural and parenchymal fibrosis: detection with high-resolution CT. *Radiology*; 166: 729–34.
- Ameille J, Mattei N, Laurent F *et al.* (2007) Computed tomography findings in urban transportation workers with low cumulative asbestos exposure. *Int J Tuberc Lung Dis*; 11: 1352–57.
- Beigelman-Aubry C, Ferretti G, Mompoin D *et al.* (2007) [Computed tomographic atlas of benign asbestos related pathology]. *Rev Mal Respir*; 24: 759–89.
- Blake CL, Dotson GS, Harbison RD (2008) Evaluation of asbestos exposure within the automotive repair industry: a study involving removal of asbestos-containing body sealant and drive clutch replacement. *Regul Toxicol Pharmacol*; 58: 324–31.
- Boillat MA, Lob M. (1973) [Risk of asbestosis in workers employed in replacing automobile brake linings]. *Schweiz Med Wschr*; 103: 1354–9.
- Butnor KJ, Sporn TA, Roggli VL. (2003) Exposure to brake dust and malignant mesothelioma: a study of 10 cases with mineral fiber analysis. *Ann Occup Hyg*; 47: 325–30.
- Cohen HJ, Van Orden DR. (2008) Asbestos exposures of mechanics performing clutch service on motor vehicles. *J Occup Environ Hyg*; 5: 148–56.
- Dahlqvist MR, Alexandersson R, Hedenstierna G. (1992) Lung function and exposure to asbestos among vehicle mechanics. *Am J Ind Med*; 22: 59–68.
- De Raeve H, Verschakelen JA, Gevenois PA *et al.* (2001) Observer variation in computed tomography of pleural lesions in subjects exposed to indoor asbestos. *Eur Respir J*; 17: 916–21.
- Finley BL, Richter RO, Mowat F *et al.* (2007) Cumulative asbestos exposure for U.S. automobile mechanics involved in brake repair (circa 1950s–2000). *J Expo Sci Environ Epidemiol*; 17: 644–55.
- Friedman AC, Fiel SB, Fisher MS *et al.* (1988) Asbestos-related pleural disease and asbestosis: a comparison of CT and chest radiography. *AJR Am J Roentgenol*; 150: 269–75.
- Gevenois P, de Maertelaer V, Madani A *et al.* (1998) Asbestosis, pleural plaques and diffuse pleural thickening: three distinct benign responses to asbestos exposure. *Eur Respir J*; 11: 1021–7.
- Gevenois PA, De Vuyst P, Dedeire S *et al.* (1994) Conventional and high-resolution CT in asymptomatic asbestos-exposed workers. *Acta Radiol*; 35: 226–9.
- Goodman M, Teta MJ, Hessel PA *et al.* (2004) Mesothelioma and lung cancer among motor vehicle mechanics: a meta-analysis. *Ann Occup Hyg*; 48: 309–26.
- Imbemon E, Marchand JL, Garras L, Goldberg M. (2005) [Quantitative assessment of the risk of lung cancer and mesothelioma among automobile mechanics]. *Rev Epidemiol Santé Pub*, 53: 491–500.
- Jiang GC, Madl AK, Ingmundson KJ *et al.* (2008) A study of airborne chrysotile concentrations associated with handling, unpacking, and repacking boxes of automobile clutch discs. *Regul Toxicol Pharmacol*; 51: 87–97.
- Lemen RA. (2004) Asbestos in brakes: exposure and risk of disease. *Am J Ind Med*; 45: 229–37.
- Marcus K, Järholm BG, Larsson S. (1987) Asbestos-associated lung effects in car mechanics. *Scand J Work Environ Health*; 13: 252–4.
- Neri S, Antonelli A, Falashi F *et al.* (1994) Findings from high-resolution computed tomography of the lung and pleura of symptom free workers exposed to amosite who had normal chest radiographs and pulmonary function tests. *Occup Environ Med*; 51: 239–43.
- Paris C, Thierry S, Brochard P *et al.* (2009) Pleural plaques and asbestosis: dose and time response relationships based on HRCT data. *Eur Respir J*; 34: 72–9.
- Paur R, Weitowitz HZ, Rodelsperger K, John H. (1985) [Pleural mesothelioma following asbestos exposure during brake repair in the automobile trade: case report]. *Prax Klin Pneumol*; 39: 362–6.
- Paustenbach DJ, Finley BL, Lu ET *et al.* (2004) Environmental and occupational health hazards associated with the presence of asbestos in brake linings and pads (1900 to present): a “state-of-the-art” review. *J Toxicol Environ Health*; 7: 25–80.
- Paustenbach DJ, Richter RO, Finley BL *et al.* (2003) An evaluation of the historical exposures of mechanics to asbestos in brake dust. *Appl Occup Environ Hyg*; 18: 786–804.
- Plato N, Tornling G, Hogstedt C *et al.* (1995) An index of past asbestos exposure as applied to car and bus mechanics. *Ann Occup Hyg*; 39: 441–54.
- Rodelsperger K, Jahn B, Bruckel J *et al.* (1986) Asbestos dust exposure during brake repair. *Am J Ind Med*; 10: 63–72.
- Rohl AN, Langer AM, Wolff MS *et al.* (1976) Asbestos exposure during brake lining maintenance and repair. *Environ Res*; 12: 110–28.
- Rolland P, Gramond C, Lacourt A *et al.* (2010) Occupations and industries in France at high risk of pleural mesothelioma: a population based-study (1998–2002). *Am J Ind Med*; 53: 1207–19.
- Staples CA, Gamsu G, Ray CS *et al.* (1989) High resolution computed tomography and lung function in asbestos-exposed workers with normal chest radiographs. *Am Rev Respir Dis*; 139: 1502–8.
- Schwartz DA. (1991) New developments in asbestos-induced pleural disease. *Chest*; 99: 191–8.
- Société française de médecine du travail, Société de pneumologie de langue française. (1999) Conférence de consensus pour l’élaboration d’une stratégie de surveillance médicale clinique des personnes exposées à l’amiante. *Rev Mal Respir*; 16: 1190–362.
- Welch LS. (2007) Asbestos exposure causes mesothelioma, but not this asbestos exposure: an amicus brief to the Michigan Supreme court. *Int J Occup Environ Health*; 13: 318–27.
- Wong O. (2001) Malignant mesothelioma and asbestos exposure among auto mechanics: appraisal of scientific evidence. *Regul Toxicol Pharmacol*; 34: 170–7.