Family of 4 consumes 1.8 Liters / month

$\frac{1.8}{4} = 0.45$

(1)

450

1.41

450

1800

450

634.50

21.15

30 / 634.5

60

34

30

45

30

150

30.5

70 / 21.15

210

150

.45 L / Mo / Person - Rice Oil

.45 L = 450 ml / mo / person

450 × 56 1.41 = 634.5 grams

634.5 grams / mo / person

$\frac{634.5}{30} = 21.15$ grams / day / person

$\frac{634.5}{30} = 21150$ mgs

S.G. 1.41 Used
Same As PCB-1248

Assume weight 154 #/person

$\frac{154}{2.2} = 70$ Kg / person

Assume wt 110 # or 50 Kg

21,150 mgs / day / person

$\frac{21150}{50} = 423$ mgs / Kg / Day Fed

200 ppm = 423 × .0002 = 0.0846 mgs

On Page 43

$\frac{21,150}{70} = 302$ mgs / Kg / Day

Amount of oil consumed

302 × .0002 = 0.0604 mgs / Kg / Day

The First Interim Report by the Analytical Lab - Committee Nov 18, 1968 -

Translation of Japanese Article

Discussion a) From the above (a) and (b) results the patients oil was confirmed to contain 1000 - 1500 ppm (as Cl) organic chlorine

Control oil Page 42 - Contained Trace or more.

0277637

It can be concluded since the Cl content of Kano-Chlor is about 48% then the patient oil contains 2000 - 3000 ppm Kano-Chlor

$\frac{1000 - 1500}{.48}$ = Approx 2000 - 3000 ppm Kano-Chlor

PCB-ARCH0744086

(2)

On Page 7 "The Symptomatology of Yusho ("Chlorachlorphenyls" Poisoning) in Dermatology"
Translation of Japanese article

3 paragraphs states about 200 ppm of Kanechlor 400 was contained in oils taken by patients.

Note: This translation is not confirmed with Page 63 1st Interim Report.

On Page 7 report above states

The amount taken by patients (families (4 persons on average)) was 1.8 L within a month

2 = Liters
1 L = 1000 ml

1.8 L per family of 4 is
0.45 L per person per month
$\frac{0.45 L}{30} = 0.015 L$ per person / day

0.015 L / person / day is 15.2 ml

The S.G. of 1248 from Table A-3 Page A 213 of Monsanto Specification is 1.405 @ 65°C
1.415 @ 15.5°C

Assume wt 50 Kg = 110#

$\frac{21.15 gm}{50 mg/Kg/day} = 423$

0277638

a S.G. of 1.41 is used for the calculation below.

15.2 ml x 1.41 = 21.15 grams / person / day oil

On Page 5 "Epidemiologic Study on "Yusho" a poisoning caused by Rice Oil Contaminated with Polychlorinated Biphenyls"
Translation of Japanese article

PCB-ARCH0744087

(3)

Top of page 5: rice oil produced in Shiffard and used by patients had been contaminated with about 2000 ppm of Kanechlor 400.

In middle of page 5: "The minimum total dose of Kanechlor 400 to cause the diseases was estimated to be approx 0.5 g. Note this is 500 mg

On Page E 33 (E Biological Effects) in summary at back of book,

"More recently, Kuratsune et al among Yusho case studied the smallest toxic dose was 500 mgs ingested over approximately 50 days by an individual weighing approximately 50 Kg."

This calculates to $\frac{500}{50\ Kg.}$ = 10 mgs/Kg over 50 days

$\frac{10\ mg/Kg}{50\ Days}$ = 0.2 mgs/Kg/day

On Page E 43 - E Biological Effects

In man it has been estimated that as little as 10 mg/Kg over 50 days causes chloracne.

Note 0.2 mgs/Kg/day × 50 days = 10 mg/Kg

0277639

PCB-ARCH0744088


(4)

2 ppm inc.
5 ppm in feeds


Continued Table 5
Page E-34 & E-35

Relation between the amounts of K rice oil used by patients and other clinical severity


Amount of Oil	Non-Affected	Light Case	Severe Case	Total
Less than 720 ml	10 (12.5%)	39 (48.5%)	31 (39.0%)	80 (100%)
720 - 1440 ml	0 (0.0)	14 (31.8)	31 (69%)	45 (100%)
More than 1440 ml	0 (0.0)	3 (14.0)	18 (86.0)	21 (100%)



Chlorinated biphenyl
m = 1, 2, 3, 4, 5



Chlorinated dibenzofuran
n = 1, 2, 3, 4
Highly Toxic



Chlorinated dibenzodioxin
N = 1, 2, 3, 4 &
Highly Toxic

FDA Pesticide Ind. Rep. 37(34) March 18, 1977

Milk (Fat Basis)	2.5 ppm
Dairy Products (F.B.)	2.5
Poultry (F.B.)	5.0
Eggs	0.5
Finished Animal feed	0.5
Animal Feed Components	5.0
Fish (Edible Portion	5.0

0277640

PCB-ARCH0744089

Weights of Persons
100 #s = 45.5 Kg Assume all eat 4.4 lbs/day or 2 Kg
110 #s = 50.0 Kg
154 #s = 70.0 Kg

See Page 4 My Notes
Using amounts of Rice Oil From Table 5 and assuming A Specific Gravity of 1.2 For Rice Oil

Amount of Oil = 144.0 Grams Using 1.2 SG.

720 x 1.2 = 864.0 grams oil

1440 x 1.2 = 1728

Using the lowest figure of 720 ml oil or 864 g oil and 200 ppm PCB in oil and 50 days exposure
864 g x 1000 mg/g = 864,000 mgs of Rice oil

864,000 mg x 200 ppm PCB mark =

864,000 mg x .0002 = 172.8 mgs PCB

172.8 mgs PCB / 50 Kg = 3.46 mgs PCB/Kg

3.46 mgs PCB/Kg / 50 days = 0.069 mgs PCB/Kg/day

Using the lowest figure 720 ml oil or 864 grams

864,000 mgs rice oil x 2000 ppm = 1728 mgs PCB

1728 mgs PCB / 50 Kg = 34.6 mgs PCB/Kg

34.6 mgs PCB/Kg / 50 days = 0.69 mgs PCB/Kg/day

0277641

PCB-ARCH0744090

Using the highest figure taken 5 of 1440 mL of oil, Temperature and assuming Specific Gravity of the oil as 1.2 kg of patient 50 Kg and 50 days exposure

Calculation using 200 ppm PCB in oil

1440 mL oil x 1.2 = 1728 grams of oil

1728 x 1000 = 1,728,000 mgs oil

1,728,000 mgs oil x 200 ppm PCB in oil = 345.6 mgs PCB

$\frac{345.6 \text{ mgs PCB}}{50 \text{ Kg}}$ = 6.91 mgs PCB/Kg

$\frac{6.91 \text{ mgs PCB/Kg}}{50 \text{ days}}$ = 0.138 mgs PCB/Kg/day

or 138 µg PCB/Kg/day

Calculation using 2000 ppm PCB in oil

1,728,000 mgs oil x 2000 ppm PCB in oil = 3456 mgs PCB

$\frac{3456 \text{ mgs PCB}}{50 \text{ Kg}}$ = 69.1 mgs PCB/Kg

$\frac{69.1 \text{ mgs PCB/Kg}}{50 \text{ days}}$ = 1.38 mgs PCB/Kg/day

or 1380 µg PCB/Kg/day

0277642

PCB-ARCH0744091