

ATMOSPHERIC TRANSPORT OF POLYCHLOROBIPHENYLS TO THE NORTH ATLANTIC*

GEORGE R. HARVEY and WILLIAM G. SVEINHAFUR

Woods Hole Oceanographic Institution, Woods Hole,
Massachusetts 02543, U.S.A.

(First received 5 August 1973 and in final form 22 November 1973)

Abstract. Concentrations of polychlorobiphenyls (PCBs) were measured in the atmosphere over the western North Atlantic. Both the particulate and vapor phase were extracted using a new sampler designed for this work. Previously PCB levels ranged from 5 ng m^{-3} near the northeast coast of the U.S. to 0.05 ng m^{-3} on the Grand Banks. The PCB concentration in the atmosphere decreases exponentially with distance from land. It is concluded that wind transport is the major source of chlorinated hydrocarbons in the oceans.

INTRODUCTION

Polychlorobiphenyl (PCB) concentrations have been measured in the water and organisms of the open Atlantic Ocean (Harvey *et al.*, 1973). It has generally been assumed that PCBs would be delivered to the open ocean mainly by the wind (Nisbet and Sarofim, 1972) similar to DDT (Seba and Prospero, 1971) with fluvial and sewage inputs contributing smaller quantities (Corcoran and Lee, 1970; Holden, 1970). However, no atmospheric PCB measurements have been reported. Risebrough (1972) was unable to detect PCBs on equatorial Atlantic dust collected on Barbados (Risebrough *et al.*, 1968). Therefore, it seemed likely that PCBs were transported mainly in the vapor or aerosol phase and we set out to design a sampling system capable of trapping both particulate and vapor phase chlorinated hydrocarbons. Atmospheric PCB measurements were important to our studies because we had previously observed no inshore-open ocean gradient of PCBs in the surface waters of the North Atlantic (Harvey *et al.*, 1973). Thus, we wished to determine if the atmospheric concentrations of PCB reflected this observation.

In this paper we present a new atmospheric sampling technique, and the results of our PCB measurements at four stations over the western North Atlantic. The sampling was performed during the first half of 1973.

PROCEDURES

The sampling system consisted of a Hurricane Air Pump (Bendix Corp.) fitted with a 102-mm filter holder on the intake port. The exhaust pipe was fitted with several meters of extension so as not to interfere with the intake. A $0.3 \mu\text{m}$ glass fiber filter, which had been baked at 500°C for 12 h, was used on the intake. The filter holder chamber, behind the filter was filled with $1/4$ in. ceramic saddles (distillation column packing, Fisher Scientific) which were coated with 0.25 per cent OV-17 silicone oil. The pump was run on low speed and generally achieved a flow rate of $22\text{--}24 \text{ ft}^3 \text{ min}^{-1}$.

The saddles had to be cleaned initially by distilling several liters of water through a column packed with about 500 g of the commercial product. The OV-17 was purified by applying 1.25 g to the top of a $1.5 \times 15 \text{ cm}$ column of silica. Interfering impurities were

* Woods Hole Oceanographic Institution Contribution No. 3173.

eluted with 80 cm³ of hexane. The OV-17 was eluted with 150 cm³ of benzene. The benzene solution was diluted with 100 cm³ of hexane and 500 g of steam cleaned saddles were added. The solvent was evaporated to dryness on a rotary evaporatory at 50° with a water aspiratory. The coated saddles could be stored in glass jars or used immediately. About 125 g are required to fill the filter holder chamber.

Recovery of the chlorinated hydrocarbons from the coated saddles was accomplished by shaking with two 100 cm³ portions of hexane in a separatory funnel. The hexane solution was dried with sodium sulfate and concentrated to 10 cm³ by distillation. The extract was applied to the top 1.5 × 15 cm column of silica and the chlorinated hydrocarbons were eluted with 80 cm³ of hexane. The hexane eluate was concentrated to 1 cm³ in a Kuderna-Danish concentrator. If necessary, the hexane was further concentrated with a stream of Florisil-purged nitrogen. No further cleanup was necessary before gas chromatography.

The concentrated extract was analyzed on a gas chromatograph equipped with a ⁶³Ni electron capture detector. The column was a 2 m × 2 mm (i.d.) glass coil packed with a 1:1 mixture of 8 per cent QF-1 and 2 per cent OV-17 on 100-120 mesh Anakrom. The oven temperature was 190°. The chromatogram was quantified by comparing the heights of seven peaks in the sample with the same matching peaks in a standard of commercial Aroclor 1254 (Monsanto) as shown in Fig. 1. The blank (a) was produced by leaving the entire apparatus at the sampling site for 24 h without running the pump. The saddles were then extracted and chromatographed as described above. The sample chromatogram (b)




Fig. 1. Typical PCB chromatogram of a Grand Banks air sample (b). The standard Aroclor 1254 is shown in (c) and the blank is shown in (a). The seven dotted peaks were used for the quantification.

is from the Grand Banks. The presence of the PCBs was confirmed by perchlorination to decachlorobiphenyl according to the procedure of Armour (1973). The decachlorobiphenyl synthesized from the Bermuda samples was further confirmed by mass spectrometry.

Extraction of the glass fiber filters with hexane followed by concentration and chromatography revealed PCB concentrations which ranged from undetectable in the Grand Banks samples to 0.04 ng m^{-3} in the Vineyard Sound samples.

The efficiency of trapping was tested by applying known quantities of 3,4,2'-trichlorobiphenyl; 2,4,5,2',4',5'-hexachlorobiphenyl (Analabs, Inc.) or Aroclor 1254 (Monsanto Co.) on the glass fiber filter. The average recovery after 20 h of pumping was 70 per cent for the pure isomers or the mixture.

The sampling stations are shown in Fig. 2. Vineyard Sound (Massachusetts) samples were collected from the roof of the pilot cabin of the W.H.O.I. vessel *R/V Asterias*. The Georges Bank collections were made from the 01 deck of the *R/V Albatross IV* (National Marine Fisheries Service). The Bermuda samples were taken from a 20-m tower on the south-west shore of the island. Finally, the collections on the Grand Banks were performed from the exploratory drilling rig *Sednet* 1 of Amoco Canada Ltd. at Station Razorbill. Procedural blanks from each station averaged 0.03 ng m^{-3} for 500 m^3 .




Fig. 2. Sampling sites over the western North Atlantic.

RESULTS

The analytical results for the four stations are set out in Table 1. In addition, the DDT family (*p*-DDT) was detected in the Bermuda samples. The concentration of DDT in the four samples decreased from $5.3 \times 10^{-11} \text{ g m}^{-3}$ on 12 February, to $1.7 \times 10^{-11} \text{ g m}^{-3}$ on 15 February (1973). The most likely source was the southeastern U.S. cotton fields directly to the west. At the other sampling sites the *p*-DDT concentration was less than $1 \times 10^{-12} \text{ g m}^{-3}$. The PCB/*p*-DDT ratio was also found to be greater than 30 in the surface water (Harvey *et al.*, 1973). The PCB concentrations were highest in Vineyard Sound which lies only 150–250 km from the nearest windward industrial sources, i.e. the Boston-Hartford-New York-New Jersey complex. The concentrations of PCB over the Grand Banks which are over 2000 km from the industrial complex of the northeast United States were 100 times less than in Vineyard Sound. In fact, when the average concentrations at each station are plotted against the average distance of the site from major industrial

Table 1. PCB concentrations over the western North Atlantic

Station	Date (1973)	Sample volume (m ³)	Wind direction	PCB ng m ⁻³ (calc. as Aruclor, 1254)
Bermuda	12 February	560	WNW	0.5
(32°20'N; 64°40'W)	13 February	480	W	0.4
(32°20'N; 64°40'W)	14 February	820	Variable	0.16
(32°20'N; 64°40'W)	15 February	500	S	0.15
Georges Bank	10 April	105	NW	1.4
(41°40'N; 67°30'W)	13 April	675	NW	0.82
(41°40'N; 67°30'W)	15 April	660	NE	0.58
(41°40'N; 67°30'W)	17 April	655	NW	0.61
(41°40'N; 67°30'W)	19 April	640	SW	0.80
(41°40'N; 67°30'W)	21 April	650	SW	1.60
Vineyard Sound	13 April	105	SW	3.9
(41°20'N; 70°50'W)	30 April	224	SW	5.3
Grand Banks	25 June	780	SSW	0.05
(45°16'N; 52°08'W)	26 June	960	SW	0.07
(45°16'N; 52°08'W)	27 June	840	WSW	0.10
(45°16'N; 52°08'W)	28 June	940	WSW	0.16
(45°16'N; 52°08'W)	29 June	540	W	0.05

centers (Bermuda is equidistant from both northeast and southeast U.S. industrial complexes) the seaward decrease is actually exponential (Fig. 3). Similar observations have been made for wind transport of industrial particulates over the ocean (Folger, 1970) and bomb fallout over the United States after nuclear tests (Eisenbud and Harley, 1953).




Fig. 3. Relationship of atmospheric PCB and distance from industrial sources.

DISCUSSION

In our opinion the collection method used can only approximate the relative concentrations of chlorinated hydrocarbons in the particulate and vapor aerosol phase. Hydrocarbons in the vapor or aerosol phase will easily pass through the glass fiber filter and be trapped in the silicone oil at an efficiency similar to the spiking experiments. Hydrocarbons adsorbed on solid particles which are trapped by the filter still have equilibrium

vapor pressures with the surrounding air. At the high flow rates used these adsorbed hydrocarbons could be entrained in the flow and swept into the vapor trap. The process is similar to that utilized in gas-liquid chromatography. However, if we can assume that particulates are being collected throughout the sampling period, the observation that only 1 per cent or less of the total chlorinated hydrocarbon concentration in the sampled air is found on the filter argues strongly against significant particulate transport.

It seems likely that during the atmospheric lifetime of chlorinated hydrocarbons the vapor, aerosol and particulate states will equilibrate and exchange many times depending on temporal meteorological parameters. Thus, the state in which they happen to be captured at a particular geographical location is relatively unimportant to considerations of total atmospheric transport. Thus, the concentrations reported in Table 1 should be regarded as total PCB m^{-3} .

The strong dilution and fallout gradient observed over the western North Atlantic is not reflected in the surface water concentrations. However, since the highest concentrations of suspended matter are in coastal waters (Manheim *et al.*, 1970) adsorptive scavenging by organisms and sedimentation must deplete these waters of dissolved PCBs at rates comparable to the delivery. In contrast, only a small fraction of the discharged PCBs reach the open ocean, but that which does must remain dissolved in the surface waters for a long time because of the low concentration of suspended matter and the slower sedimentation rate. The result of these processes is that the concentration of dissolved PCBs from near shore water and the open sea is about the same (Harvey *et al.*, 1973). Sediments should reveal a gradient of PCBs from inshore to the abyssal plains. Analyses of bottom feeders support this prediction (Harvey *et al.*, 1974).

Since the relative unimportance of river transport of chlorinated hydrocarbons has been noted (Seba and Prospero, 1971; Hartung and Klinger, 1970; Corcoran and Lee, 1970; Seba and Corcoran, 1969) wind transport is the major source of chlorinated hydrocarbons in the oceans.

Acknowledgements: We thank Drs. R. Duce and T. Bidleman, University of Rhode Island for permission and assistance in using their Bermuda tower. Dr. M. Grosslein made space available aboard *R/V Albatross IV*, National Marine Fisheries Service. We are especially grateful to Mr. G. Medeiros for collecting the samples on *Albatross IV*. Finally, we are indebted to Mr. Neil Greer (Amoco Canada Ltd.) for making accommodations available on the *Sednerk I*. This work was supported by the International Decade of Ocean Exploration, National Science Foundation, Grant GX-35212.

REFERENCES

- Armour J. A. (1974) Quantitative perchlorination of polychlorinated biphenyls as a method for confirmatory residue measurement and identification. *J. Assoc. Off. Agric. Chem.* (in press).
- Corcoran E. F. and Lee C. C. (1970) The distribution of chlorinated hydrocarbon pesticides in the Mississippi River Delta and adjacent Gulf of Mexico. Paper presented before American Chemical Society, 4 December, New Orleans, Louisiana.
- Eisenbud M. and Harley J. H. (1953) Radioactive dust from nuclear detonations. *Science* 117, 141-147.
- Folger D. W. (1970) Wind transport of land derived mineral, biogenic and industrial matter over the North Atlantic. *Deep Sea Res.* 17, 337-352.
- Hartung R. and Klinger G. W. (1970) Concentration of DDT by sedimented polluting oils. *Environ. Sci. Technol.* 4, 407-410.
- Harvey G. R., Steinhauser W. G. and Teal J. M. (1973) Polychlorobiphenyls in North Atlantic Ocean water. *Science* 180, 643-644.
- Harvey G. R., Miklas H. P., Bowen V. T. and Steinhauser W. G. (1974) Observations on the distribution of chlorinated hydrocarbons in Atlantic Ocean organisms. *J. Mar. Res.* (in press).
- Holden A. V. (1970) Source of polychlorinated biphenyl contamination in the marine environment. *Nature* 228, 783-784.

- Manheim F. T., Meade R. H. and Bond G. C. (1970) Suspended matter in surface waters of the Atlantic continental margin from Cape Cod to the Florida Keys. *Science* 167, 371-376.
- Risebrough R. W., Huggett R. J., Griffin J. J. and Goldberg E. D. (1968) Pesticides: Trans-Atlantic movements in the northeast trades. *Science* 159, 1233-1235.
- Risebrough R. W. (1972) Personal communication.
- Seba D. B. and Corenran E. F. (1969) Surface slicks as concentrators of pesticides in the marine environment. *Pest. Monit. J.* 3, 190-193.
- Seba D. B. and Prospero J. M. (1971) Pesticides in the lower atmosphere of the northern equatorial Atlantic Ocean. *Atmospheric Environment* 5, 1043-1050.