Bird killer could be a chemical



Peter Davis

Cause of death unknown: a technician at the Nature Conservancy Experimental station examines a dead guillemot

WHAT killed 10,000 sea birds in the northern Irish Sea during the past three weeks, will probably never be known. Scientists from many laboratories and organisations who met in London last week, concluded that oil pollution and severe gales could have played a part in the wildlife disaster.

But the most disturbing results to come out of the inquiry were the high levels of industrial chemicals called polychlorinated biphenyls found in eight out of nine birds examined at the Nature's Conservancy's station at Monk's Wood. The hundreds of parts per million of PCB's in the birds were almost certainly enough to be toxic, possibly to be lethal.

Last November, The Sunday Times published an article on the possible risk to birds from this widly-used group of industrial chemicals. Their presence in dead birds was discovered in the first place purely by chance, for they happen to be similar to organochlorine pesticides and were identified by the same analytical procedures.

There is no guarantee that even if poisoning is suspected, analysis of a dead bird will reveal the substance responsible. Analysts can only find what they are looking for, and if some unknown poison is involved, the chances that it will identified are slim.

The organophosphorus insecticides, for example, which are closely related to nerve gases, cannot be detected in dead animals, and only with great difficulty in live ones. Because of the shortage of live specimens no tests for these poisons have yet been carried out, but the scientists hope to do so in the near future.

If nothing else the disaster has exposed the lack of any kind of organisation for dealing with incidents of this kind, a lack which was even more glaring at the time of the Torrey Canyon. It was only informal contact between scientists that ensured a reasonably thorough investigation.

At all stages it has been held up by lack of specimens, lack of people and lack of money. To analyse a single bird for organochlorine pesticides costs from £10-£20. It seems a trivial sum, but to a laboratory with modest resources and other work to do, it is far from negligible.

Bryan Silcock

Sunday Times 26/10/69.

DSW 282778

STLCOPCB4064107