

THE BRITISH JOURNAL OF CANCER

(The official journal of the British Empire Cancer Campaign)



EDITOR
R. W. Scarff

ASSISTANT EDITOR
F. R. Selbie

ADVISORY EDITORS

S. P. Bedson
Sir Stanford Cade
E. C. Dodds
Malcolm Donaldson
C. E. Dukes
L. H. Gray
W. E. Gye
A. Haddow

The Lord Horder
J. P. Lockhart-Mummery
J. McIntosh
W. D. Newcomb
Sir Heneage Ogilvie
M. J. Stewart
Percy Stocks
B. W. Windeyer

DIRECTOR
F. L. Hopwood

VOLUME ONE
1947

LONDON
H. K. LEWIS & CO. LTD.
136 COWER STREET, W.C.1

THIS DOCUMENT WAS NOT A RECORD OF
PPG INDUSTRIES, INC. DID NOT COME FROM
IT'S FILES AND CANNOT BE AUTHENTICATED
BY PPG INDUSTRIES, INC.

PP.274-5

GG 15249

1 BB 0007240 1

EX 10

NOTE: THIS DOCUMENT DID
NOT GET INTO THE FILES

A FURTHER STUDY OF THE INCIDENCE OF CANCER OF THE LUNG AND LARYNX.

E. L. AND N. M. KENNAWAY.

(From the Pathological Department, St. Bartholomew's Hospital, London.)

Received for publication July 25, 1947.

MATERIAL EXAMINED.

THE data given below are derived from an examination of the death certificates of cases of cancer of the lung and of the larynx in males from England and Wales for the years 1921 to 1938 inclusive. The figures for the first 12 years of this period (1921 to 1932) were dealt with in an earlier paper (Kennaway and Kennaway, 1936); the material from the subsequent six years has now been examined in the same way, and the data for the whole 18-year period 1921-38 can now be presented together. The whole number of certificates examined individually was 38,418, while the annual total figures for cases in women have been utilized for comparison. This total was made up as follows:

Certificates Examined.			
Males.	1921-32.	1933-38.	Total.
Cancer of lung	8808	14,741	23,549
Cancer of larynx	9472	5,397	14,869
			38,418

The investigation was carried out on the general lines followed in an earlier study of the occupational incidence of cancer of the bladder and prostate (Henry, Kennaway and Kennaway, 1931).

The chief objects of the enquiry were: (1) to ascertain whether cancer of the lung showed any special incidence upon particular occupations, and (2) to seek for evidence to show whether the increase in deaths attributed to cancer of the lung, now of the order of over 16-fold (from 361 deaths of males in 1921 to 5982 in 1945) indicates any real increase in the prevalence of this form of cancer (Fig. 1). The figures for cancer of the larynx serve as a basis of comparison with those for cancer of the lung, in that the larynx is a part of the same (respiratory) tract which is more accessible to inspection, and is not examined by diagnostic methods which have changed greatly in recent years. The death certificates of persons under 20 years of age were omitted, as the investigation was concerned primarily with the effect of occupation.

Method of Calculation of Ratios.

The comparative incidence of cancer of the lung and larynx upon various occupations was estimated in the manner described in detail in the earlier paper (Kennaway and Kennaway, 1936). In judging of the occupational incidence of

GG 15250

1 BB 0007241 1

UNIVERSITY OF MICHIGAN

NOTE: THIS DOCUMENT DID
NOT COME FROM PPG FILES

incidence of cancer of the lung is the same in silicotic and non-silicotic (Table XI).

The account given here of the literature on the relation of silicosis to cancer of the lung does not claim to be complete (see the review by Klotz, 1955).

(6) *Asbestos workers.*

In the years 1939 to 1944 in England and Wales the fatal cases of cancer of the lung and asbestosis numbered 2699 and 75 respectively. The average duration of employment in fatal cases was, in silicosis, 34 years, and in asbestosis 15 (Factories Report, 1944). In his report for 1953, the Senior Medical Officer of Factories states that "among 103 fatal cases in which asbestosis or asbestosis with tuberculosis was present, cancer of the lung was associated in 12 (11.6 per cent)." Table XII records the cases of cancer of the lung in asbestos workers which have been found in the materials dealt with in this paper.

TABLE XII.—*Cancer of Lung. Asbestos Workers. Males. Death Certificates 1932-1938.*

	Age.
1935 Asbestos weaver	62
1935 " cloth weaver	54
1938 " weaver	46
1938 Boiler and pipe asbestos coverer	58
1938 Asbestos packer	56
1936 Disintegrating room hand in asbestos works	65
1934 Departmental Manager in asbestos company	45
1936 Asbestos works manager	58

These eight cases of cancer of the lung occurred in the years 1934 to 1938 among men of ages from 45 to 65. If one takes the number of deaths from cancer of the lung in these years among the general population of males of approximately the same age groups (45-64) one can make the following comparison:

- 4,068,382 males (ages 45-64) produced 8191 deaths from cancer of the lung.
- = approximately 4 million males (ages 45-64) produced 8000 deaths from cancer of the lung.
- = 4000 males produced 8 deaths from cancer of the lung.

The number of male asbestos workers in this age-group appears to be considerably less than 4,000 but as we have been unable to obtain any exact figures for the workers at risk one cannot carry the calculation further.

Lynch and Smith (1935) report from South Carolina a bronchiogenic carcinoma in a man who had been for 21 years an asbestos weaver.

Egbert and Geiger (1936) record a bronchiogenic carcinoma, with characteristic pulmonary fibrosis and asbestosis bodies, in a man who had been for 18 years an asbestos weaver at New Haven, Connecticut.

Gloyne (1935) records two cases of fairly advanced asbestosis, with squamous carcinoma of the lung, in women workers in England. One survived nine years after an eight years' exposure to asbestos dust as a spinner.

GG 15251

1 BB 0007242 1

University of Pittsburgh

other lived for fifteen years after two short periods of six months and thirteen months' exposure in the mangle and opening departments of the factory. . . . The malignant lesions of the lung were, in each case, very small and not recognized during life." Subsequently the patient died with asbestosis and an oat-cell carcinoma of the lung in a man who had been for 20 years foreman in the stores department of an asbestos works.

(7) *Alcohol and tobacco.*

There is a considerable difference between occupations associated with the supply of alcohol, and of tobacco: the former show a very high incidence upon the larynx, and a normal incidence upon the lung (Table XIII). The ratio for the lung is the same for licensed victuallers as for physicians and surgeons (Table XV). The occupations of barman and cellarman show by far the highest figures of any in Table V for cancer of the larynx. The incidence of cancer of the lung upon the two occupations connected with tobacco has decreased distinctly in the later period. The increasing use of cigarettes (Fig. 10, 11) has caused the tobacconist to become more and more a vendor of closed packets which can hardly have any occupational effect. The decline of pipe-smoking makes the mixing by the tobacconist by hand of a blend to suit a customer's taste to be less frequent.

TABLE XIII.—*Cancer of Lung and Larynx. Occupations Associated with the Supply of Alcohol and Tobacco.*

	Ratio.			
	Lung.		Larynx.	
	1921-32.	1933-38.	1921-32.	1933-38.
Barmen	104	122	469	275
Cellarmen	92	87	317	502
Licensed victuallers	128	124	185	199
Tobacco manufacturers	196	110	65	129
Tobacconists and their assistants	175	82.5	142	171

(8) *Coal-gas and tar.*

The occupations associated with coal-gas and tar are important on account of the known liability of the workmen to cancer of the skin, and of the bladder (Henry, Kennaway and Kennaway, 1931), but they offer unsatisfactory statistical material because some of the most important occupations employ very few men, and hence produce but few cases of cancer, so that the sampling error is high even over an 18-year period. This difficulty can only be met by omitting some of the smaller classes. Thus, in Table XIV there are ten occupations in which there is especial exposure to coal-gas and tar. If from these ten one excludes the four which have produced less than ten deaths from cancer of the lung, one obtains, by this quite arbitrary procedure, the results shown in Table XIV.

Thus all the six occupations retained, which employ about 82,000 men (1931 census) and yield 267 cases of cancer of the lung, have an increased (up to 284) ratio for this disease, the lowest figure being that for chimney sweeps (119).

GG 15252

BB 0007243

NOT COME FROM PG FILES

Kennaway-
Mixed Pages 1947

INCIDENCE OF CANCER OF LUNG AND LARYNX

261

any form of cancer it is of course necessary to correct for age distribution, for an occupation employing a large proportion of older men will yield more cases of




FIG. 1.—Deaths from cancer of lung and larynx. Males. 1921-1945.

cancer apart from any aetiological factor. The census returns give the age distribution, in 5- and 10-yearly periods, of the whole population of males, and

GG 15253

BB 0007244

NOTE: THIS DOCUMENT DID
NOT COME FROM THE FILES

also of those following each one of the recognized occupations, at the time of the census. The number of deaths attributed to cancer of the lung and larynx occurring in the whole male population in each of these age groups during the years in question was obtained from material at the General Register Office. The comparison of the various occupations with the general population is then a sum in proportion, which gives the ratio of the number of deaths in any given occupation to those, reckoned as 100, which occurred in the same number of males in the same age groups in the whole population. An example of this calculation is given in the earlier paper.

Sources of Error in Statistical Work on Death Certificates.

These were examined in the earlier paper (Kennaway and Kennaway, 1936). The use of death certificates for statistical purposes is liable to errors which are inherent in the nature of the material and cannot be wholly eliminated. One must either recognize and admit these errors and correct them in any way possible, or abandon any attempt to obtain information from these data.

(1) *Correction for period between census.*—In order to compare the incidence of cancer upon different occupations one must know the number of persons following each occupation. This is learned from the census returns. The longer the inter-census interval the greater is the risk of important alteration in these numbers. The figures for populations used for the earlier period (1921-32) were those of the 1921 and the 1931 census, which were combined by means of a formula described in the earlier paper (Kennaway and Kennaway, 1936): those in the present paper are from the 1931 census, which is the last taken. No census was made in 1941, on account of the war.

The ratios for the whole 18-year period are calculated from the sums of the registered, and the calculated, deaths for the 12-year and 6-year periods as in the following examples:

TABLE I.—Combination of 12-year and 6-year Periods. *Cancer of the Lung.*

	1921-1932			1933-1938			1921-1932 + 1933-1938		
	Calculated deaths.	Total registered deaths.	Ratio of regd. to 100 calculated deaths.	Calculated deaths.	Total registered deaths.	Ratio of regd. to 100 calculated deaths.	Calculated deaths.	Total registered deaths.	Ratio of regd. to 100 calculated deaths.
Carpenters.	169	123	73	276.6	200	72	445.6	323	72.5
Printers	39.7	61	102	88.8	116	131	148.5	177	119

(2) *Sampling error.*—In some of the most interesting occupations from the present standpoint (e.g. patent fuel workers, tar distillers) a few thousand men only are employed, and these will yield but very few cases of cancer of any one organ even if the incidence is high (Table XIV). Hence the sampling error will be large even over the period studied here of 18 years. The figures given in Table XVIII show the range of fluctuations which occurs. Even the large group of agricultural workers, numbering over 900,000 men, does not give so smooth a curve representing deaths from cancer of the lung as does the whole male population (Fig. 2).

BB 0007245

GG 15254

Factors which may affect the value of the individual death certificate are:

(3) *Location of employment.* The individual entry in the death register does not give, and is not intended to give, any information about the length of




FIG. 2.—Deaths from cancer of lung. Males.

A. General population. 1921-1944.

B. Coal miners.

C. Agricultural workers. } $\times 20$. 1921-1938.

time during which the deceased followed the occupation named, nor about the nature or duration of any previous employment; the uncertainty could be removed only by enquiry into each individual case, and that is impracticable

BB 0007246

GG 15255

NOTE: THIS DOCUMENT
NOT COPY

when large numbers are concerned. The immense amount of labour involved in such enquiries can be seen in the monograph on cancer of the scrotum by Henry (1946). But as none of the occupations included in the present study, with one exception, shows a very high incidence of cancer of the lung, comparable to that seen in the occupational cancers of the skin and bladder, this source of error is not a very serious one.

(4) *Incorrect diagnosis* is of course a source of error in these, as in all other, medical data. In the case of death certificates received in bulk, perhaps some years after the dates of the deaths recorded, one can do nothing but exclude any certificate of which the wording suggests (1) that the cause of death did not come within the category under examination, or (2) that the diagnosis was uncertain or ill-founded (see next section: such certificates are made the subject of individual inquiry by the General Register Office). When such exclusions have been made, one must treat the remaining certificates as correct if any use is to be made of the material.

Death Certificates Taken for Examination.

The material utilized in this paper has been restricted as far as possible to cases of primary malignant growth of the lung, and of the larynx. The forms of words in the certificates taken for examination, and in those rejected, may be stated thus:

Cancer of lung.

(a) *Retained.*—Cancer, carcinoma, or sarcoma of lung, bronchus, pleura, root of lung, hilum of lung, lung and mediastinum, or lung and pleura. Pulmonary, or bronchial cancer, carcinoma or sarcoma. Cancerous pleurisy.

(b) *Rejected.*—Any new growth of mediastinum, of mediastinal or of bronchial glands. New growths of chest or thorax. Peribronchial new growth. Any new growth qualified by "probable," "doubtful," or "query."

Where another organ is mentioned, in addition to the lung, as the site of new growth, the practice has been to retain those cases where this organ is unlikely to be the primary site (e.g. cancer of lung and brain, or of lung and liver) and to reject the rest (e.g. sarcoma of arm and lung, sarcoma of lung and testicle, cancer of lung and oesophagus).

Cancer of larynx.

(a) *Retained.*—The category of cancer of the larynx in the tabulation used in the General Register Office includes malignant growths of which the situation is described by the following terms:

Aryteno-epiglottidean folds.	Post-crioid, retro-crioid.
Arytenoid glands and cartilage.	Pyriform fossa. Sinus pyriformis.
Cricoid.	Thyroid cartilage.
Epiglottis.	Vallecula epiglottica.
Glossio-epiglottic fold.	Vocal cords.
Glottis.	Trachea, windpipe.

All these were retained with the exception of vallecula.

University of Manchester

GG 15256

1 BB 0007247 1

Some cases of which the nature is doubtful (e.g. "tumour of cricoid cartilage and thyroid gland").

Autopsies.

The opinion is sometimes expressed that statistics of lung cancer are of no value which comprise cases where no autopsy has been made, and some authorities (see below, p. 289) maintain further that, even when autopsies are performed, many cancers of the lung escape notice. Table II shows the numbers of autopsies recorded in the death certificates for cancer of the lung* discussed here (i.e. after the exclusions stated in the preceding section). Similar data for 1921-32 are given in the earlier paper (Kennaway and Kennaway, 1936). These cases are classified as follows:

- No P.M. : (1) No P.M.
(2) Inquest. No P.M.
P.M. : (3) P.M.
(4) Inquest and P.M.
(5) Coroner's P.M. without inquest.

TABLE II.—Autopsies. Cancer of Lung. (Males, 1933-38.)

Year.	I. No P.M.	II. Inquest. No P.M.	III. Sum of I and II.	IV. P.M.	V. Inquest and P.M.	VI. Coroner's P.M. No Inquest.	VII. Sum of IV, V and VI.	Total.
1933	1,247	6	1,253	439	34	28	501	1,754
1934	1,784	3	1,487	494	22	39	555	2,042
1935	1,646	—	1,646	574	36	34	644	2,290
1936	1,958	4	1,962	539	37	34	610	2,572
1937	2,235	5	2,240	522	61	40	623	2,863
1938	2,524	5	2,529	598	42	53	693	3,222
Total	11,094	23	11,117	3,166	232	228	3,626	14,743
1921-32	6,569	34	6,603	2,005	132	68	2,205	8,808
1921-38	17,663	57	17,720	5,171	364	296	5,831	23,551

TABLE III.—Proportion of Autopsies. (Males, 1921-38.)

		Total deaths.	Autopsies.	Mean autopsies.	
				Per annum.	Per cent of deaths.
Cancer of lung,	1921-32	8,803	2,205	184	25.0
"	1933-38	14,743	3,626	604	24.6
" of larynx,	1921-32	2,472	995	50	6.3
"	1933-38	3,398	448	75	8.3

Table III shows (1) that the number of autopsies has increased much more in the lung series than in the larynx series; and (2) that the average proportion of cases of cancer of the lung examined post mortem is the same in the earlier

* A corresponding table for cancer of the larynx is omitted to save space.

GG 15257

BB 0007248

UNIVERSITY OF MICHIGAN

266

TABLE IV.—Cancer of Lung. 1921-1932 and 1933-1938.

The occupations are arranged in descending order of the ratio of registered to 100 calculated deaths, 1921-38.

Occupation.	1921-1932.				1933-1938.				1921-38.	
	Population.	Calculated deaths.	Total regd. deaths.	Ratio of regd. to 100 calculated deaths.	Population.	Calculated deaths.	Total regd. deaths.	Ratio of regd. to 100 calculated deaths.	Ratio of regd. to 100 calculated deaths.	Ratio of regd. to 100 calculated deaths.
Labourers, patent fuel works	847	0.40	2	433	00	0.005	1	1,529	571	
Gas stokers and coke-oven chargers	12,818	10.8	37	342	13,620	19.1	48	251	284	
Gas producer men	3,195	2.5	4	162	2,451	3.4	8	232	202	
Metal grinders	15,220	8.3	19	220	17,000	13.9	20	141	176	
Gas works foremen and inspectors	6,443	5.1	10	197	6,076	9.3	15	161	174	
Boaters' mill workers; ship makers and joiners	1,486	1.04	1	96	1,760	1.8	4	218	174	
Miscellaneous labourers, road sweepers and dustmen	39,155	30.2	68	173	40,184	61.8	103	107	169	
Gas fitters	15,079	8.8	16	182	19,376	15.8	25	158	167	
Painters, street masons, concretors, asphalters	7,419	5.7	12	209	8,915	10.2	14	137	163.5	
Motor drivers, goods and passengers	254,350	96	136	142	305,805	196.9	301	153	119	
Electric furnace engine men	965	6.74	1	131	398	0.05	1	153	119	
Cabinet makers	31,155	24.7	34	138	41,116	39.4	60	152	147	
Tobacco manufacturers	7,698	4.0	0	100	8,192	7.2	9	116	114	
Stomachians, cutters and dressers	37,177	35.5	53	149	40,087	60.8	85	140	113	
Tanners, leather dressers, curriers, skilled workers	18,001	14.2	21	148	18,580	22	30	136	141	
Potters; ware-makers, casters and finishers	7,723	5.9	9	152	8,511	9.9	13	132	139.5	
Gas works engine and crane drivers	1,854	1.7	1	60	1,809	2.7	5	186	138	
Sandblasters	920	0.5	1	201	1,285	0.97	1	103	136	
Gas works managers	1,303	1.2	3	214	1,049	1.7	1	58	136	
Drivers of horse-drawn vehicles	151,554	123	177	144	138,556	183.5	210	131	136	
Gas works labourers	33,391	23.5	46	162	33,909	40	50	109	129	
French polishers	17,783	11.4	19	167	22,072	19.7	21	107	129	
Painters	178,837	143	176	123	202,710	245.5	323	132	129	
Licensed victuallers	75,832	83.8	167	128	76,440	138.3	172	124	126	
Printers	83,489	59.7	61	162	82,147	88.8	116	131	119	
Chimney sweeps	5,900	0.5	11	170	6,190	11.1	10	66	119	
Flourishers	19,050	31.3	33	165	60,301	53.9	80	123	116	
Barnmen	21,059	9.9	10	161	27,161	10.1	20	122	115	
Tobaccoists and their assistants	11,031	12	21	175	11,075	21.2	17	82.5	112	

NOTE: THIS DOCUMENT DID NOT COME FROM PRO FILES
F. L. AND N. M. KENNEDY

66 15758
1 BB 0007249 1

TABLE IV (contd.).

Occupation.	1921-1932.				1933-1948.				1921-34.	
	Population.	Calculated deaths.	Total real. deaths.	Ratio of real. to 100 calculated deaths.	Population.	Calculated deaths.	Total real. deaths.	Ratio of real. to 100 calculated deaths.	Ratio of real. to 100 calculated deaths.	Ratio of real. to 100 calculated deaths.
Hairdressers	36,197	23.0	20	84	44,430	43	47	109	101	
Bakers and pastry cooks	63,617	43.7	37	85	75,010	73.8	70	103	86	
Professional men	103,097	119.3	101	81	108,912	181.3	176	97	96	
Collar-men	6,333	1.3	1	92	7,050	6.9	6	87	89	
Potters, etc., kiln and ovenmen, kiln setters and piers	12,188	8.2	8	98	11,112	14.6	12	82.6	84	
Groomers, horsekeepers	31,918	28.4	30	106	29,521	42.6	30	71	83	
Black kiln and ovenmen	6,225	4.3	1	23	7,112	7.4	0	121	85	
Stationary engine and crane drivers	72,875	58.0	52	71	69,680	61.0	75	82	78	
Cotton strippers, grinders and card-room jollers	5,242	3.4	1	30	5,312	5.8	0	101	76	
Locomotive engine drivers, firemen and cleaners	87,514	54.1	41	76	87,812	91.1	68	73.5	71	
Carpenters	210,131	109	123	73	243,928	276.6	200	72	72.5	
Stone miners and quarrymen	28,471	21	22	105	32,380	31.0	17	53	70	
Coal miners, hewers and jollers	419,311	276	174	63	448,813	442.7	310	72	69	
Lithographic and process engravers	5,701	3.0	3	83	7,280	6.0	4	64	67	
Blind turnmen and labourers	9,391	7.2	6	84	6,135	7.9	4	50	66	
Gardeners	200,071	201	118	58	223,310	347.6	213	61	60	
Cotton spinners and piecers (mule, ring cap or flyer)	38,817	21.5	7	32	43,991	30.1	27	73	59	
Coal miners- workers above ground	83,671	65.1	46	71	90,056	101.0	49	47	50	
Tar distillery workers and coke-oven workers	9,697	6.8	2	29	7,824	8.5	0	70	52	
Brick and plain tile makers	9,825	6.2	3	40	12,403	10.5	5	47	48	
Coal miners, other workers below ground	81,403	65.2	28	44	80,247	112.4	52	46	45	
Coal miners, conveying material to the shaft	91,561	31.4	12	38	127,170	80.2	28	47	41	
Coal miners, making roads	62,156	64.8	20	53	69,512	100	37	37	43	
Cotton weavers	40,078	27.8	13	47	41,016	46.2	17	37	40.5	
Farmers	263,180	286	153	53	260,062	460.5	148	32	40	
Agricultural labourers, including sheep-herds	434,300	351	119	34	453,221	532.7	137	26	29	
Farm bailiffs and foremen	20,270	18.8	5	26	17,983	27.1	6	22	21	
Total	56		2,249		3,550					

INCIDENCE OF CANCER OF UTERUS AND LARYNX

267

CG 15259

BB 0007250 1

NOTED BY CURRENT DID
NOTED BY FILES

TABLE V.—*Cancer of Larynx. 1921-1932 and 1933-1938.*

The occupations are arranged in descending order of the ratio of registered to 100 calculated deaths, 1921-38.

Occupation.	1921-1932.				1933-1938.				1921-38.	
	Population.	Calculated deaths.	Total regl. deaths.	Ratio of regl. to 100 calculated deaths.	Population.	Calculated deaths.	Total regl. deaths.	Ratio of regl. to 100 calculated deaths.	Ratio of regl. to 100 calculated deaths.	Ratio of regl. to 100 calculated deaths.
Barmen	21,059	8.3	39	469	27,181	4.7	13	276	396	
Cellar-men	6,333	4.4	14	317	7,050	2.4	12	502	382.5	
Gas-works engine and crane drivers	1,854	1.7	4	233	1,800	0.93	2	214	227	
Gas stokers and coke-oven chargers	12,818	10.7	20	186	13,026	6.7	17	255	213	
Mainly casual labourers, road sweepers and dustmen	39,455	45.3	78	172	40,184	23.6	43	267	205	
Licensed victuallers	75,832	93.4	173	165	76,310	61.2	102	190	190	
Metal grinders	16,220	7.1	16	141	17,009	3.8	9	234	174	
Drivers of horse-drawn vehicles	154,554	132	212	184	138,656	69.8	100	143	100.5	
Chimney sweeps	5,900	7.8	15	102	6,496	4.6	6	120	169	
Potters; wine-makers, caskers and butlers	7,723	6	11	184	8,511	3.4	4	118	100	
Potters, etc.; kiln and ovenmen, kiln setters and placers	12,188	7.7	9	110	14,442	4.6	10	220.5	150	
Tobaccoists and their assistants	14,031	13.4	10	112	14,075	8.2	14	171	163	
Cabinet makers	31,455	28	40	164	41,116	15.7	20	127	151	
Painters, street masons, concretors, asphalters	7,419	6.8	9	154	8,945	3.4	4	116	140	
French polishers	17,783	11.1	15	135	22,072	6.4	9	141	137	
Lithographic and process engravers	6,701	3.6	3	84	7,289	2.3	5	214.5	135	
Painters	178,837	150	195	130	202,746	87	120	138	133	
Cotton spinners and piecers (mule, ring, cap or flyer)	38,817	20.9	31	148	43,991	12.4	11	91	127	
Groomers, horse-keepers	31,948	32.1	43	134	29,521	17	17	100	122	
Plumbers	49,050	31.1	36	116	69,301	18.3	22	120	117	
Printers	83,489	101.2	71	116	82,147	30.3	33	109	111	
Gas fitters	35,079	8.3	8	66	19,375	5	6	120	106	
Motor drivers, goods and passengers	254,350	64.8	65	100	365,805	42.5	48	113	105	
Hairdressers	36,497	22.0	28	124	41,439	13.5	10	74	105	
Gas-works managers	1,303	1.3	1	76	1,619	0.63	1	157	162	
Bakers and pastry cooks	63,547	46.8	49	105	75,016	27.6	24	87	98	

E. L. AND N. M. KENNEDY

NOTE: THIS DOCUMENT DID NOT COME FROM PROFILES

1 BB 0007251 1

CG 15260

44-38861-10002

iii

205

66 15261

NOTE: THIS DOCUMENT DID
NOT COME FROM PPG FILES

and later periods. Hence the continuing increase in deaths attributed to cancer of the lung is not due to any increase in this proportion.

COMPARATIVE INCIDENCE UPON VARIOUS OCCUPATIONS.

In the sections below, the periods 1921-32, 1933-38 and 1921-38 are referred to as A, B and C respectively.

The 56 occupations named in Table IV employed during the period B 3,880,001 out of the 13,305,668 males aged 20 and upwards (census of 1931), and they include most of the men engaged in the two largest industries in the country, namely agriculture and coal mining. The tables include 3550 out of the 11,741 cases of cancer of the lung, and 1451 out of the 5397 cases of cancer of the larynx, occurring in period B, in the whole population in question.

Tables IV and V show the data referring to the occupations which have been studied in detail. In Table IV the occupations are arranged in the order of magnitude of the ratio for cancer of the lung, and in Table V in the corresponding order for cancer of the larynx.* The population engaged in each occupation, and the numbers of calculated, and of actual, deaths which are required for the estimation of the ratios are included. The number of deaths in each year of period B in each of these occupations is given in Table XVIII.

(1) *Agriculture.*

A group of open-air occupations (gardeners, farmers, agricultural labourers, including shepherds, farm bailiffs and foremen), which together include about 954,000 men, has 503 cancers of the lung and 342 cancers of the larynx in B. This group shows very low ratios for lung (from 26 to 58 in A and from 22 to 61 in B) and low ratios for larynx (from 41 to 73 in A and 29 to 79 in B). In this open-air class the highest figure for both lung and larynx is given by gardeners, of whom many live in or near large towns.

(2) *Coal mining.*

The chief coal-mining occupations (workers above ground, hewers, road-makers, persons conveying material to the shaft, other workers below ground), which employed nearly 770,000 men, show low ratios (lung, 38 to 71 in A and 37 to 72 in B; larynx, 44 to 58 in A and 42 to 73 in B), which are very like those of the open-air group considered in the preceding paragraph. The air in a coal mine is pumped in from outside, and collieries are not always situated in urban districts (e.g. the Kent coalfield). The close agreement in the ratios given in A and B by the agricultural and coal mining groups is noteworthy.

These two industries (agriculture and coal mining) have also a low incidence of cancer of the bladder (Henry, Kennaway and Kennaway, 1931). The rate of increase of deaths attributed to cancer of the lung in these two groups is shown in Fig. 2.

* The difference between the total numbers of occupations in Table IV (56) and Table V (52) is due to four occupations (patent fuel workers, potters' mill workers, blast furnace engine men, sandblasters) in which no deaths from cancer of the larynx occurred in either period.

GG 15262

BB 0007253

NOTE: THIS DOCUMENT DID
NOT COME FROM PPG FILES

ing of open-air occupations, where the cancer came to the dust of roads, is above the normal for the lung and larynx, with the exception that motor drivers have nearly normal figures for larynx (Table VI). None of these occupations shows a high incidence of cancer of the lung, and the pavloirs, street masons, concretors and asphalters show a distinct fall in the later period.

(4) *Grooms, horsekeepers, carpenters.*

Grooms and horsekeepers, who are exposed to various forms of dust arising from horses and fodder, have an incidence of cancer of the lung (A 106, B 71) and larynx (A 134, B 100) which is not very far from normal. Carpenters (Table I and Fig. 7) also show low ratios (lung A 73, B 72; larynx A 82, B 86). These instances suggest that exposure to dust *per se*, irrespective of its chemical nature, does not conduce to cancer of the lung.

TABLE VI.—*Cancer of Lung and Larynx. Occupations with Exposure to Road Dust.*

	Ratio.			
	Lung.		Larynx.	
	1921-32.	1933-38.	1921-32.	1933-38.
Pavloirs, street masons, concretors and asphalters	209	137	154	116
Council labourers, road sweepers, dust-men	173	167	172	267
Drivers of horse-drawn vehicles	144	131	184	143
Motor drivers (goods and passengers)	142	153	100	113

(5) *Silicosis.*

The occupations in which there is the greatest liability to silicosis are named in Table VII. Of these occupations, those mentioned in Table VIII, and the various subdivisions of coal-mining, were included in the present study. Table VIII shows that in the period 1933-38, the highest incidence of cancer of the lung is in potters' mill workers, slip makers and arkmen (ratio 218), but in this occupation the numbers are small (see Table IV), and hence the sampling error is considerable. In the other occupations in Table VIII the tendency is towards a decrease in the ratio. The stonemasons, and stone miners and quarriers of course include many workers with non-siliceous rocks. The figures for deaths from silicosis in occupations of this class are falling; in the metal grinders this is due to the adoption of non-siliceous abrasive wheels, which has been advocated by the Senior Medical Inspector of Factories (Factories Report, 1938). In the various groups of coal miners (Table IV) the low ratios show no uniform tendency to rise or fall. The general indication of these results is that the factors which lead to silicosis are not very active in producing cancer of the lung or larynx. There are, however, in the literature various accounts of cases in which

* C. J. Girdling (*Lancet*, 1946, 251, 891), in a study of pneumoconiosis in South Wales anthracite miners, found no cases of cancer of the lung in 227 necropsies on certified silicotics, and one case associated with silicosis in a further series, making nearly 400 necropsies in all.

GG 15263

BB 0007254

cancer of the lung was associated with silicosis. Such individual observations are difficult to assess statistically as the two conditions must sometimes occur together, and the close examination of silicotic lungs in special investigation favours the detection of a high proportion of carcinomas.

TABLE VII.—Deaths from Silicosis in England and Wales, 1939 to 1945 inclusive. (From the 'Annual Report of the Chief Inspector of Factories for the Year 1945,' p. 79.)

1. Refractories industries	68
2. Pottery, manufacture of	310
3. Sandstone quarrying and dressing	173
4. Sandstone masons	363
5. Metal grinding	98
6. Sandblasting	53
7. Steel dressing and cleaning of castings	45
8. Stone, pebble, flint and sand crushing	15
9. Scouring powders, manufacture of	5
10. Abrasive wheel manufacture	4
11. Glass cutting and bevelling	3
12. Millstone dressing	1
13. Slate quarrying and dressing	45
14. Granite quarrying and dressing	13
15. Tunnel mining (sewage works, etc.)	21
16. Coal mining	1731

TABLE VIII.—Cancer of Lung in Occupations with Liability to Silicosis.

	Ratios of registered to calculated deaths.			Number of deaths.
	1921-32.	1933-38.	1921-38.	
Potters' mill workers; slip makers and arkmens	96	218	174	5
Potters; ware-makers, casters and finishers	152	132	130.5	22
Metal grinders	229	144	176	39
Stonemasons, cutters and dressers	149	140	143	86
Stone miners and quarriers	105	49	70	39
Sand-blasters	201	103	136	2
Coal-miners (see Table IV)	58-71	37-72	43-69	774

Some of the literature on the relation of silicosis to cancer of the lung is summarized below.

Dible (1934), in the course of 14 autopsies upon persons of both sexes suffering from silicosis, found malignant disease in three cases in stonemasons, as follows:

- Case 1. Carcinoma of bile-duct. Spheroidal-cell carcinoma of prostate.
- " 2. Oat-cell carcinoma of lung. Columnar-cell carcinoma of colon.
- " 3. Anaplastic carcinoma of lung.

NOTE: THIS DOCUMENT DID NOT COME FROM THE FILES

GG 15264

BB 0007255

Andersen and Dible (1938) have reported similar results in the lungs (A and B) showing primary carcinoma (Table V) of histological abnormalities of blood, the mean content of these being nearly 40 times that of A, while the mean content of the remaining 38 (D) of B was the same as that of A. Microscopically, C showed typical silicotic nodules in 7, areas of confluent fibrosis in 3, and fibrosis not more than suggestive of silico-sis in 2 of the 12 cases. Effects in A and D "did not in any sense reproduce the picture found in those organs in which silica was present in excess." The authors conclude that "a group of cases of pulmonary carcinoma exists in which the organs contain an excess of silica and show histological evidence of silicotic fibrosis. The conclusion, we think, is that the role of the silicosis is aetiological."

TABLE IX.—*Silicosis and Cancer of Lung (Klotz, 1939).*

	Number of autopsies.	Carcinoma of lung.	
		Number.	Per cent of autopsies.
Cases of silicosis	50	4	8.0
Unselected cases	4500	53	1.17

Klotz (1939) gives some data from Toronto (Table IX) of which the statistical character, and indications, are similar to those of the material of Andersen and Dible.

In the report for 1938 of the Chief Inspector of Factories (p. 81) the Senior Medical Inspector (Dr. J. C. Bridge) gave the following data derived from 1290 autopsies carried out to ascertain whether silicosis was present (Table X). Hence presumably the subjects had shown in life some respiratory symptoms.

TABLE X.—*Silicosis and Cancer of Lung (J. C. Bridge).*

Number of autopsies.	Silicosis.	Cancer of lung.	
		Number of cases.	Per cent of autopsies.
943	Present	23	2.4
347	Absent	17	4.9
1290			

TABLE XI.—*Cancer of the Lung in Witwatersrand Miners (Miners' Phthisis Medical Bureau).*

Europeans who had not worked underground (Miners' Phthisis Medical Bureau).			Non-silicotic European miners (Medical Bureau).			Silicotic European miners (Medical Bureau).		
Post mortems.	Cases of primary cancer of lung.	Percentage of all post mortems.	Post mortems.	Cases of primary cancer of lung.	Percentage of all post mortems.	Post mortems.	Cases of primary cancer of lung.	Percentage of all post mortems.
1296	13	0.93	1679	12	0.71	1435	10	0.7
			1458					
			3117					

The data obtained during 15 years from 3117 post-mortems on European miners on the Witwatersrand (Miners' Phthisis Bureau, 1936) indicate that the

GG 15265

BB 0007256

NOTE: THIS DOCUMENT DID
NOT COME FROM PPG FILES

276

E. L. AND N. M. KENNAWAY

TABLE XIV.—*Cancer of Lung. Occupations Associated with Coal Gas and Tar.*

	1921-32. Deaths.	Ratio.		
		1921-32.	1933-38.	1921-38.
Gas stokers and coke-oven chargers	85	342	251	284
Gas producermen	12	162	232	202
Gas-works foremen and inspectors	25	197	161	174
Gas fitters	41	182	158	167
Gas-works' labourers	96	162	109	129
Chimney sweeps	21	170	90	119
	267			
Excluded :				
Labourers, patent fuel works	3	433	1529	571
Gas-works engine and crane drivers	6	60	186	133
Gas-works managers	4	244	58	136
Tar distillery workers and coke-oven workers	8	29	70	52
	21			

In five of the six cases, and especially in sweeps, the ratio has fallen in the later period. The ratios for cancer of the larynx are irregular throughout this class.

(9) *Professional occupations.*

A group of professional workers (stockbrokers and stockjobbers ; clergymen ; Roman Catholic priests and monks ; ministers of other religious bodies ; judges, stipendiary magistrates and barristers ; solicitors ; physicians, surgeons and registered medical practitioners ; dental practitioners) numbering over 108,000 men and showing 176 deaths from cancer of the lung and 39 from cancer of the larynx, give altogether ratios rather below the average (lung A 94, B 97 ; larynx A 84, B 50 ; Tables IV, V and XV). These ratios for cancer of the lung, being near 100, are in accordance with Stevenson's (1923) discovery of the absence of social gradient in this form of cancer. When the individual occupations are examined considerable variations are found. In the earlier paper attention was drawn to the high ratio for cancer of the lung in judges, stipendiary magistrates and barristers (173) and in stockbrokers and stockjobbers (187) ; in the later period (1933-38) the former fell to 112 and the latter rose to 253. The stockbrokers and jobbers also have the highest ratio among the professional workers for cancer of the larynx. The figures for Roman Catholic priests and monks are too small to be significant. The ratio for ministers of other religious bodies has risen (Table XX). In regard to the view that the rapid increase in recorded deaths from cancer of the lung is due to the detection of more cases by improved diagnosis, the very moderate and consistent ratio (A 129, B 125) for cancer of the lung in medical men is noteworthy, for this is an occupation where the availability of the existing methods for the detection of cancer is presumably at a maximum. Dental practitioners give very similar figures (A 103, B 124).

BB 0007257 1

CG 15266

University of Liverpool

The ratios for cancer of the larynx in these occupations are low, and in some cases considerably below 100, except in stockbrokers and stockjobbers (158).

TABLE XV.—*Cancer of Lung. Professional Occupations.*

Occupation.	1921-32			1933-38			1921-38	1923-38	
	Population.	Total read. deaths.	Ratio of read. to 100 calculated deaths.	Population.	Total read. deaths.	Ratio of read. to 100 calculated deaths.	Ratio of read. to 100 calculated deaths.	Cancer of larynx.	Deaths. Ratio.
Stockbrokers and stockjobbers	6,893	14	187	7,122	32	353	229	8	158
Roman Catholic Priests and Monks	3,604	1	31	4,050	—	—	11	5	24
Clergymen (Anglican Church)	23,422	15	53	22,754	22	49	50	1	45.5
Ministers of other Religious Bodies	11,569	3	38	11,960	15	69	57	6	64
Judges, Stipendiary Magistrates and Barristers	3,658	7	173	3,632	7	112	136	1	34.5
Solicitors	17,306	19	100	17,853	26	84	90	9	65.5
Physicians, Surgeons and Registered Medical Practitioners	27,951	35	129	29,908	56	125	127	7	38
Dental Practitioners	10,494	8	103	11,633	18	124.5	117	2	44
									39

(10) *Cotton workers.*

Cotton strippers, grinders and card-room jobbers, among whom severe bronchitic and asthmatic affections occur, show a very low ratio (30) for cancer of the lung in A, which rose to the normal level (104) in B (see p. 286).

Cotton spinners and piecers have a low but increasing ratio for lung (A 32, B 75) and medium ratio for larynx (A 148, B 91). The total population of 43,991 (1931 census) includes ring spinners as well as mule spinners, but the latter occupation certainly makes up a very large proportion of the total, for the number of mule spinners aged 20 and over was estimated in 1926 as 41,000 by the Departmental Committee appointed to consider evidence as to the Occurrence of Epitheliomatous Ulceration among Mule Spinners. In view of the exposure to fumes which may be carcinogenic the low incidence of cancer of the lung is remarkable, but the very high incidence of cancer of the skin, and the increased liability to cancer of the bladder (Henry, Kennaway and Kennaway, 1931) in this occupation, must be taken into account. The question of the penetration into the lung of oil droplets in the air of the mule room will be dealt with in another publication.

RATE OF INCREASE OF CANCER OF THE LUNG AND LARYNX IN MEN AND WOMEN.

Cancer of lung.

Cancer of the lung has increased from 361 to 3982 cases per annum in men (Fig. 1) between 1921 and 1945, and from 189 to 1480 cases in women. These figures are in the ratio of 1:19.6 (men) and 1:8 (women). The increase became more rapid in 1926 and again in 1939 in men (Fig. 1) and about 1929 in women. The annual increase, in men, between 1932 and 1937 was at the rate of from 200 to 300 cases per annum; since then it has been very irregular

CG 15267

BB 0007258

NOTE: THIS DOCUMENT DID NOT COME FROM PPG FILES

(Table XVI). The ratio between deaths from cancer of the lung in men, and in women, is now (i.e. in 1945, the latest year for which figures are available) 1 : 4.0, which is approaching the latest available (1939) ratio for cancer of the mouth (Table XVII). Cancer of the lung differs from the other forms of cancer in Table XVII in that the proportion of males is increasing. During the whole




FIG. 3.—Deaths from cancer of lung. Males and Females. 1935-1945.

period under consideration here (1921-1945) the ratio of female to male deaths from cancer of the lung has changed from 1 : 1.9 to 1 : 4.0. A similar change has occurred in the United States: the ratio of the standardized death rates for cancer of the lung and pleura in white men and women was 3 : 1 in 1928, and 20 years earlier was less than 1 : 5 : 1 (Statistical Bulletin, 1939).

Fig. 3 shows that the rate of increase in recent years of deaths attributed to cancer of the lung is much greater in men than in women for the curve of deaths

GG 15268

| BB 0007259 |

of males rises much more steeply than does that of deaths of females. But this apparent great difference between the sexes is due in large part to the fact that, in a graph of this type, a given increase in a larger quantity is much more conspicuous than is the same percentage increase in a smaller quantity. If one reckons the annual numbers of deaths as a percentage of those in any given initial year (Table XVI), the difference between the two sexes is not nearly so conspicuous (Fig. 4).




FIG. 4.—Cancer of lung. Males and Females, 1935-1945. Deaths per annum as percentage of deaths in 1935.

TABLE XVI.—Totals of Death Certificates (England and Wales) and Percentage Increase in Cancer of the Lung.

	Cancer of lung.		Cancer of larynx.	
	Men.	Women.	Men.	Women.
1935	2337 = 100	558 = 100	903	224
1936	2606 = 112	917 = 107	916	275
1937	2914 = 125	927 = 108	918	232
1938	3273 = 140	1049 = 122	887	259
1939	3391 = 145	1149 = 134	929	280
1940	3898 = 164	1180 = 137	903	267
1941	4069 = 174	1156 = 133	929	269
1942	4579 = 196	1267 = 148	837	255
1943	5081 = 217	1296 = 151	972	281
1944	5491 = 235	1360 = 158	852	280
1945	5982 = 256	1480 = 172	841	279

CG 15269

BB 0007260

NOTE: THIS DOCUMENT DID
NOT COME FROM THE FILES

280

E. L. AND N. M. KENNAWAY

TABLE XVII.—*Cancer of Lung, Oesophagus and Mouth Region in Men and Women. (England and Wales).*

	Site of cancer.	Number of deaths.		Ratio.
		Women.	Men.	
1921	Lung	186	361	1 : 1.9
1932	"	365	1553	1 : 2.7
"	Oesophagus	661	1774	1 : 2.7
"	Mouth region (lip, tongue, mouth, tonsil, jaw, pharynx)	536	3040	1 : 5.7
1937	Lung	927	2914	1 : 3.1
1939	"	1020	3544	1 : 3.4
"	Oesophagus	729	1590	1 : 2.2
"	Mouth region	585	2538	1 : 4.3
1945	Lung	1480	5982	1 : 4.0

Cancer of larynx.

Cancer of the larynx has increased between 1921 and 1945 from 641 to 841 cases per annum in men, with a maximum of 972 in 1943, and from 138 to 279 cases in women (Table XVI). These figures are in the ratio of 1 : 1.3 in men and 1 : 2.0 in women. But the figures for men have been fluctuating in the region between 850 and 970 since 1930, and in women there has been no uniform increase since 1928. Hence if the adoption of smoking by women has had any effect upon their liability to the disease, this factor has reached equilibrium.

COMPARISON OF THE RATIOS IN THE EARLIER AND LATER PERIODS.

The calculation of these ratios (Tables IV and V) for the two periods 1921-32 (A) and 1933-38 (B) provides material for a comparison. If the data from which the ratios are derived are reliable, this comparison should show whether the incidence of cancer of the lung or larynx upon any given occupation has undergone changes differing significantly from any which have occurred in the general population, and the general degree of agreement between the ratios for the two periods is some indication of the value of conclusions based on death certificates. In Fig. 5 the comparison is made for cancer of the lung by calculating the ratio between the earlier and later ratios ($B/A \times 100$) and setting down the figures in order of magnitude. Occupations with not more than one death in either period are omitted here, and are considered elsewhere (p. 286). The results shown in the graph may be summarized thus:

Ratios below 66	11
" 66 to 133	35
" above 134	3
	49

Thus 35/49 of the ratios of one period, or 71.4 per cent, are within ± 33 per cent of those of the other, which is perhaps as close an agreement as one could look

BB 0007261

GG 15270

University of Kentucky




FIG. 5.—Cancer of lung. Males in 49 occupations.

A. Ratio of registered to 100 calculated deaths, 1921-1932.

B. Ratio of registered to 100 calculated deaths, 1933-1938.

Dots represent $B/A \times 100$.

Occupations with not more than one death in either period are omitted.

for in such material. When the individual points on the graph are examined, the larger occupations are found to lie chiefly in the middle part of the series, as one would expect from the lower sampling error.

A similar treatment of the figures for cancer of the larynx (Fig. 6) gives the following result:

Ratios below 66	7
„ 66 to 133	28
„ above 144	14
	—
	49

In this case 28/49 or 57 per cent of the earlier ratios are within ± 33 per cent of the later ones. There are thus more occupations in which the ratio has increased than in the case of cancer of the lung.

OCCUPATIONAL INCIDENCE OF THE INCREASE OF CANCER OF THE LUNG.

If there is a real increase in cancer of the lung, and if this is due to some external factor, one might find differences in the rate of increase in classes of persons exposed to different environments. The annual numbers of deaths from cancer of the lung, and of the larynx, in each one of the occupations studied in detail in this paper can be seen in Table XVIII. The increase in the total of recorded cases has been so great (Fig. 1) that, if this increase were due to a contribution from any of the smaller occupations, the incidence upon these would have to be very high indeed.

GG 15271

BB 0007262

NOTE: THIS STATEMENT DID

NOT COME FROM THE FILES

TABLE XVIII.—Number of Deaths from Cancer of the Lung and Larynx
in 36 Occupations in each Year 1933-1938.

The occupations are placed in the same order as in Table IV.

		1933.	1934.	1935.	1936.	1937.	1938.	Total.
Labourers, patent fuel works	Lung	—	—	—	—	—	1	1
	Larynx	—	—	—	—	—	—	—
Gas stokers and coke-oven chargers	Lung	9	5	4	8	11	11	48
	Larynx	2	3	3	4	2	3	17
Gas producer men	Lung	1	—	4	—	—	3	8
	Larynx	—	—	—	—	—	—	—
Metal grinders	Lung	1	3	4	1	5	6	20
	Larynx	3	2	—	1	2	1	9
Gas-works foremen and inspectors	Lung	2	2	3	5	—	3	15
	Larynx	—	—	—	1	1	—	2
Potters' mill workers; slip makers and arkmens	Lung	—	1	1	1	1	—	4
	Larynx	—	—	—	—	—	—	—
Mainly council labourers, road sweepers and dustmen	Lung	18	10	16	18	16	25	103
	Larynx	6	7	9	16	12	13	63
Gas fitters	Lung	—	4	4	8	3	6	25
	Larynx	1	—	1	2	—	2	6
Paviours, street masons, concretors and asphalters	Lung	4	3	2	1	2	2	14
	Larynx	—	1	1	2	—	—	4
Motor drivers, goods and passengers	Lung	20	29	42	43	72	85	301
	Larynx	7	6	8	8	8	11	48
Blast furnace engine men	Lung	—	—	1	—	—	—	1
	Larynx	—	—	—	—	—	—	—
Cabinet makers	Lung	19	—	9	14	14	13	60
	Larynx	5	2	5	6	—	2	20
Tobacco manufacturers	Lung	3	2	—	1	—	2	8
	Larynx	—	—	3	1	—	—	4
Stone masons, cutters and dressers	Lung	11	15	13	16	17	13	85
	Larynx	1	1	4	3	2	1	12
Tanners, leather dressers, curriers, skilled workers	Lung	4	5	3	5	4	9	30
	Larynx	—	1	2	—	2	—	5
Potters; ware-makers, casters and finishers	Lung	1	2	1	1	5	3	13
	Larynx	—	1	—	—	1	2	4
Gas-works engine and crane drivers	Lung	—	1	2	1	—	1	5
	Larynx	—	1	—	—	1	—	3
Sandblasters	Lung	—	—	1	—	—	—	1
	Larynx	—	—	—	—	—	—	—
Gas-works managers	Lung	—	—	—	—	—	1	1
	Larynx	—	1	—	—	—	—	1
Drivers of horse-drawn vehicles	Lung	25	49	33	51	45	44	240
	Larynx	15	17	23	17	18	7	100
Gas-works labourers	Lung	4	8	8	9	9	12	50
	Larynx	1	1	3	5	1	—	11
French polishers	Lung	5	2	4	4	2	4	21
	Larynx	2	2	4	—	—	1	9
Painters	Lung	29	47	43	51	76	77	323
	Larynx	27	24	20	11	18	20	120
Licensed victuallers	Lung	18	33	21	29	38	33	172
	Larynx	17	21	14	22	14	14	102
Printers	Lung	14	13	23	14	26	26	116
	Larynx	6	4	6	6	7	4	33
Chimney sweeps	Lung	—	—	3	1	2	4	10
	Larynx	1	—	1	1	2	1	6
Plumbers	Lung	5	10	8	12	14	17	66
	Larynx	3	1	2	5	4	4	22
Burners	Lung	2	1	6	1	2	5	20
	Larynx	3	3	2	2	1	2	13

TABLE XVIII (contd.).

		1932.	1933.	1934.	1935.	1936.	1937.	1938.	Total.
Tobaccoists and their assistants	Lung	1	4	1	3	3	3	3	17
	Larynx	3	3	3	—	3	2	2	14
Hairdressers	Lung	5	8	4	11	6	13	13	47
	Larynx	3	2	3	—	—	2	2	10
Bakers and pastry cooks	Lung	10	9	12	15	12	16	16	76
	Larynx	3	7	2	4	5	3	3	24
Professional men	Lung	25	29	27	34	26	35	35	176
	Larynx	6	3	4	9	10	5	5	39
Cellar-men	Lung	2	—	—	1	1	1	1	6
	Larynx	2	2	3	1	2	2	2	12
Pottery, etc. kiln and ovenmen, kiln setters and placers	Lung	1	1	2	2	3	3	3	12
	Larynx	—	5	—	2	1	2	2	10
Grooms and horsekeepers	Lung	3	8	5	5	2	7	7	30
	Larynx	4	2	4	—	5	2	2	17
Brick kiln and ovenmen	Lung	3	—	2	1	—	3	3	9
	Larynx	—	2	—	—	1	—	—	3
Stationary engine and crane drivers	Lung	9	9	15	11	14	17	17	75
	Larynx	7	4	6	2	2	3	3	24
Cotton strippers, grinders and card room jobbers	Lung	—	—	—	2	1	3	3	6
	Larynx	—	—	—	—	1	1	1	2
Locomotive engine drivers, firemen and cleaners	Lung	11	6	10	13	13	13	13	66
	Larynx	3	5	1	2	5	2	2	18
Carpenters	Lung	25	34	35	31	35	40	40	200
	Larynx	21	14	17	18	15	12	12	97
Stone miners and quarrymen	Lung	2	2	1	5	3	4	4	17
	Larynx	6	2	2	1	1	—	—	12
Coal miners—lowers and getters	Lung	51	49	51	40	62	66	66	319
	Larynx	20	14	24	14	20	15	15	107
Lithographic and process engravers	Lung	2	1	—	—	—	1	1	4
	Larynx	—	2	—	1	2	—	—	5
Blast furnacemen and labourers	Lung	—	—	1	1	1	1	1	4
	Larynx	1	—	—	1	1	1	1	4
Gardeners	Lung	21	31	38	40	41	42	42	213
	Larynx	16	23	11	25	20	28	28	123
Cotton spinners and piecers (mule, ring, cap or flyer)	Lung	4	4	3	6	7	3	3	27
	Larynx	1	2	2	3	—	3	3	11
Coal miners—workers above ground	Lung	6	11	5	11	7	9	9	49
	Larynx	7	2	3	5	3	6	6	26
Tar distillery workers and coke-oven workers	Lung	1	2	2	—	—	1	1	6
	Larynx	—	—	—	—	—	—	—	—
Brick and plain tile makers	Lung	—	—	2	1	2	—	—	5
	Larynx	—	—	—	—	—	—	—	—
Coal miners—other workers below ground	Lung	6	7	10	10	7	12	12	52
	Larynx	2	1	4	5	2	4	4	18
Coal miners—conveying material to the shaft	Lung	3	2	5	3	4	11	11	28
	Larynx	2	—	—	—	1	4	4	7
Coal miners—making roads	Lung	6	11	2	7	4	7	7	37
	Larynx	2	4	4	6	3	3	3	24
Cotton weavers	Lung	3	5	3	3	3	—	—	17
	Larynx	3	—	1	2	2	4	4	12
Farmers	Lung	17	24	21	29	27	30	30	148
	Larynx	20	12	8	14	16	11	11	81
Agricultural labourers, including shepherds	Lung	17	21	17	31	23	28	28	137
	Larynx	21	19	28	25	22	22	22	135
Farm builders and foremen	Lung	2	—	1	1	1	1	1	6
	Larynx	1	—	2	—	—	—	—	3
	Lung								3550
	Larynx								1451

NOT TO BE REPRODUCED
OR COME FROM THE FILES

GG 15273

BB 0007264




FIG. 6.—Cancer of larynx. Males in 49 occupations.

A. Ratio of registered to 100 calculated deaths, 1921-1932.

B. Ratio of registered to 100 calculated deaths, 1933-1938.

Dots represent $B/A \times 100$.

Occupations with not more than one death in either period are omitted.

TABLE XIX.—Increase of Deaths Attributed to Cancer of Lung in Certain Occupations (1927-1938).

		Deaths from cancer of lung.		
		1927-32.	1933-38.	Difference.
See Table VI.	Paviours, street masons, concrete- tors, asphalters	7	14	7
	Council labourers, road sweepers, dustmen	47	103	56
	Drivers of horse-drawn vehicles	111	240	129
	Motor drivers	102	301	199
		267	658	391
General population, males		6296	15,045	8749

For example, if one takes the numbers of deaths from cancer of the lung in those occupations where there is especial exposure to road dust (Table VI) and divides them between two successive periods of 6 years (1927-32 and 1933-38) and compares these figures with those given by the whole male population, it becomes quite clear that even these occupations account for only a small fraction (391 cases or 4.5 per cent) of the whole increase of 8749 cases (Table XIX).

The following example shows the effect of the increase in deaths attributed to cancer of the lung upon the data for a single occupation.

GG 15274

BB 0007265

210,431 carpenters produced 123 deaths from cancer of the lung in 12 years (1921-33) or 10.2 deaths per annum, and in the next 6 years (1933-38) 243,928 carpenters produced 200 such deaths, or 33.3 deaths per annum (Table IV). The number of carpenters producing 1 death in 1 year was therefore $210,431 \div 10.2 = 20,640$ in the first period and $243,928 \div 33.3 = 7324$ in the second. The ratio between the two figures, namely 3.3 : 1, is reasonably near the inverse of the ratio (1 : 2.8) between the mean annual deaths (755 and 2507) in these two periods from cancer of the lung in the whole male population. The result indicates that the increase of cancer of the lung has fallen upon carpenters, who undergo considerable exposure to some kinds of dust, in much the same measure as it has upon men in general (Fig. 7).




FIG. 7.—Deaths from cancer of lung. Carpenters, 1921-1938.

Occupational exposure to three agents which are perhaps liable to induce cancer of the lung (asbestos, arsenic, nickel [Factories Report, 1938, p. 72]) occurs in employments which comprise very few workers, and hence have no appreciable effect upon the total. Of these three substances, arsenic seems to be the one to which the general population is most likely to be exposed. No attempt is made here to deal with all the data on the carcinogenic action of arsenic, as this subject is under investigation by a Committee of the Industrial Health Board, but the following quotations from recent Annual Reports of the Senior Medical Inspector of Factories show the possible importance of arsenic in relation to cancer of the lung. "In the case of arsenical poisoning due to sodium arsenite, which was fatal, the post-mortem examination revealed that in addition to pigmentation of the trunk and limbs, warty growths all over the body, and perforation of the nasal septum, there was a primary cancer of the right lung with metastatic growths in neighbouring glands and in the liver" (Factory Report, 1939, p. 22). "The other (case) occurred in a filling machine operator aged 57, for 43 years in a factory manufacturing sheep dip containing sodium arsenite, the cause of death being due to carcinoma of the right lung."

"Three similar cases of pulmonary carcinoma occurring in arsenical sheep-dip workers have been notified since 1939" (Factory Report, 1943, p. 45).

Neubauer (1947) collected the literature relating to 142 cases of medicinal arsenical epithelioma; in one of these cancer of the lung occurred (Montgomery, 1935; Montgomery and Waisman, 1941). The patient had taken

a medicine for eczema for six years in childhood, which was assumed to contain arsenic on account of the appearance from the age of 42 onwards of numerous epitheliomas of the skin of the arsenical type. Later an intraurethral epithelioma developed, and finally an epithelioma of the bronchus with "malignant dyskeratosis characteristic of the arsenical type of epithelioma," and extensive metastases.

Semon (1945) records a case of bronchial carcinoma in a man who, during a period of about 20 years, had taken large doses—up to 1 drachm 3 times daily—of Fowler's solution (equivalent to 1 per cent arsenic trioxide) for recurrent attacks of severe dermatitis herpetiformis. His skin was unaffected. There is, of course, nothing to show that the bronchial carcinoma was due to the arsenic, but unless attention is drawn to such a case others may not be noticed.

TABLE XX.—*Cancer of the Lung. Occupations Showing a Considerable Rise in the Ratio in the Later Period.*

	1921-32. A.			1933-38. B.			Ratio B A x 100.
	Population.	Number of deaths.	Ratio.	Population.	Number of deaths.	Ratio.	
Potter's mill workers, slip-makers and arkmen	1,436	1	96	1,766	4	218	227
Gasworks engine and crane drivers	1,854	1	60	1,800	5	186	310
Cotton spinners and piecers	38,817	7	32	43,991	27	75	234
Cotton strippers and grinders and card room jobbers	5,242	1	39	5,242	6	104	347
Tar distillery workers and coke-oven workers	9,697	2	29	7,824	6	70	242
Brick kiln and ovenmen	6,225	1	23	7,112	9	121	528
Ministers of other religious bodies	11,769	1	88	11,769	15	157	

NOTE: THIS DOCUMENT DID NOT COPY FROM FILES

(1) *Cancer of the lung.*

In 7 occupations the ratios for B are higher than those for A by more than 50 per cent (Table XX and Fig. 5). In all of these the ratios in A are lower, and in most cases much lower, than 100, and hence the considerable rise raises the ratio in B above 150 in two cases only (potter's mill workers, slip makers and arkmen 218; gas-works engine and crane drivers 186). In 4 of the 7 occupations there is only one death in A, and hence the comparison of the two periods is liable to considerable sampling error. The fact that the rise starts from a low level does not, of course, prove that it is unimportant, and these occupations must be watched in the subsequent years to see if the rise continues.

Cases of cancer of the lung have occurred among stokers of oil-fired boilers, but no figure for the population of these workers is available.

(2) *Cancer of the larynx.*

In 9 occupations the ratios for B are higher than those for A by more than 50 per cent (Table XXI and Fig. 6). In 6 of these there are not more than 3 deaths in the first period, so that the sampling error is high, and the ratio in the first period is low.

GG 15276

BB 0007267

TABLE XXI.—*Cause of the Excess of Deaths Showing a Considerable Rise in the Ratio in the Later Period.*

	1921-22, A.			1943-45, B.			Ratio B/A x 100.
	Population.	Number of deaths.	Ratio.	Population.	Number of deaths.	Ratio.	
Celermen	6,533	14	317	7,050	12	501	158
Metal grinders	15,220	10	141	17,009	9	234	166
Brick-kiln and oven men	6,225	3	71	7,112	3	123	173
Pottery, steel kiln and oven men	12,188	9	116	14,442	10	220.5	190
Cotton strippers and grinders and card room jobbers	5,242	2	60	5,342	2	118	197
Tobacco manufacturers	7,695	3	65	8,192	4	129	198.5
Gas works managers	1,393	1	78	1,049	1	187	207
Lithographic and process engravers	5,704	3	84	7,289	5	214	255
Blast furnacemen and labourers	9,391	3	40	6,135	4	138	340

ERRORS IN THE DIAGNOSIS OF CANCER OF THE LUNG.

Some hold that cancer of the lung is not increasing, and that the ascent of the curve (Fig. 1) is due to improved diagnosis.

This statement is easy to make and difficult, perhaps impossible, to disprove, and is apt to serve as a substitute for any investigation of the question. If one accepts it, at least one conclusion follows which is of some interest. Let us suppose that the steeply rising line (Fig. 1) represents the detection of an increasing fraction of a pool of cases awaiting discovery. When the means of diagnosis, and their availability, can be improved no further, the curve must flatten. We cannot say when this will happen, so for the present we must take the latest, and highest, figure (5982 deaths of males in 1945) as representing the total. If the actual number has altered little during recent years a similar number of deaths took place in, say, 1921; but only 361 of these deaths (Fig. 1) were assigned to the right cause. Hence in that year in at least $5982 - 361 = 5621$ fatal cases of cancer of the lung the cause of death entered upon the death certificate was wrong. This is of course possible, but it would be interesting to know what were the 5621 erroneous descriptions of fatal disease.

Some conditions which might give rise to false diagnoses in cases of cancer of the lung are* :

(1) *Mediastinal tumour*.—The increase of cases of cancer of the lung cannot depend wholly, or even largely, upon a diversion to that category of cases formerly set down as mediastinal tumour because the number of these has never been sufficient.

Thus the death certificates for males show no fall in deaths attributed to cancer of the mediastinum during the years following 1931 (when this form of cancer is first mentioned in the Registrar-General's Statistical Reviews), which would account for the increase of 7.24 in cases of cancer of the lung which took place in the years 1931-37 (Table XXI).

* I am indebted to Professor D. W. Slaughter, Professor R. A. Willis and Dr. Eric Scowen for information on this subject.

NOTE: THIS DOCUMENT DID NOT COME FROM THE FILES

GG 15277

BB 0007268

NOTE: THIS DOCUMENT DID
NOT COME FROM PMA FILES

288

E. L. AND N. M. KENNAWAY

TABLE XXII.—*Cancer of Lung and of Mediastinum.*

	Deaths of males. England and Wales.	
	Cancer of lung.	Cancer of mediastinum.*
1931	1,055	135
1932	1,553	262
1933	1,820	262
1934	2,095	244
1935	2,345	248
1936	2,611	222
1937	2,930	308
Total	14,409	1681
7 × 1055	7,385	
Increase	7,024	

* These figures are taken from the Registrar-General's Statistical Reviews for the years in question.

(2) *Pulmonary infections.*—A long list could be compiled of infective processes and their results which might have to be considered in the diagnosis of cancer of the lung in the living subject, but those most likely to be named in error on death certificates are tuberculosis, lung abscess, pneumonia, and bronchiectasis. In theory, the numbers of deaths attributed to these conditions should be falling if increasing numbers of cancers of the lung are being detected among such cases. But actually the numbers of deaths attributed to respiratory tuberculosis, and to pneumonia, are so large, ranging from 2 to 10 times those assigned to cancer of the lung, that the proportional fluctuations prevent any diminution which could be due to this cause from being identified (Fig. 8). In this figure the two curves representing pulmonary infections show no uniform relationship to the third representing cancer of the lung. The numbers of deaths from abscess of the lung (Table XXIII) are too small (mean 100 per annum) to be of importance, and moreover they show no continuous fall during the period. Bronchiectasis is not separated from bronchitis in the Registrar-General's Statistical Reviews.

TABLE XXIII.—*Deaths of Males from Abscess of Lung.*
England and Wales.

1932	92	1937	96
1933	82	1938	132
1934	91	1939	119
1935	111	1940	84
1936	102	1941	92
	1942	100	

GG 15278

BB 0007269 1

University of Manchester

NOTE: THIS DOCUMENT DID
NOT COME FROM THE FILES

there was a special surgical clinic dealing with lung cases, and where full X-ray and bronchoscopic investigation was practised, 42 per cent of the clinical diagnoses were wrong (Table XXIV). Hence the diagnosis of cancer of the lung appears to be still very incomplete, even under favourable conditions, in spite of the great improvements which are generally stated to have been made. At any rate one cannot well emphasize at the same time both the high proportion of cases (e.g. 42 per cent) which escape detection through inaccurate diagnosis, and the adequacy of the improvements in diagnosis to explain the apparent increase in cancer of the lung, though the two statements are of course compatible. Willis does not give any information about false positive diagnoses of pulmonary cancer.

Professor Willis (private communication) has recorded a very instructive example of the statistical effect of diagnostic errors. In a series of 1000 consecutive autopsies which showed the presence of cancer, 155 were found to be cases of cancer of the stomach, and of these, 109, or 70 per cent, had been diagnosed correctly and 46 had therefore received incorrect negative diagnoses. There were also 40 cases diagnosed as cancer of the stomach in which this condition was not present. In this group of 1000 cases, therefore, $109 \div 40 = 149$ cases had been diagnosed as cancer of the stomach, and the actual number present was 155. In so far as the total is concerned therefore the clinical answer to the statistical question—how many cases of cancer of the stomach were there in these 1000 cases?—was very nearly correct. The far greater error in the diagnoses of individuals, of whom 86 were described wrongly, is concealed statistically by the approximate cancellation of the 46 false negatives by the 40 false positives.

This process of attaining accuracy by the balance of positive and negative errors is, of course, neither desirable nor reliable, but it is unavoidable in such material, and is bound to happen to a greater or less extent whether we like it or not. When any decisions are reached on inadequate evidence in any subject, there will be the same tendency for errors in opposite directions to cancel each other.

FACTORS WHICH MIGHT CAUSE AN INCREASE IN THE DEATHS ATTRIBUTED TO CANCER OF THE LUNG.

The increase in recent years in the number of deaths attributed to cancer of the lung might be due to one, or more than one, of the following factors:

(1) An increase in the actual number of cases of the disease.

It is obviously difficult to obtain any direct evidence of this factor; one is driven to seek for indirect evidence by means of enquiry how far the second and third factors can be excluded. Some possible causes of a real increase are considered in the next section.

In the earlier paper (Kennaway and Kennaway, 1936) we drew attention to the evidence afforded by cases of cancer of the lung in which an erroneous diagnosis was made based on nervous or mental symptoms due to metastases. Thus Fried and Buckley (1930) record that metastases in the central nervous system were found in 15 of 38 proved cases of cancer of the lung. In 11 of the 15 a diagnosis of primary tumour of the brain was made and the lung tumour overlooked; an intracranial operation was performed in 11 and the metastases

University of Manchester

GG 15280

BB 0007271

removed in 10 cases. Obviously neither improvement nor fashion in the diagnosis of cancer of the lung (see below) come into play here, and hence if it can be shown that such cases are, as some physicians believe, increasing, a real increase in cancer of the lung is indicated.

(2) Improvement in diagnosis, whereby a larger proportion of the actual number of cases is detected.

The methods of examination of the lungs, and the availability of apparatus for such purposes have undergone very great development in the last twenty years, and it would be remarkable if these changes had not increased the number of cases of cancer of the lung discovered. This matter is discussed under "Errors in diagnosis" above. According to some authorities (e.g. Willis) these improved methods, even when fully available, still leave about half the cases undetected, so we may look for a further considerable increase from this source.

(3) A tendency to identify as cancer of the lung cases which are really of a different nature; that is, a fashion in diagnosis.

We have not found any data which would enable one to judge whether this factor is of any importance; if it is to account for the form of the curve shown in Fig. 1, two or three hundred *more* of such errors must occur every year. Willis (p. 289) does not mention such positive errors in the diagnosis of cancer of the lung.

SOME POSSIBLE CAUSES OF CANCER OF THE LUNG IN MAN.

Experimental pathology has not yielded any conclusive evidence on this question, as it has done in the case of the occupational cancers of the skin. Most of the laboratory work, such as that of Argyll Campbell, has been done on the mouse, in which species the neoplastic factors in the lung are certainly different from those in man.

(1) In default of experimental evidence, indications of at any rate one direction in which to seek are derived from the material of the General Register Office. These data are compatible with the importance of some factor in town air, e.g. coal smoke.

(a) Cancer of the lung differs from that of the upper alimentary tract, larynx and skin in showing no inverse relationship to ascent in the social scale (Stevenson, 1923). Owing to the mixing action of the wind* there is less difference in the outdoor air breathed by different classes than in other social conditions such as food and cleanliness.

(b) The higher incidence in towns (Stocks, 1936). Table XXV shows that there is at every age a steep downward gradient of rates from London through

* An elaborate study of the mixing of air-borne particles was made at Leicester. The effect of wind appears to be not so great as one would expect. "The maximum effect of wind is to displace the point of maximum concentration by 1 mile" (Leicester, "Atmospheric Pollution," 1945, p. 123). But it is difficult to believe that, say, an east or west wind blowing across a city does not have a considerable effect in equalizing the concentration of the air in the east and west ends. Great importance is attached also to turbulence as a mixing agent. "No exact definition of turbulence has yet been made, but the word has been used, in this report and elsewhere, to describe the readiness with which air mixes with itself. During the last nine months of the Leicester Survey, special attention was given to the measurement of variables connected with turbulence, because of the importance of turbulence in determining the concentration of suspended pollution. It was thought that one of these variables might prove suitable for more general use in connection with measurements of atmospheric pollution" (p. 114). Apparently turbulence includes all forms of disturbance due to unequal local heating.

NOTE: THIS DOCUMENT IS NOT TO BE REPRODUCED OR TRANSMITTED IN ANY FORM OR BY ANY MEANS

CG 15281

BB 0007272

large and small towns to rural districts." There are, however, exceptional large towns with a low rate (e.g. Bristol). Empirically, smoke concentration in a town is in proportion to the square root of the population (Leicester. 'Atmospheric Pollution,' 1945, p. 124).

(c) The low incidence in agricultural workers (Kennaway and Kennaway 1936). There is an equally low incidence in coal miners.

(b) and (c) might be explained by the better conditions for diagnosis in urban areas.

Coal smoke does not account well for an increase in cancer of the lung during a period when, as one would think, atmospheric pollution has not increased owing to greater use of gas and electricity. The domestic fire is the chief source of soot in the atmosphere,* and it is in the domestic field that the substitution of gas and electricity for coal has been most general.

One must, however, note that ten years ago the authorities on atmospheric pollution were not unanimous on the question whether there had been any measurable diminution.†

NOTE: THIS DOCUMENT IS NOT TO BE USED FOR FILES

TABLE XXV.—Cancer of Lung. Males. England and Wales. 1921-1930. (Sticks.)

	Rates per million at ages shown.								
	25-	35-	45-	50-	55-	60-	65-	70-	75-84.
London	15	55	121	193	253	263	236	199	215
County Boroughs	7	35	73	103	135	168	148	144	93
Other Urban Districts	5	23	45	77	96	128	150	118	93
Rural Districts	7	18	30	34	63	73	87	92	71

(2) One must look, therefore, for some other factor to explain the increase, if there is a real increase, in cancer of the lung in recent years. Many such factors have been suggested, e.g. influenza, tobacco, popular drugs, tarred roads, exhaust gases and other emanations, including lead ethyl, from motor vehicles driven by petrol or Diesel engines.

No attempt is made here to discuss all these possible factors.

* " . . . the percentage of tar in domestic smoke is very high, reaching 30-40 per cent. while in industrial smoke from boiler furnaces and such there is practically no tar " ('Atmospheric Pollution,' 12th Report, p. 46). " A high deposit of tar can only be due to pollution from domestic chimneys or very inefficient industrial furnaces, since in a properly constructed and worked furnace a high temperature is maintained for a sufficient distance from the main place of combustion for all combustible vapours to be fully burned before the furnace gases leave the chimney " ('Atmospheric Pollution,' 23rd Report, p. 9). Soot from the chimney of a dwelling house may contain 40 per cent of tar (Cohen and Ruston, 1925), and the dust suspended in the air of a town ('Leicester Atmospheric Pollution,' 1945) may contain 14 per cent of tar.

† In the '22nd Report on the Investigation of Atmospheric Pollution' (1935-1936, p. 5) " . . . Chairman sums up some rather equivocal evidence thus: " . . . on the whole there has been a definite reduction in the extent of pollution of the atmosphere in Britain during the past twenty years." On p. 11 of the same report, however, the conclusion is drawn that " no general statement can be derived from the increases and decreases in different areas during 1925-1935. The special study of atmospheric pollution in Leicester ('Leicester Atmospheric Pollution,' 1945) showed no difference between the two periods 1927-1932 and 1933-1939 in the amount of tar in deposited matter: the report makes the suggestion that a decrease in tar might be masked by an increase in material from motor cars. But the inconclusive nature of the data shows that there has been no increase in atmospheric pollution in any way proportional to the increase in cancer of the lung.

GG 15282

1 BB 0007273 1

The subject of tarred roads was dealt with in the earlier paper (Kennaway and Kennaway, 1936).

Although the incidence of lung cancer in those occupations where there is exposure to road dust (see above, Table VI) is rather above the general level, it seems very questionable whether the recent increase of this disease among the general population can be attributed to the tarring of roads. For scores of years before this increase began coal tar was being discharged into the atmosphere in the form of soot by the domestic fire in quantities vastly greater than any that could now be derived from roads. Tarring of the roads has undoubtedly increased the amount of dust derived from tar in the air, but this increase appears negligible in comparison with the amount of soot present already. One sees clouds of smoke drifting over any large town, but one does not see any similar clouds of tar-laden vapour arising continually from tarred roads. The annual consumption of coal for all purposes in Great Britain in 1936-1938 was about 188 million tons, of which 40 to 45 million tons was burned by domestic users, who are by far the greatest producers of smoke and soot.* It is difficult to believe that tarred roads made any important addition to this total.

One must, however, bear in mind that the total amount of soot per unit volume of air may not be the only important factor. The size and physical state of tar particles derived from the domestic fire, and from a tarred road, are no doubt different, and this difference may determine the amount of penetration into the deeper air passages (see, for instance, Boyland, Gaddum and MacDonald, 1947; Owens, 1953).

(b) *Tobacco*.—A possible connection between tobacco, and especially cigarettes, and cancer of the lung, has been suggested many times, perhaps most recently in the case of Turkey (*Lancet*, 1946, i, p. 435).

Peacock (1943) has pointed out that cancer of the stomach is far more common in man than, so far as we know, in any other species, and has suggested that this is due to his use of heated foods. A similar argument might be applied in the case of cancer of the lung, which is not known to be prevalent in any of the lower animals. The adenoma of the lung of the mouse, and certain affections of the lung of sheep in South Africa and Iceland, are neoplasms of which the exact nature is uncertain. We know one instance at any rate of the susceptibility of the lung of an animal to a carcinogenic agent, namely the lung of the cat in relation to 2-acetylaminofluorene given by the mouth (Harding, 1946): hence there is no reason to think that animals are immune to any such agents. One obvious factor, possibly carcinogenic, to which the lung of man alone is exposed is tobacco smoke.

During the present century very considerable changes have taken place in this country in the social distribution of the various methods of smoking. There were, of course, exceptions to any rule, but roughly one might say that in the earlier part of this period men of the richer classes smoked pipes, cigarettes and cigars, and men of the poorer classes smoked pipes; cigarette smoking was increasing among the richer women, while women of the poorer classes did not smoke. One never saw a woman scrubbing her front door-step, or hanging out the washing, with a cigarette in her mouth. The great change which has taken

* A. Parker. Chadwick Public Lecture. March 13th, 1945.

† In 1929 in England and Wales, carcinoma of the stomach was assigned as the cause of 7050 out of 27,629 deaths from carcinoma of all sites in males ('Registrar-General's Statistical Review').

NOTED FOR FILES
NOT COMPLETED

GG 15283

BB 0007274

place in the later years is the general increase of cigarette smoking,* and adoption by women of classes which formerly did not smoke at all. Men




FIG. 9.—Deaths. England and Wales. 1932-1942.

A. Males } Diseases of coronary arteries, angina pectoris.
B. Females }
C. Males } Cancer of lung.
D. Females }

(Data from the Registrar-General's 'Statistical Review of England and Wales for the Year 1942. Tables. Part I. Medical'.)




FIG. 10.—Consumption in the United Kingdom of tobacco (A) and cigarettes (B).

* The Secretary of State for War stated in the House of Commons on January 23, 1945 (Parliamentary Debates (Hansard), 407, 626) that "In making up the stocks of tobacco and cigarettes for the Army overseas it is assumed that there are ten cigarette smokers to every one pipe smoker. This proportion is based on experience."

GG 15284

BB 0007275

things which do not suggest them in any restriction, e.g. factory workmen, painters, window cleaners, dustmen and road sweepers, who used to smoke a pipe at meal times, now smoke cigarettes while at work. Cigarette smokers are said to inhale more than do smokers of pipes, but it is very difficult to get any conclusive evidence upon this matter, which might be important. There is some American literature, which has been summarized (*Brit. med. J.*, 1946, i, 94) upon the arsenic content of tobacco and tobacco smoke. The occurrence of cancer of the lung in makers of arsenical sheep-dip (p. 285) indicates the possible importance of this factor.

Some writers have contended that smoking cannot be associated with cancer of the lung, because this form of cancer has increased more among men, while the use of tobacco has increased more among women. But the sexual distribution of cancer involves unknown factors (Table XVII), and does not provide a




FIG. 11.—Cigarettes as percentage of tobacco consumed in the United Kingdom, 1924–1943.

very secure basis for an argument of this kind. Of course no claim is made here that the simultaneous increases in the consumption of tobacco, and in cancer of the lung, proves any etiological connection between the two. Other changes which have taken place in the same period, which no one proposes to associate with cancer of the lung, e.g. the increase in the issue of wireless licences, show a very similar curve, and such correlations are a common subject for statistical witticisms. Thus wireless licences have increased at a rate (about 10-fold in the last 20 years) similar to that shown by deaths from cancer of the lung, or from coronary disease (Fig. 9).

The annual consumption of tobacco in the United Kingdom has increased from 128 million pounds in 1924 to 250 million pounds in 1946, and the percentage of these amounts smoked in the form of cigarettes has risen from 56 in 1924 to over 80 in 1943–5* (Fig. 10 and 11). Thus the consumption of cigarettes shows a considerable increase, both absolute and relative.

* We are indebted to Mr. J. R. Willis, M.C., of the Board of Trade for these data. The proportion of cigarettes to total tobacco consumption is available only in those years for which a Census of Production is taken. The monthly net clearances of tobacco mainly by manufacturers for home consumption in the United Kingdom of Great Britain and Northern Ireland are given in the *Monthly Digest of Statistics*.

CG 15285

BB 0007276

NOTE: THIS DOCUMENT DID
NOT COME FROM THE FILES

206

E. L. AND N. M. KENNEDY

To summarize the evidence brought together in this section on the incidence of cancer of the lung: the higher mortality in towns, the low mortality in agricultural occupations, and the absence of social gradient, are compatible with a factor in the air, such as coal smoke. In any such comparison of urban and rural areas, the question of facilities for diagnosis must be considered. By coal smoke, which is probably a decreasing contaminant of the air, does not account well for the increase of cancer of the lung. Among various possible factors is tobacco smoke; the consumption of tobacco has risen, and so has the percentage of it smoked in the form of cigarettes, of which the smoke is often inhaled. An effect of tobacco would accord well with the absence of social gradient.

SUMMARY.

(1) The death certificates for cancer of the lung and of the larynx in males from England and Wales for the years 1921-38 inclusive, numbering 38,418 have been investigated and the periods 1921-32 and 1933-38 are compared. The 63 occupations examined employ about 30 per cent of the male population aged 20 and upwards.

(2) Sources of error in statistical work on death certificates are discussed.

(3) The increase in the recorded cases of lung cancer cannot be attributed to any increase of data obtained by autopsy.

(4) The agricultural and coal-mining industries show a low incidence of cancer of the lung and of the larynx.

(5) A group of open-air occupations, where there is exposure to the dust of roads, has ratios above 100 for cancer of the lung and of the larynx, with the exception that motor drivers have a normal liability to cancer of the larynx. But the comparative incidence of cancer of the lung is not increasing distinctly in any of these occupations, and in the pavements, street masons, concretors and asphalters there has been a distinct fall in the ratio.

(6) The occupations in which there is a liability to silicosis do not show high incidence of cancer of the lung, but there are in the literature some studies of small numbers of cases in which the two conditions were associated.

(7) Cases of cancer of the lung have occurred in some occupations involving exposure to asbestos.

(8) In the death certificates examined, and in the Reports of the Chief Inspector of Factories, no occupations involving exposure to any kind of dust except those concerned with asbestos, arsenic and nickel, which employ very small numbers, have been found in which there might be an increased incidence of cancer of the lung.

(9) Workers exposed to coal-gas and tar tend to show an increased prevalence of cancer of the lung, but in the later period studied the incidence does not exceed two-and-a-half times that on the general population.

(10) Occupations concerned with the supply of alcohol have a high incidence of cancer of the larynx.

(11) The later period studied shows a considerable decrease in the occurrence of cancer of the lung in those engaged in the preparation and sale of tobacco.

(12) The very moderate ratio (1.25) for cancer of the lung in medical men is important in regard to the view that the recent rapid increase in recorded deaths from cancer of the lung is due to the detection of more cases by improved diagnosis.

University of Liverpool

GG 15286

BB 0007277

areas for this is an occupation where the availability of the existing methods for the detection of cancer is presumably at a maximum.

(13) No special occupations have been found, among the 63 examined, to which the increase in the total of cases of cancer of the lung can be attributed. This increase is now so great that the incidence upon any such occupations would have to be very high, indeed.

(14) No evidence has been found that tarring of roads has affected the incidence of cancer of the lung. Such data as are available suggest that coal-tar in the atmosphere, whether derived from roads, domestic chimneys or any other source, does not cause an exceptionally high incidence of cancer of the lung. Cotton mule spinners show an especially small liability to cancer of the lung, although they inhale air sprayed with an oil which produces cancer of the skin. Much further work is required on the factors which regulate the penetration of particles and droplets of various shapes and sizes into the air passages.

(15) The higher mortality from cancer of the lung in towns (Stocks), the low mortality in agricultural occupations, and the absence of social gradient (Stevenson) are compatible with an etiological factor in the air such as coal smoke. But in any comparison of urban and rural areas, the question of facilities for diagnosis must be considered.

(16) Soot is probably a decreasing contaminant of the air owing to the substitution of other sources of heat for the domestic fire, which is the chief source of soot-containing smoke. Hence coal smoke does not account well for any recent increase in cancer of the lung. Among various possible factors which have been suggested to account for the increase is tobacco smoke; the consumption of tobacco has risen, and so has the percentage of it smoked in the form of cigarettes, of which the smoke is often inhaled; such an effect of tobacco would accord well with the absence of social gradient.

We are indebted to the Registrar-General for the data considered in this paper, and also for kindly permitting us to use some figures which have not yet been made public officially. We have received much assistance in the classification of occupations, and on various other questions, from some members of the staff of the General Register Office, to whom we wish to express our gratitude. We wish to thank Dr. E. R. A. Merewether, Senior Medical Inspector of Factories, for information upon various matters, and especially upon asbestosis, and also Professor R. A. Willis for unpublished material.

We are indebted to the Anna Fuller Fund for secretarial assistance, to the Sir Halley Stewart Trust for a grant, and to Miss Blanche Goode, who has taken part in the classification of death certificates.

REFERENCES.

NOTE.—No attempt is made in this paper to review the statistical literature relating to cancer of the lung in this or other countries.

- ANDERSON, C. S., AND DIBLE, J. H.—(1938) *J. Hyg. Camb.*, 38, 185.
 'Atmospheric Pollution. The Investigation of.' Reports 1-23. London: H.M. Stationery Office.
 ROYLAND, E., GADDUM, J. H., AND MACDONALD, F. F.—(1947) *J. Hyg. Camb.*, 45, 299.
 COHEN, J. B., AND RUSTON, A. G.—(1925) 'Smoke. A Study of Town Air.' London.
 DIBLE, J. H.—(1934) *Lancet*, ii, 982.

NOTE: THIS DOCUMENT DID NOT COME FROM THE FILES

Pamphlet

GG 15287

BB 0007278

- EGBERT, D. S., AND GEIGER, A. J.—(1936) *Amer. Rev. Tuberc.*, 34, 143.
 'Factories, Annual Report of the Chief Inspector of.' London: H.M. Stationery Office.
- FRIED, B. M., AND BUCKLEY, R. C.—(1930) *Arch. Path.*, 9, 483.
- GLOYNE, S. R.—(1933) *Tubercle*, 17, 5.
- HARDING, H. E.—(1946) 23rd Ann. Rep. Brit. Emp. Cancer Campaign, p. 92.
- HENRY, S. A.—(1946) 'Cancer of the Scrotum in Relation to Occupation.' Oxford Univ. Press.
- Idem, KENNAWAY, N. M., AND KENNAWAY, E. L.—(1931) *J. Hyg. Camb.*, 31, 125.
- KENNAWAY, N. M., AND KENNAWAY, E. L.—(1936) *Ibid.*, 36, 236.
- KLOTZ, M. O.—(1939) *Amer. J. Cancer*, 35, 38.
- 'Leicester, Atmospheric Pollution in. A Scientific Survey.' London: H.M. Stationery Office, 1945.
- LYNCH, K. M., AND SMITH, W. A.—(1935) *Amer. J. Cancer*, 24, 56.
- 'Miners' Phthisis Medical Bureau, Report of the. for the 3 years ended July 31, 1935.' Pretoria: Union of South Africa Government Printer, 1936.
- MONTGOMERY, H.—(1935) *Arch. Derm. Syph.*, 32, 218.
- Idem, AND WAISMAN, M.—(1941) *J. Invest. Derm.*, 4, 365.
- NEUBAUER, O.—(1947) *Brit. J. Cancer*, 1, 192.
- OWENS, J. S.—(1923) *Trans. med. Soc. Lond.*, 45, 79.
- PEACOCK, P. R.—(1943) *Glasg. med. J.*, 139, 117.
- 'Registrar-General's Statistical Review of England and Wales.' Tables. Text. London: H.M. Stationery Office.
- SEMON, H. C.—(1945) *Proc. Roy. Soc. Med.*, 38, Section of Dermatology, p. 128.
- 'Statistical Bulletin,' Metropolitan Life Insurance Co., New York (1939), 20, 7.
- STEVENSON, T. H. C.—(1923) *Biometrika*, 15, 382.
- STOCKS, P.—(1936) 13th Ann. Rep. Brit. Emp. Cancer Campaign, 232.
- WILLIS, R. A.—(1941) *Med. J. Austral.*, Sept. 6, p. 258.

NOTE: THIS DOCUMENT DID
NOT COME FROM THE FILES

THE ESTIMATION OF TUMOUR SUSCEPTIBILITY IN PURE LINES.

C. C. SPICER.

From the Imperial Cancer Research Fund, London, and the
Department of Biometry, University College, London.

Received for publication August 29, 1947.

It is the purpose of this paper to discuss the problem of assigning measures of cancer susceptibility to inbred strains of mice, selected for their proneness to mammary cancer. The question has been previously discussed by Murray and Hoffman (1941), and the related problem of the assessment of potency of carcinogenic compounds by Irwin and Goodman (1946), Twort and Twort (1939, 1933), Bryan and Shimkin (1941) and Lea (1945). It is suggested that the problem is most clearly examined by using an actuarial function, the "force of mortality," on whose characteristics the mortality of a strain depends. This function, which is usually denoted by the symbol $\mu(x)$, measures the rate at which the members of a life table are dying out. It can be calculated for a single cause,

GG 15288

1 BB 0007279 1