

DRAFT 10/17/80

MALIGNANT MESOTHELIOMA IN CONNECTICUT 1935-1977

by

H. C. Lewinsohn

J. W. Meigs

M. J. Teta

Presentation to the Connecticut Thoracic Society, November 4, 1980

A00572

UCC 004937

I. Introduction and Aims

The combined sex age-adjusted mesothelioma incidence rate for Connecticut was reported in 1977 to have increased ten-fold since 1935.^① Although available statistics might be subject to diagnostic error, the apparent rise was attributed to the increase in the State's 'cumulative asbestos consumption'.^{②③} Complete occupational histories for the cases of mesothelioma were not presented. // A detailed review of the available pathological material by an independent pathologist to investigate the degree of diagnostic certainty was not undertaken. // The present study attempts to determine the role of various etiological factors, such as occupational and environmental asbestos exposures, and includes a review of available pathological material.

A00573

II. Methodology

The CTR has identified 229 cases of malignant mesothelioma as well as 38 other pleural tumors, not mesothelioma, which were diagnosed in the state between 1935 and 1977 (Table 1).

Medical, demographic, and occupational data have been collected for the cases and for the respective spouses of cases diagnosed 1955-1977. Similar information has been gathered for a random sample of approximately 700 decedents (1935-75) aged 20 to 98 years from the Division of Health Statistics of the Connecticut Department of Health Services.

Descriptive epidemiology for this research pertains primarily to the forty-three year time interval. ^{The} ~~Future case~~-control comparisons will ~~include~~ ^{comprise} ~~include~~ cases (215) diagnosed after 1954 (Table 1), and controls (604) whose deaths occurred during this same time period. This procedure yields a case-control ratio of approximately 1:3, while reducing sources of error resulting from the limited occupational and medical data prior to 1955 and the lack of awareness of mesothelioma associated with this earlier time period (1935-1954). The Price and Lee City Directories were searched for job title and name of specific employer or industry for cases, controls and spouses at 1, 10, 20, 25, 30, 40 and 50 years prior to date of diagnosis, death or until the subject was less than twenty years old. An occupational history search was attempted for spouses to coincide with these intervals for their corresponding cases.

1970 U.S. Census industrial and occupational codes (U.S. Dept. of Commerce, 1971) ⁽⁵⁾ were assigned to the employment information ascertained from medical histories, death certificates, and City Directories. A computerized

Table 1. Sex, age and diagnostic characteristics of 229 cases of malignant mesothelioma and 38 other pleural tumours diagnosed in Connecticut, 1935-1977

Diagnosis ^a	Males		Females		Total		Mean age	
Pleural mesothelioma	102	(94) ^b	45	(37)	147	(131)	61	(62)
Pleural tumour (other than mesothelioma)	24	(12)	14	(8)	38	(20)	62	(63)
Peritoneal mesothelioma	20	(18)	13	(11)	33	(29)	58	(59)
Mesothelioma at other sites & at unknown sites	29	(21)	20	(14)	49	(35)	52	(54)
Total	175	(145)	92	(70)	267	(215)	59	(60)

^aWHO (1976) ^(c)

^bNumbers in brackets refer to the period 1955-1977

A00575

list of job and industry titles has been developed for all cases, spouses, and controls (1955-1977). This will form the basis for classification of study subjects into asbestos exposure categories for future case-control comparisons (sample from listing Fig. 1).

A00576

UCC 004941

III. Results

Descriptive Epidemiology

The following ~~descriptive~~ results are based upon ^{available} ~~the information~~ ^{information} ~~available from the~~ CTR and will require adjustment following ~~the~~ completion of ^{the} slide review. Using the 1950 US population as a standard, the age-adjusted incidence rate for mesothelioma in Connecticut is 2.1/million for the years 1935-1977. Rates per 100,000 population increased for both sexes, but there was a rapid rise from about 1960 for males (Fig. 2). The male-female ratio is approximately 2:1; the mean age at diagnosis is fifty-nine years. Average survival time from date of diagnosis to date of death is ten months.


~~Ten~~ ^{Ten (10)} ~~cases~~ ^{New London Labor Market Area (LMA)} were reported in ~~the LMA (New London)~~ ^(New London) where shipyards are located, and ~~five~~ ^{5 (50%)} of these were identified between 1975 and 1977 (Fig. 3).

The geographical distribution of mesothelioma shows evidence of urban clusters in the 5 largest cities (Bridgeport, Hartford, Waterbury, Stamford, New Haven) where 30 % of the cases resided at time of diagnosis (Fig. 3). Since these locations have comprised 20-30% of Connecticut's population (1940-1977), the suggested urban effect may reduce to a factor of population density. These 5 large cities and New London are ^{all} ~~also~~ centers for ^{LMA's} ~~Labor Market Areas~~ (LMA's). All 6 areas exhibit a similarly increasing age-adjusted incidence rate for males. The Stamford LMA's mesothelioma rate shows an unexplained sharp increase since 1965 (Fig. 4).

A00577


Fig 2

AGE-ADJUSTED INCIDENCE RATES OF MESOTHELIOMA IN CONNECTICUT BY YEAR OF DIAGNOSIS AND SEX




A00578

GEOGRAPHICAL MAPPING OF CASES OF MESOTHELIOMA (1935 - 1977) BY TOWN OF RESIDENCE AND 1960 LABOR MARKET AREA



AGE-ADJUSTED INCIDENCE RATES OF MESOTHELIOMA FOR CONNECTICUT MALES BY YEAR OF DIAGNOSIS AND LABOR MARKET AREA



SLIDE REVIEW

Malignant mesothelioma is a very rare disease, ^{with} only 2 cases per million population expected annually. ⁷ Resistance to the acceptance of mesothelioma as a disease entity persisted until the late 1960's. ⁸ Wagner's widely publicized association of mesothelioma with Cape Crocidolite asbestos exposure created the potential for the introduction of diagnostic bias. ⁹

Positive diagnosis of mesothelioma is often complicated by its confusion with other forms of cancer.. (Vidone discussion of simulators).

The controversy over criteria for positive diagnosis is well documented in the literature. ¹⁰ Most experts agree, however, that a full autopsy is required to positively distinguish diffuse mesothelioma of the pleural or peritoneum from other primary or secondary neoplasms. ¹¹

We examined, for all CTR reported cases of mesothelioma and for all pleural tumors other than mesothelioma (1935-77), the histological basis for diagnosis (Tables 2,3). For 12% (32) of the cases, no tissue was examined at time of diagnosis, while the origin of available material varied. Overall, ⁽¹⁹³⁵⁻⁷⁷⁾ only 105 (44%) ~~105~~ (44%) of the 238 deceased cases are known by the CTR to have been autopsied, but only 83 of the autopsied cases were microscopically confirmed. The percentages for all categories, except peritoneal mesothelioma, are low ranging from 33% to 40% autopsied,

These results suggested the advisability of a ^{case} ~~slide~~ review. Dr. Romeo Vidone, chief pathologist of St. Raphael's Hospital, is presently studying the medical records (except occupational data) and slides we have obtained for cases diagnosed after 1954. Cooperation was sought from 37 hospitals, of which 30 have thus far provided us with the materials requested.

^{preliminary}
(Summary of Dr. Vidone's Findings)

A00581

Table 2: Basis for Diagnostic Evaluation
for cases of Malignant Mesothelioma
and Pleural Tumors other than
Mesothelioma Diagnosed in Connecticut, 1935-1977

<u>Diagnosis^a</u>	<u>Tissue^b Available</u>	<u>%</u>	<u>No Tissue Available</u>	<u>%</u>	<u>Total</u>	<u>%</u>
Pleural Mesothelioma	136	(93)	11	(7)	147	(100)
Pleural Tumor (other than mesothelioma)	27	(71)	11	(29)	38	(100)
Peritoneal Mesothelioma	31	(94)	2	(6)	33	(100)
Mesothelioma at other sites and unknown sites	41	(84)	8	(16)	49	(100)
Total	235	(88)	32	(12)	267	(100)

^aWHO (1976)

^bspecimen from biopsy, frozen section, surgery, autopsy, D and C

A00582

Table 3: Frequency of Autopsy for Cases of Malignant Mesothelioma and Pleural Tumors other than Mesothelioma Diagnosed in Connecticut, 1935-1977

(1) Diagnosis ^a	(2) Autopsy, Microscopic Evidence of Cancer	(3) Autopsy, Gross Evidence, Unknown Whether Cancer	(4) Autopsy, but only Indirect Evidence of Cancer	(5) Autopsy, no Report	(6) No Autopsy	(7) Unknown	(8) Number Dead	Autopsy ^b %
Pleural Mesothelioma	43		1	7	62	18	134	40
Pleural Tumor (other than mesothelioma)	8	1		3	13	11	37	35
Peritoneal Mesothelioma	21				2	1	28	89
Mesothelioma at other sites and unknown sites	11			2	21	5	39	33
Total	83	1	1	12	98	35	238	44%

^aWHO (1976)

^b $\frac{\text{cols}(1) + (2) + (3) + (4) + (5)}{\text{Col } (8)}$

A00583

IV. Discussion

Although this study was initially undertaken to discover whether it would be feasible to identify the environmental factors responsible for the increase of malignant mesothelioma in the State of Connecticut, it has suffered from lack of detail in available records. Major deficiencies in the data are the low autopsy rate for the pleural mesotheliomata and the inadequacy of the pathological material available and used for diagnostic purposes. ^{Sc. report}

⁽¹²⁾ Felton has pointed out several needs for post-mortem review. A problem may arise in workers' compensation adjudication where a decision has to be made, years to decades after initial work exposure, in those instances where death has resulted from pulmonary cancer or a thoracic or peritoneal mesothelioma. It is therefore important, in such cases, to be able to confirm by the presence of asbestos related effects that there has been exposure to respirable asbestos fibers.

A further need for autopsy arises to confirm the diagnosis of malignant mesothelioma made on limited biopsy material obtained during life. This is important for workers' compensation purposes and also to improve the epidemiological data required to investigate the etiological factors involved with this disease.

Valuable information can be gained from a comparison between radiographic appearances and histological changes in the lung. Such information will help to eliminate present diagnostic uncertainties.

It has been our experience that occupational histories are not routinely obtained and included in hospital records. In our search through records stored at the CTR (i.e. hospital records), we found job titles for only 17% of the mesothelioma cases. If we exclude from this group those designated as "retired", "housewife", or "student", the figure is reduced to 12%. We were only able to ascertain type of industry in 7% of these cases. The sample size for these statistics was 220 cases, whose diagnoses were between 1955 and 1977. A recently passed Ct. statute, whose method of implementation is under study, will hopefully alleviate the paucity of employment data from medical records.

A00504

UCC 004949

Two other areas of epidemiological interest which are not routinely recorded for patients admitted with suspected malignant disease are smoking habits and hobbies and or part-time activities.

(Comment on the significance of the Slide Review)

It is evident that until the physicians investigating cases of malignancy appreciate the importance of occupational factors~~-on~~ and environmental factors ~~on-the-etiology-of-these-diseases,-no~~ (lifestyle, habits, hobbies, part-time activities^e, detailed^a job histories) , any attempt to apportion blame to any particular factor for disease causation or promotion will be severely hampered. It is particularly essential in the case of a rare tumor, which may present diagnostic difficulties, to obtain sufficient tissue for study before a diagnosis is made. Whenever a tumor is found and an association is suspected with a particular occupational or environmental factor, every attempt should be made to document all relevant facts and to subsequently verify the diagnosis by means of a full autopsy.

A00585

20.

References

Bruckman, L., Rubino, R. A., and Christine, B. (1977a) Asbestos and Mesothelioma in Connecticut. APCA Journal, 27, 121-126

Bruckman, L. (1977b) A Study of Airborne Asbestos Fibers in Connecticut. Paper presented at the Workshop on Asbestos: Definitions and Measurement Methods, National Bureau of Standards, Gaithersburg, Maryland

Bruckman, L. and Rubino, R. A. (1978) Monitored Asbestos Concentrations in Connecticut. APCA Journal, 28, 1221-1226

Price and Lee City Directory (1890-1977) The Price and Lee Company, New Haven, Connecticut

U. S. Department of Commerce, Bureau of the Census: 1970 Census of the Population. Alphabetical Index of Industries and Occupations (1971), Washington, D. C., U. S. Government Printing Office

World Health Organization: ICD-O International Classification of Diseases for Oncology, (1976) Geneva, Switzerland

Felton —

A00586

UCC 004951



Hilton C. Lewinsohn, Corporate Medical Director, Raybestos-Manhattan, Inc.
and Lecturer, Yale School of Epidemiology and
Public Health

(address: Raybestos-Manhattan, Inc.,
100 Oakview Drive,
Trumbull, Connecticut 06611, U. S. A.)

J. Wister Meigs, Director, Connecticut Cancer Epidemiology Unit and
Clinical Professor of Epidemiology, Yale School of
Epidemiology and Public Health

(address: 30 College Street,
New Haven, Connecticut 06520, U. S. A.)

Mary Jane Teta, Associate in Research, Connecticut Cancer Epidemiology
Unit, Yale School of Epidemiology and Public Health

(address: 30 College Street,
New Haven, Connecticut 06520, U. S. A.)

AUG 5 1967