

300 Osgangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. David Hanstein  
P. O. Box 2352  
Leucadia, CA 92024

Dear Mr. Hanstein:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

 THANKS for PARTICIPATING  
IN THE SAFETY  
PROGRAM.

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Mack Tighe  
28303 Golden Meadow Drive  
Rancho Palo Verdes, CA 90274

Dear Mr. Tighe:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Dean Quirion  
5 Stobie Street  
Winslow, ME 04901

Dear Mr. Quirion:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Charles Turner  
ARCO Independence

Dear Mr. Turner:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Jerry Thomas  
ARCO Independence

Dear Mr. Thomas:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Ms. Carol Taylor  
2634 Superior Street  
Bellingham, WA 98226

Dear Ms. Taylor:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Richard Stinson  
13682 Sioux Road  
Westminster, CA 92683

Dear Mr. Stinson:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Russell Santos  
P. O. Box 030512  
Ft. Lauderdale, FL 33303

Dear Mr. Santos:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Robert Rush  
ARCO Spirit

Dear Mr. Rush:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Robert Nims  
ARCO Independence

Dear Mr. Nims:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Ms. Elizabeth Parrent  
P. O. Box 713  
Castine, ME 04421

Dear Ms. Parrent:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Kevin Garnier  
753 W. 36th Street  
San Pedro, CA 90731

Dear Mr. Garnier:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec



October 4, 1991

Mr. David Tothill  
ARCO Sag River

Dear Mr. Tothill:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

*Mark*  
Mr. Mark Tighe  
28303 Golden Meadow Drive  
Rancho Palo Verdes, CA 90274

Dear Mr. Tighe:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Svetko Lisica  
57000 Zadar  
Jurja Krizanica 25  
Yugoslavia, Europe

Dear Mr. Lisica:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Ms. Jean Holmes  
3-E Bass Street  
Neumarket, NH 03857

Dear Ms. Holmes:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Clifford Collier  
ARCO Independence

Dear Mr. Collier:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Michael Burke  
315 Sweetman Road  
Ballston SPA, NY 12020

Dear Mr. Burke:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Scott Williams  
16142 Twinkle Circle  
Huntington Beach, CA 92649

Dear Mr. Williams:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Salvador Valencia  
ARCO Independence

Dear Mr. Valencia:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. David Heebner  
2343 East Pole Road  
Everson, WA 98247

Dear Mr. Heebner:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Billy Hamblen  
1001 SE 16th Street, Apt 8  
Fort Lauderdale, FL 33316

Dear Mr. Hamblen:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Michael Doyle  
ARCO Spirit

Dear Mr. Doyle:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Glenn Crenshaw II  
1659 Cyrene Drive  
Carson, CA 90746

Dear Mr. Crenshaw:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

ARCO Marine, Inc.  
300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Mr. Larry Cadle  
3301 Hughes Ln 82  
Dickinson, TX 77539

Dear Mr. Cadle:

While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec

300 Oceangate  
Post Office Box 22617  
Long Beach, California 90801-5617  
Telephone 213 590 4400  
TWX 910 341 6829



October 4, 1991

Ms. Kristie Black  
635 6th Street North  
Shelby, MT 59474

Dear Ms. Black:


While you were on board the ARCO Independence from August 2-7, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, which involves riding the eight AMI vessels which are known to be constructed with some asbestos bearing materials, a summary report will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Independence showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeter (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Independence 22 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab ranged from less than .00017 to .009 f/cc.

The complete report is on board the Arco Independence and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

  
Jay C. Kitchener  
Manager, Safety, Health &  
Environmental Protection

JCK/cec