

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



PLAINTIFF'S
EXHIBIT
ARCO 14

October 29, 1991

Mr. Gary Pierce
1659 E. Bakerview Road
Bellingham, WA 98226

Dear Mr. Pierce:

While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Ms. Arlene Johnson
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Ms. Johnson:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. James Valerio Sr.
115 Norwood Avenue
Long Branch, NJ 07740

Dear Mr. Valerio:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Timothy Fellows
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Fellows:

While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Ms. Tamara Thorenson
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Ms. Thorenson:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Michael Mann
436 Yale Avenue
Bacaville, CA 95687

Dear Mr. Mann:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Suong Tra
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Tra:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Robert Wenz
P. O. Box 3450
Valdez, AK 99686

Dear Mr. Wenz:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec



October 29, 1991

Mr. Michael Sullivan
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Sullivan:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Marvin Sims
3500 Wise Avenue
Long Beach, CA 90810

Dear Mr. Sims:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Robert Nazario
14430 NE 25th Ave., Apt. L301
Bellevue, WA 98007

Dear Mr. Nazario:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Gerald Murray
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Murray:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec



October 29, 1991

Mr. John Morgan
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Morgan:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Leo Mire
4460 Flamingo
Beaumont, TX 77705

Dear Mr. Mire:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Michael Lay
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Lay:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Fredrick Jolly
14221 441st Place., SE
North Bend, WA 98045

Dear Mr. Jolly:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Thomas Hilken
424 Brookside Place
Crawford, NJ 07016

Dear Mr. Hilken:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Mark Hawkins
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Hawkins:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Federico Gonzales
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Gonzales:

While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Kenton Feeney
P. O. Box 213, Celtic Cove
Beals, ME 04611

Dear Mr. Feeney:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Blaine Christeson Jr.
1704 Thomasville Road, Apt. #162
Tallahassee, FL 32303

Dear Mr. Christeson:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Ms. Victoria Boutilier
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Ms. Boutilier:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec



October 29, 1991

Mr. Ozabe Banks Sr.
Arco Marine, Inc.
Vessel: ARCO Spirit

Dear Mr. Banks:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Mark Shuler
Arco Marine, Inc.
Vessel: ARCO Juneau

Dear Mr. Shuler:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Lawrence Kelly
3137 Valemoor Drive
Palm Harbor, FL 34685

Dear Mr. Kelly:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Ronald Hooker
149 Prestwick Court
Vallejo, CA 94590

Dear Mr. Hooker:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. James Hayden
8 Lily Pond Drive
Camden, ME 04843

Dear Mr. Hayden:

While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Jerald Dyrek
RR2 Box 1798
Washington, IA 52353

Dear Mr. Dyrek:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. John Carroll
4613 Versailles Avenue
Plano, TX 75075

Dear Mr. Carroll:

While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Michael Connolly
24 Old County Road
Scarboro, ME 04074

Dear Mr. Connolly:


While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



October 29, 1991

Mr. Richard Vidoli
5357 Bayshore Avenue
Cape Coral, FL 33904

Dear Mr. Vidoli:

While you were on board the ARCO Juneau from August 29 to September 4, 1991, Clayton Environmental Services, Inc., was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Juneau showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the Arco Juneau 20 samples were taken and analyzed to determine the presence of airborne asbestos fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne asbestos levels ranging from .00023 to .002 f/cc.

The complete report is on board the Arco Juneau and available for your review. If you would like any additional information, please feel free to call me at (213) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health &
Environmental Protection

JCK/cec