

Ten Cases of Angiosarcoma of the Liver in Shawinigan, Quebec

Fernand Delorme, M.D., and Gilles Thériault, M.D.

Ten cases of angiosarcoma of the liver have been diagnosed in Shawinigan, Quebec since 1955. All have occurred in men who worked in a vinyl chloride polymerizing plant. Cigarettes and alcohol do not seem to be associated with this tumor. The amount of vinyl chloride to which these people were exposed, according to information obtained through questionnaires, appears elevated. Angiosarcoma of the liver is accompanied by a fibrosis of the liver which may precede its appearance. In describing the Canadian cases of angiosarcoma, this study attempts to shed more light upon the causal relationship between vinyl chloride monomer and angiosarcoma of the liver and thus to provide some clues to understanding the pathogenesis of this disease.

In January 1974, three cases of angiosarcoma of the liver were discovered among workers of a chemical plant.¹ This finding appeared abnormal and the agent suspected of being responsible for the disease was identified as vinyl chloride monomer (VCM). Since then, many reports on the occurrence of this cancer among vinyl chloride workers have appeared around the world²⁻⁵ and the United States Occupational Safety and Health Administration has enacted a new threshold limit value for concentration of this chemical in the air of the working environment.⁶

Ten cases of angiosarcoma of the liver were diagnosed among workers of a vinyl chloride polymerizing plant in Shawinigan, Quebec. This is the largest number of cases ever to take place in a single plant. The present study describes occupational history, drinking and smoking habits and some histological features of these Canadian cases. Data on some of these Canadian cases jointly with cases of angiosarcoma from the United States have already been described in another article by one of the authors.⁷ Since more accurate and complete information was retrieved on the Canadian cases, it is being presented separately in the present article.

From the Department of Pathology, Centre Hospitalier Régional de la Mauricie, Shawinigan, Quebec (Dr. Delorme) and the Department of Social and Preventive Medicine, Laval University, Ste-Foy, Quebec (Dr. Thériault).

Methods

Case Finding. — A search for cases of angiosarcoma of the liver beginning as far back as 1955 and extending over two decades was carried out by scrutinizing files of the regional hospitals in the Shawinigan vicinity. All cases of primary tumors of the liver were collected and the histological material (slides, biopsy reports, autopsy reports) were reviewed for the presence of angiosarcoma. To confirm the diagnosis of angiosarcoma of the liver, the cases were submitted to consultants from the United States familiar with this disease.

Occupational History. — The questionnaires used to obtain a detailed occupational history of each case culled information concerning past and present residence, medical history, smoking and drinking habits along with past occupations and employers in chronological order. The answers to most of questionnaires were given by spouses of the men who died from the disease. In one instance, however, the answers were given by the sister and brother-in-law, and in another, by the worker himself. Two former co-workers of the men who died of an angiosarcoma of the liver confirmed and completed the information obtained through the families.

Plant Description. — The Shawinigan vinyl chloride plant started operating in 1943. At that time, it produced and polymerized vinyl chloride monomer. Later in the 1950s, it stopped producing VCM and kept operating the polymerization unit. The workforce varied from 100 to 350 workers. At the moment, it employs 250 men. Production of PVC has fluctuated with time. Present production is estimated at 50 million pounds annually. Exposure to VCM in the working atmosphere, though not documented, is believed to have been very high in the past. Sharp improvement has taken place in recent years and it is estimated at less than 3 ppm at the moment.

Results

Table 1 shows the age at death, the smoking habits and alcohol consumption for the ten cases of angiosarcoma of the liver diagnosed in the Shawinigan region over the last two decades. All

Table 1.

Case No.
1
2
3
4
5
6
7
8
9
10
T

were males, the average of 55.

It is somewhat of a tumor; five were moderate or less per cigarette per moderate smokers or less.

The men were very light drinkers; bottle of beer regularly. As be associate workers.

Past disease: the development of history of hepatic trouble.

Table 2: vinyl chloride being directed they worked for an average.

Table 3. — Sex, Age at Death, Smoking Habits and Alcohol Consumption of Ten Cases of Angiosarcoma of the Liver in a Canadian Community.

Case No.	Sex	Age at Death	Smoking Habits			Estimation of Cigarette Consumption		Alcohol Consumption	
			Smoker	Ex-smoker	Non-smoker	No. of Cigarettes per Day	No. of Years Smoked	No. Bottles of Beer Per Week	No. of Years Drinking
1	male	41	x			20	27	6	4
2	male	43	x			12	27	1	25
3	male	42			x	—	—	1	25
4	male	48	x			10	30	—	—
5	male	56	x			10	42	1	40
6	male	51	x			40	33	4	5
7	male	52	x			25	34	21	27
8	male	53	x			25	36	18	20
9	male	53	x			20	35	4	10
10	male	61		x		25	43	—	—
T		50.5							

were males, the age at death ranging from 41 to 61 years with an average of 50.5 years.

It is somewhat doubtful that smoking has much to do with this type of tumor. Out of the ten cases, one man was a nonsmoker, five were moderate smokers, smoking an average of 20 cigarettes or less per day, and the remaining four smoked one pack of cigarettes per day or more. The tumor appeared earlier in the moderate and nonsmokers (x age 47 years) than in the heavy smokers (x 54 years).

The men who developed an angiosarcoma of the liver were very light drinkers. Only two men out of ten drank more than one bottle of beer per day. None of the ten men drank wine or liquor regularly. As with cigarette smoking, drinking does not appear to be associated with an angiosarcoma of the liver in vinyl chloride workers.

Past diseases of the liver may also be ruled out as potentiating the development of angiosarcoma of the liver since the medical history of the ten cases showed no previous liver diseases nor hepatic troubles.

Table 2 shows the length of time the men were exposed to vinyl chloride either by working in the polymerization unit or by being directly exposed to vinyl chloride. It also shows the occupations they performed and whether they worked overtime. They worked for periods ranging from 11 to 29 years in the plant with an average of 21 years and 4 months. Direct exposure to vinyl

chloride monomer was obviously shorter, ranging from 5 years 3 months to 26 years with an average of 16 years 10 months.

From the job descriptions, it can be seen that most of the men worked either as operators or maintenance workers, or both. Seven men worked as "reactor cleaners," an operation which seems to have been the worst in exposing people to vinyl chloride monomer. One woman said of her husband's exposure, "He went down into the autoclaves many times. Once, they had to take him out unconscious."

It is difficult to get a fair idea of the amount of vinyl chloride monomer to which these people were exposed. Many spouses (six out of ten) insisted upon mentioning that their husbands worked overtime.

As shown in Table 3, the time elapsed between the first exposure to vinyl chloride monomer and the diagnosis of the disease ranges from 11 years to 28 years with an average of 20.5 years. This concurs with previous findings that an angiosarcoma of the liver can develop after a minimum exposure of four years, but that actual manifestation does not occur for at least ten years.⁴

Residence history revealed that all workers lived within a two mile radius of the VCM plant since the beginning of their employment at that plant. However, no particular pattern in the residential history could be identified that could lead to finding environmental factors common to everyone.

Past occupational history showed that all these workers spent

Table 2. — Occupational History of Ten Cases of Angiosarcoma of the Liver in a Canadian Community.

Case No.	Time Exposed				Occupation in Contact with Vinyl Chloride	Worked Overtime
	Worked in the Plant		Worked in Contact with VC			
	Years	Months	Years	Months		
1	11		11		Reactor cleaner, operator polymerization and	
2	14	2	14	2	Maintenance, operator, midnight, pipe fitter	yes
3	20	5	20	5	Operator, foreman	yes
4	22		22		Reactor cleaner, operator, foreman, water cooling unit	yes
5	17	5	5	3	Reactor cleaner, water cooling unit	yes
6	25	4	23	4	Reactor cleaner, maintenance, midnight, reactor	yes
7	26	6	25	6	Reactor cleaner, operator VCM fabrication and polymerization unit, machine shop	
8	25	8	5	8	Reactor cleaner, maintenance, machine shop	
9	29		26		Furnace operator, pipe fitter, maintenance	yes
10	18	10	14	9	Reactor, furnace operator, packaging	
Σ	21	4	18	10		

Table 3. — First Exposure, Year of Diagnosis, Survival Time, Type of Tumor, Accompanying Benign Liver Lesions, and Means of Diagnosis of Ten Cases of Angiosarcoma of the Liver in a Canadian Community.																
Case No.	First Exposure	Disease Found	Survived Months	Type of Tumor				Other Lesions Accompanying								Means of Diagnosis
				Scissoidal	Nodulo-papillary	Sarcomatous	Cavernous	Thickening of Liver Capsule	Portal Fibrosis	Sclerosis of the Wall of Portal Vein	Proliferation of Fibrous Tissue	Dilatation of Sinusoids	Proliferation of Sinusoidal Cells	Regenerated Liver Cells		
1	1944	1955	2		x		x	x				x	x		Biopsy	
2	1943	1957	1	x	x	x	x	x	x	x	x	x	x	x	Autopsy	
3	1941	1962	4				x				x	x	x	x	Autopsy	
4	1945	1967	8	x	x		x	x	x	x	x	x	x	x	Biopsy-autopsy	
5	1944	1968	3				x			Not enough non-tumoral material to appreciate					Biopsy	
6	1947	1971	2				x	x	x	x	x	x	x	x	Autopsy	
7	1946	1972	7	x		x	x	x	x	x	x	x	x	x	Biopsy	
8	1961	1973	6				x	x	x	x	x	x	x	x	Autopsy	
9	1946	1974	2				x	x	x	x	x	x	x	x	Biopsy-autopsy	
10	1953	1976	alive													
11			4.1			x	x	x	x	x	x	x	x	x	Biopsy	

the greater part of their active lives working for the same employer. The longest period anyone worked outside the plant was four years.

Once the tumor is diagnosed, the survival period is very short. In the present study, the longest survival period was eight months, the average 4.7 months.

It must be pointed out that angiosarcoma is not the only liver pathology found in vinyl chloride monomer workers.⁹⁻¹⁰ There are two types of benign lesions — sclerosis and cellular transformation. Sclerosis manifests itself by a thickening of the capsule of the liver, a portal fibrosis with proliferation of the bile ducts and lymphocytic infiltrate, a sclerosis of the wall of the portal vein and centrilobular veins and a perisinusoidal fibrosis with sinusoid dilatation. Cellular transformation shows a sinusoidal cell proliferation with atypic, dysplastic and reactional liver cells. The tumor itself is a malignant angiosarcoma type tumor appearing in four definite histological patterns: a cavernous pattern, a sinusoidal pattern, a nodulo-papillary pattern and a sarcomatous pattern. In our experience, the cavernous pattern is the most constant and frequent one (Table 3).

Discussion

The present study is an epidemiological and a histological description of the ten cases of angiosarcoma of the liver found in the region of Shawinigan, Quebec. Although the selection of cases was made from hospital records, it happened that all were men who worked in the same factory — a vinyl chloride polymerization plant. This type of tumor being very rare, a causal relationship between the exposure to vinyl chloride and the development of an angiosarcoma of the liver is established.

The present study reveals that the exposure necessary to develop the cancer must be relatively high. Cleaning reactors and operating production units were occupations carried out in an environment with a high concentration of vinyl chloride monomer. Furthermore, it is possible that in the 1940's, the men worked more than 40 hours a week. To this, one must add overtime work, another contributing factor. It will be necessary to assess with as much accuracy as possible the exposure of these people to vinyl

chloride monomer to discover a dose-response relationship between the chemical and the disease.

It seems that benign non-tumoral reactions are the first response of the liver to the chemical. Eventually they give way to malignant transformations.⁷⁻¹¹ Obviously, more studies will be necessary to discover whether the benign and malignant lesions are two steps of a unique phenomenon or two different diseases.

References

1. Creech JL and Johnson MN: Angiosarcoma of liver in the manufacture of polyvinyl chloride. *IOA* 16:150-151, 1974.
2. Lee FI and Harry DS: Angiosarcoma of the liver in a vinyl chloride worker. *Lancet* 1316-1318, June 29, 1974.
3. Block, Bradford J: Angiosarcoma of the liver following vinyl chloride exposure. *JAMA* 229:53-54, 1974.
4. Réty J, Sauvage, Tuallion, and Habrard A: Le 3ième cas français d'angiosarcome hépatique chez un ouvrier du chlorure de vinyle. *Arch Mal Prof* 551-555, 12 déc. 1975.
5. Center for Disease Control. Morbidity and Mortality. Weekly Report. 23(8):1-2, Feb 1976.
6. Department of Labor, Occupational Safety and Health Administration: Exposure of vinyl chloride. *Federal Register* 39(194):35890-35898, Part II, Oct 4, 1974.
7. Makk L, Delorme F, et al: Clinical and morphologic features of hepatic angiosarcoma in vinyl chloride workers. *Cancer* 37:149-163, 1976.
8. Laszlo M, Delorme F, et al: Clinical and morphologic features of hepatic angiosarcoma in vinyl chloride workers. *Cancer* 37(1), Jan 1976.
9. Delorme F and Makk L: Angiosarcome du foie chez les ouvriers ayant été en contact prolongé avec le chlorure de vinyle. Description morphologique des lésions. *Union Méd Canada* 104:834-841, 1973.
10. Delorme F and Makk L: Angiosarcome du foie chez les ouvriers ayant été en contact prolongé avec le chlorure de vinyle. Description morphologique des lésions. *Revue franc gastroent* 119 May 1976.
11. Thomas LB and Popper H: Pathology of angiosarcoma of the liver among vinyl chloride-polyvinyl chloride workers. *Ann NY Acad Sci* 246:268-277, 1975.
12. Thomas LB, Popper H, et al: Vinyl-chloride-induced liver disease. *New Engl J Med* 292:17-22, 1975.
13. Gedigk P, Muller R, et al: Morphology of liver damage among polyvinyl chloride production workers. A report on 51 cases. *Ann NY Acad Sci* 246:278-285, 1975.
14. McDonald RB: Primary carcinoma of the liver. *Arch Int Med* 99:266-279, 1957.
15. Popper H and Thomas LB: Alterations of liver and spleen among workers exposed to vinyl chloride. *Ann NY Acad Sci* 246:172-194, 1975.

Angiosarcoma of the Liver in Shawinigan, Quebec/Delorme and Thériault

Please
As
to
of
Willia

The
two a
withou
only c
amosi
perloc
in Pl
thicke
nor w
there
pleur
relati
pleur
appei
asbes

In
cigar
autop
repor
In 19
geno
lowe
foun
Sir
woul
cigar
diser
obse
som
less
ings
Ann
of p
fro
Colle
Phila
Jour

SEP 23, 1980

FILE NUMBER : 150

AUTHOR(S) : HOLMBERG, B.; KRONEVI, T.; WINELL, M.

TITLE : THE PATHOLOGY OF VINYL CHLORIDE EXPOSED MICE.

PUBLICATION SOURCE : ACTA VET. SCAND. 17:328-342. 1976.

LANGUAGE : ENG.

COUNTRY : SWE.

757

VC
Animal studies
Biology

***** EXPERIMENTAL DESIGN *****

THIS INHALATION STUDY USED OUTBRED ALBINO NMRI MICE (12 WKS. OF AGE). TWENTY FOUR MICE (12 MALES AND 12 FEMALES) GROUP WERE EXPOSED TO A VC CONCENTRATION OF 50 PPM (FOR 52 WKS.) OR 500 PPM (FOR 26 WKS.), 6HRS./DAY, FIVE DAYS/WK. CONTROL GROUPS OF 24 ANIMALS WERE RUN CONCURRENTLY WITH EACH SERIES, HOWEVER, ONLY ONE GROUP WAS EXPOSED TO AIR IN AN INHALATION CHAMBER. IN THE EXPERIMENTAL GROUPS FOUR MICE OF EACH SEX WERE SACRIFICED AND SUBJECTED TO PATHOLOGICAL EXAMINATION ON THE 26th WEEK AND THE REMAINING ANIMALS EXAMINED WHEN MORIBUND OR DEAD. FOUR MICE IN THE CONTROL GROUP WERE SACRIFICED ON THE 52nd WEEK. THE VC CONCENTRATION IN THE DARK, CYLINDRICAL METAL INHALATION CHAMBER WAS ANALYZED BY G.C. REGULARLY. THE MICE WERE WEIGHED (TOTAL BODY WEIGHT) EVERY TWO WEEKS.

***** KEYWORDS *****

VINYL CHLORIDE; INHALATION; MICE; PATHOLOGY; BENIGN TUMORS; ALVEOGENIC ADENOMAS; LUNG; HEMANGIOSARCOMA; FAT TISSUE; CARCINOGENICITY; BLOOD VESSEL; RUPTURE; TELANGIECTASIS; LIVER; MAMMARY ADENOCARCINOMA; PARARINIAL HEMANGIOSARCOMA; PARAINTESTINAL HEMANGIOSARCOMA; RHABDOMYOSARCOMA; HEMOCOELIA; BROWN FAT; SUBCUTANEOUS; SUBPERITONEAL; VC ANIMAL STUDY.

***** ABSTRACTS *****

IN THIS INHALATION STUDY MICE WERE EXPOSED TO VC AT 50 OR 500 PPM FOR 6 HR./DAY, 5 DAYS/WK, FOR 52 AND 26 WKS., RESPECTIVELY. THERE WAS A DECREASE IN BODY WEIGHT IN THE EXPOSED GROUPS AFTER 40 WEEKS. THREE OF THE 40 CONTROL SERIES ANIMALS WERE OBSERVED WITH TUMORS: ONE ANIMAL HAD A MAMMARY ADENOCARCINOMA WITH PULMONARY METASTASIS; THE SECOND ANIMAL HAD DISGERMINOMA OF THE OVARY; AND THE THIRD DIED WITH A RETICULUM CELL SARCOMA OF THE SPLEEN AND MESENTERIC LYMPH NODES.

IN THE MICE EXPOSED TO 50 PPM AND SACRIFICED ON THE 26th WEEK ALVEOGENIC ADENOMAS WERE OBSERVED IN 2/4 MALES, HOWEVER NO TUMORS WERE FOUND IN THE FEMALES. THE REMAINING ANIMALS IN THE LOW EXPOSURE GROUP THAT WERE NECROPSIED BETWEEN THE 36th AND 56th WEEK ALL BORE TUMORS. IN THE WHOLE GROUP 13/24 HAD ALVEOGENIC ADENOMAS. THE LUNG TUMORS WERE SINGLE OR MULTIPLE WITH A FEW FOCI WITH

DIAMETERS UP TO 0.2 cm. THERE WERE 14/24 ANIMALS WITH SUBPERITONEAL HEMANGIOSARCOMA WHICH WERE FOUND IN THE PARARENAL, PARAINTESTINAL OR IN THE PELVIC/CAUDAL PART OF THE ABDOMINAL CAVITY. THE PARARENAL HEMANGIOSARCOMAS ADHERED TO THE KIDNEY BUT DID NOT INFILTRATE THE KIDNEY. IN THE SUBCUTANEOUS TISSUE OF THESE ANIMALS THERE WAS ONE HEMANGIOSARCOMA IN THE BROWN FAT, FOUR HEMANGIOSARCOMAS IN OTHER SITES AND A MAMMARY ADENOCARCINOMA. ONE OF THE ANIMALS THAT BORE AN ALVEOGENIC ADENOMA ALSO HAD A SECONDARY HEMANGIOSARCOMA IN THE LUNG ORIGINATING FROM A PRIMARY SUBCUTANEOUS HEMANGIOSARCOMA OF THE THROAT. THERE WAS ONE ANIMAL THAT REVEALED RHABDOMYOSARCOMA AS WELL AS ALVEOGENIC ADENOMA WHEN NECROPSIED ON THE 51st WEEK. EIGHT OF THE 24 MICE HAD RUPTURE OF HEMANGIOSARCOMAS CAUSING HEMOCOELIA TO OCCUR.

THE ANIMALS EXPOSED TO 500PPM VINYL CHLORIDE WERE NO LONGER EXPOSED AFTER THE 26TH WEEK, SINCE THEY WERE CONSIDERED IN POOR CONDITION. ALL MICE IN THIS SERIES SHOWED ALVEOGENIC ADENOMA WITH LUNG TUMORS IN ALL LOBES WITH 10 OR MORE FOCI. THE DIAMETERS OF THE FOCI WERE AROUND 0.9 cm. THE SUBPERITONEAL HEMANGIOSARCOMAS WERE FOUND IN THE PARARENAL (3 CASES) AND PELVIC/CAUDAL PART OF THE ABDOMINAL CAVITY (8 CASES). IN THE SUBCUTANEOUS TISSUE THERE WERE FOUR CASES OF MAMMARY ADENOCARCINOMA AND ONE CASE OF HEMANGIOSARCOMA IN THE BROWN FAT. IN THE ORGANS OF THESE ANIMALS THERE WAS REPORTED ONE CASE OF HEMANGIOSARCOMA OF THE LIVER, AND ONE CASE OF ADENOMA OF THE KIDNEY. THE OTHER PATHOLOGICAL CHANGES OTHER THAN NEOPLASTIC LESIONS WERE TELANGIECTASIS OF THE LIVER IN TWO ANIMALS AND HEMOCOELIA IN TWO ANIMALS EXPOSED TO THE HIGHER CONCENTRATION.

THE AUTHORS CONSIDER THESE RESULTS TO BE INDICATIVE OF A DOSE-RESPONSE RELATIONSHIP SINCE THE FREQUENCY OF ANIMALS BEARING TUMORS IS 100% IN THE 500PPM GROUP AND 71% IN THE 50PPM GROUP. THE 500PPM GROUP ALSO HAD MORE MULTIPLE TUMORS OVER THE WHOLE LUNG AND LARGER LUNG TUMOR FOCI THAN THE LOW DOSE GROUP. THE IMPORTANCE OF HEMANGIOSARCOMAS IN VARIOUS ORGANS AND FAT TISSUE OF SUBCUTANEOUS AND SUBPERITONEAL AREAS INDICATES THE ROLE OF THE BLOOD VESSELS AND FAT TISSUE IN THE BIOLOGICAL ACTIVITY OF VC.