


The lower limit of size of particle visible to the naked eye cannot be stated definitely. It depends not only upon the individual, but also upon the shape and color of the particle and upon the intensity of light. Under



Compiled by W. G. Frank and Copyrighted.

FIG. 1. SIZES AND CHARACTERISTICS OF AIR-BORNE SOLIDS

ideal conditions a particle 10 microns, or even less, may be recognized, while under less favorable conditions it may be difficult or even impossible to recognize a particle of 50 microns. The lower limit of visibility should, therefore, be considered as within a range (as above) or optimum conditions stated.

Mineral particles, such as grains of sand, bits of rock, volcanic ash, or fly-ash, can be transported long distances under unusual circumstances. Thus, the dust storms of 1935 in the Kansas district resulted in vast amounts of fine top soil being thrown high into the air. Solar illumination

TABLE 1. APPROXIMATE LIMITS OF INFLAMMABILITY OF SINGLE GASES AND VAPORS
IN AIR AT ORDINARY TEMPERATURES AND PRESSURES^a

GAS OR VAPOR	LOWER LIMIT VOLUME IN PER CENT	HIGHER LIMIT VOLUME IN PER CENT	GAS OR VAPOR	LOWER LIMIT VOLUME IN PER CENT	HIGHER LIMIT VOLUME IN PER CENT
Acetaldehyde.....	4.0	57.0	Hexane.....	1.2	6.9
Acetone.....	3.0	13.0	Hydrocyanic acid.....	5.6	40.0
Acetone ^b	2.5		Hydrogen.....	4.1	74.0
Acetylene.....	2.5	80.0	Hydrogen sulphide.....	4.3	45.5
Acetylene ^b	2.3		Illuminating gas.....	5.3	31.0
Allyl alcohol.....	2.4		Iso-amyl alcohol.....	1.2	
Ammonia.....	16.0	27.0	Iso-butane.....	1.8	8.4
Amyl alcohol.....	1.2		Iso-butyl alcohol.....	1.8	
Amyl chloride.....	1.4		Iso-pentane.....	1.3	
Amylene.....	1.6		Iso-propyl acetate.....	1.8	7.8
Benzene.....	1.4	6.7	Iso-propyl alcohol.....	2.6	
Benzine.....	1.1		Methane.....	5.3	14.0
Blast-furnace gas.....	35.0	74.0	Methane ^b	5.0	15.0
Butane.....	1.9	8.4	Methyl acetate.....	3.1	15.5
Butyl acetate (30 C).....	1.7	7.6	Methyl alcohol.....	6.7	36.0
Butyl alcohol.....	1.7		Methyl bromide.....	13.5	14.5
Butylene.....	1.7	9.0	Methylbutyl ketone.....	1.2	8.0
Carbon disulphide.....	1.2	50.0	Methyl chloride.....	8.0	19.0
Carbon monoxide.....	12.5	74.0	Methyl cyclohexane.....	1.2	
Croton aldehyde.....	2.1	15.5	Methyl ethyl ether.....	2.0	10.1
Cyclohexane.....	1.3	8.3	Methyl ethyl ketone.....	1.8	10.0
Cyclopropane.....	2.4	10.3	Methyl formate.....	5.0	23.0
Decane.....	0.7		Methyl propyl ketone.....	1.5	8.5
Dichlorethylene.....	9.7	12.8	Natural gas.....	4.8	13.5
Diethyl selenide.....	2.5		Nonane.....	0.8	
Dioxan.....	2.0	22.0	Octane.....	0.9	
Ethane.....	3.2	12.5	o-Xylene.....	1.0	6.0
Ethyl acetate.....	2.0	11.5	Paraldehyde.....	1.3	
Ethyl alcohol.....	3.3	19.0	Pentane.....	1.4	7.8
Ethyl bromide.....	6.7	11.2	Propane.....	2.4	9.5
Ethyl chloride.....	4.0	14.8	Propyl acetate.....	1.8	8.0
Ethylene dichloride.....	6.2	15.9	Propyl alcohol.....	2.5	
Ethylene.....	3.0	29.0	Propylene.....	2.0	11.1
Ethyl ether.....	1.8	36.5	Propylene dichloride.....	3.4	14.5
Ethyl formate.....	2.7	16.5	Propylene oxide.....	2.1	21.5
Ethyl nitrite.....	3.0		Pyridine (70 C).....	1.8	12.4
Ethylene oxide.....	3.0	80.0	Toluene.....	1.3	6.7
Furfural (125 C).....	2.1		Vinyl ether.....	1.7	27.0
Gasoline.....	1.4	6.5	Vinyl chloride.....	4.0	22.0
Heptane.....	1.0		Water gas.....	6 to 9	55 to 70

*Limits of Inflammability of Gases and Vapors, by H. F. Coward and G. W. Jones (*U. S. Bureau of Mines, Bulletin No 279, 1939*).

^bTurbulent mixture.

as far east as Boston was affected noticeably and particles as large as 40 to 50 microns were actually carried half way across the continent before they settled out. In similar manner volcanic ash has been carried even further. It is not surprising, therefore, that fly-ash from furnace gases, cement dust and the like, can be carried for considerable distances and