Monsanto

3/4/69

FROM (NAME & LOCATION) E. S. Tucker - R&D Laboratories - 1700 So. Second

DATE March 4, 1969

cc: R. E. Keller - Res. 1
E. P. Wheeler - GO

SUBJECT AROCLOR - WILDLIFE:
INCINERATION OF NCR PAPER

REFERENCE

TO W. R. Richard - GO

It has been proposed from the observed distribution of polychlorinated biphenyls in tissue samples taken from marine organisms that global eco-distribution of these materials occurs via aerial transport (wind currents, airborne particulate matter, etc.).

A qualitative experiment has been performed with the objective of determining if Aroclor in N.C.R. paper is destroyed or volatilized when the paper is disposed of by burning.

The experiment was carried out in the following manner. Approximately 3g. (1 sheet) of paper (CB 15# white, 8-1/2 X 11, long grain, F-3468-4-24-64, The N.C.R. Company) was placed in the incinerator chamber of the apparatus shown in Figure I.

FIGURE I



At this point the vacuum was turned on and the paper ignited. After the paper had completely burned, the hexane in each trap, and hexane extracts of the ash and the walls of the incinerator chamber were analyzed for the presence of Aroclor by EC/GC. Hexane samples from trap #1 were also analyzed after extraction with base and acid to remove any oxidized materials.

DSW 372351

42

IN-10 REV. 11-65

STLCOPCB4098884

Conclusions:

Under the conditions of this experiment:

1. Aroclor is easily volatilized when N.C.R. paper is burned.

2. Aroclor undergoes little, if any, decomposition when burned.

3. Unfortunately, it appears that significant air pollution can occur via burning of N.C.R. paper or other Aroclor containing materials even under more strenuous conditions.

E. S. Tucker

db

DSW 372352

41



FIG. 2

DSW 372353

40

STLCOPCB4098886



FIG 3

DSW 372354

STLCOPCB4098887

DSW 372355



Fig. 4

STLCOPCB4098888