

A REVIEW OF CHLORINATED BIPHENYL CONTAMINATION IN NATURAL WATERS

GILMAN D. VEITH and G. FRED LEE

Water Chemistry Laboratory
University of Wisconsin, Madison, Wis. 53706, U.S.A.

(Received 10 November 1969)

Abstract—A review of the present state of knowledge concerning environmental contamination by organochlorine compounds indicates that the chlorinated biphenyls may be one of the more widespread contaminants. The significance of this contamination has not yet been evaluated due, in part, to the lack of systematic analytical procedures for the quantitative and qualitative determinations of the components of chlorinated biphenyl mixtures. The possibility of the chlorinated biphenyls producing serious errors in chlorinated pesticide analyses is emphasized.

INTRODUCTION

CHEMICAL determinations of the types and concentrations of chlorinated pesticides in natural waters have become increasingly important to fish management, wildlife ecology, and water pollution control. Based on gas chromatographic (GLC) analyses of environmental samples, pp'DDE is generally considered to be one of the more widespread contaminants. However, implicit in every GLC analysis is the possibility that unresolved and/or unidentified components which have specific retention volumes similar to that of the desired substance may introduce serious errors in the interpretation of the chromatogram. Increasing numbers of investigators have reported the presence of organochlorine compounds in environmental samples which cannot be identified as the common chlorinated pesticides. Although the unidentified compounds have been attributed from presumptive evidence to the metabolites of common pesticides, plasticizers, or chlorinated biphenyls, proper chemical identifications have not been reported. Therefore, when studying natural water systems which receive industrial wastes, it may be more correct to consider unidentified compounds as chlorinated organic contaminants until confirmation techniques are developed.

Within the last three years, the chlorinated biphenyls (or polychlorinated biphenyls-PCB) have received considerable study. The preliminary evidence indicates that this group of organochlorine compounds may be widespread in the environment. For example, WIDMARK (1967) reported that mass-spectrum analyses of compounds eluting from GLC of fish and bird extracts indicated that unknown organochlorine compounds were present which behaved as chlorinated biphenyls of higher chlorine content. More recently, mass numbers and chlorine numbers per molecule for each peak in the chromatograph of commercial chlorinated biphenyl were reported (JENSEN *et al.*, 1968). However, to our knowledge, no spectral data or other confirmatory data for positive structural identification of these compounds have ever been published.

With this improper basis of identification, a number of other publications have followed which identify unknown compounds as chlorinated biphenyls. Since mixtures of chlorinated biphenyls may contain greater than 20 compounds which can only be

resolved by GLC with great difficulty, assignment of unidentified peaks to one of the chlorinated biphenyls has been accomplished by comparison of a chlorinated biphenyl with the nearest relative retention time. For example, HOLDEN and MARSDEN (1967), who had earlier reported the presence of unknown compounds in fish (HOLDEN, 1966), reported the presence of chlorinated biphenyls in seals with no supporting data other than relative retention data from GLC columns.

ROBINSON *et al.* (1967) reported compounds of unknown identity in marine organisms which gave GLC retention times longer than pp'DDE and microcoulometric response. The authors indicated that the chromatographic characteristics of the unknowns suggested they were not metabolites of the "cyclodiene-type" pesticides as has been proposed by ROBURN (1965).

HOLMES *et al.* (1967) attributed unknown peaks from GLC to the chlorinated biphenyls and noted that the substances seemed to accumulate in organisms associated with natural waters (seals, fish and birds) to a greater extent than in domestic animals. This work reported relative retention data with no supporting data.

RISEBROUGH *et al.* (1968 a) and RISEBROUGH *et al.* (1968 b) reported organochlorine compounds in fish, birds and bird eggs which were attributed to the chlorinated biphenyls. With no supporting data and despite the absence of structural confirmations for the chlorinated biphenyls, the investigators reported concentrations of the unknowns (as chlorinated biphenyls) as well as ratios of DDT to chlorinated biphenyls. From these ratios, it was concluded that a chlorinated biphenyl fallout pattern existed around industrial centers and that air transport of the contaminants was significant.

KOEMAN *et al.* (1969) presented GLC chromatograms of extracts from environmental samples as well as mass numbers and numbers of chlorine atoms per molecule for each peak of the chromatogram. Although no spectral data were presented, the mass number data strongly suggested that chlorinated biphenyls were present. This paper also presented the results of the analyses for each unknown as being "present" or "absent" in a particular sample, rather than attempting to quantitate the amount of material present.

THE CHLORINATED BIPHENYLS

Since contamination of the environment by chlorinated biphenyls has been implicated in the brief review presented, the need for the environmental scientist to become familiar with the present state of knowledge concerning these molecules is apparent. Mixtures of chlorinated biphenyls have been manufactured since 1929 and present registered trademarks for commercial brands include: Arochlor (Monsanto), Chloretol (Allis-Chalmers), Dykanol (Cornell-Dubilier Division, Federal Pacific Electric), Inerteen (Westinghouse), Noflamol (Wagner Electric), Pyramol (G. E.), Therminol (Monsanto) in the U.S.; and Clophen (I. G. Farbenindustrie A. G., Germany), Fencolor (Caffaro, Italy), Kannechlor (Kanegafuchi Chemical Co., Japan), Pyralene (Prodelec, France), and Soval (Russia) (HUBBARD, 1964). The chlorinated biphenyls have been used as plasticizers for adhesives, polyvinyl acetates, acrylic resins, in sealing compounds, pipe joint compounds, printing ink, rubber manufacturing, high pressure-high temperature lubricants, fungicidal insulations, anti-corrosion, nail coatings, water proofings, diffusion pump fluid, stabilization of polymers,

dielectrics, heat preventing coats (MONSANTO O/F *et al.*, 1955).

The toxicity a has not been indicated that m "safe" is 1.0 mg highly chlorinated with chlorinated reported repeat biphenyls (M effects of Arocl cited) to the el 5460 to be non

More recent biphenyls to h tion, and that factor of 1000. dieldrin and D

However, in with organoch possibility the biological cor review of the other metabol enzyme induc great as pp'D facts make tol cules have th concentration

Some cont studies of org biphenyl. In analyses. Ho have been in make the pr of chemical l

For exam step which is residues (Ea was treated It was concl nitration ste

dielectrics, heat transfer media, coax-cable insulations, paints and varnishes, ice-preventing coatings, and is an evaporation retardant for pesticide applications (MONSANTO O/PL-306; HUBBARD, 1964; HOPKINS and HOFFMAN, 1955; SULLIVAN *et al.*, 1955).

The toxicity and sublethal effects of the chlorinated biphenyls to biological systems has not been thoroughly evaluated (REICHEL, 1969). Inhalation tests on animals indicate that maximum concentrations of chlorinated biphenyls as vapor which are "safe" is 1.0 mg/m³ for the lower chlorinated biphenyls and 0.5 mg/m³ for the more highly chlorinated biphenyls (HUBBARD, 1964). Continuous or repeated skin contact with chlorinated biphenyl leads to a condition called "chloracne" which has been reported repeatedly with industrial workers handling products containing chlorinated biphenyls (MEIGS *et al.*, 1953). SULLIVAN and HORNSTEIN (1953) found no toxic effects of Arochlor 5460 on flies, but the chlorinated terphenyls were toxic (no levels cited) to the elm leaf beetle (DUDA, 1957). DEONIER *et al.* (1946) found Arochlor 5460 to be non-toxic at 0.1 ppm (24 hr) to the larvae of *Anopheles quadrimaculatus*.

More recently, LICHTENSTEIN *et al.* (1969) reported the toxicity of the chlorinated biphenyls to houseflies (*D. melanogaster*) decreased with increasing degree of chlorination, and that the chlorinated biphenyls appear to be less toxic than dieldrin by a factor of 1000. The chlorinated biphenyls were found to increase the toxic effect of dieldrin and DDT on houseflies.

However, in addition to investigating the sources of environmental contamination with organochlorine compounds, the environmental scientist is concerned with the possibility that these organochlorine molecules may exhibit detrimental effects on biological communities at sublethal levels. KUPFER (1967) has presented an extensive review of the effects of chlorinated pesticides on steroid hydroxylase induction and other metabolic systems. Also, RISEBROUGH *et al.* (1968 b) have demonstrated that enzyme induction by the chlorinated biphenyls (Arochlor 1262) was five times as great as pp'DDE or technical DDT as determined via oestradiol degradation. These facts make tolerance limits set by bioassay less significant since organochlorine molecules have the capacity to produce physiological effects in organisms at sublethal concentrations.

ANALYTICAL PROCEDURES

Some controversy has been generated over the analytical procedures utilized in studies of organochlorine contaminants which have been attributed to the chlorinated biphenyl. In general, the procedures are similar to typical chlorinated pesticide analyses. However, chemical reactions such as dehydrohalogenation and nitration have been introduced prior to electron capture gas chromatography in attempts to make the procedures more selective for the chlorinated biphenyls. It is the addition of chemical treatment steps which has caused disagreement in the literature.

For example, after extract clean up, JENSEN and WIDMARK (1967) used a nitration step which is similar to that used in removing DDT and metabolites from toxaphene residues (ERRO *et al.*, 1967; KAWANO *et al.*, 1969). Briefly, the concentrated extract was treated with a mixture of H₂SO₄ and HNO₃ (conc. 1:1, v/v) at 0°C for 5 min. It was concluded incorrectly that only PCB and BHC isomers are unaffected by the nitration step. RISEBROUGH *et al.* (1969) have reported the use of similar procedures.

- HOLDEN A. V. and MARSDEN K. (1967) Organochlorine pesticides in seals and porpoises. *Nature, Lond.* 216, 1274-1276.
- HOLMES D. C., SIMMONS J. H. and TATTON J. O'G. (1967) Chlorinated hydrocarbons in British wildlife. *Nature, Lond.* 216, 227-229.
- HOPKINS T. L. and HOFFMAN R. A. (1955) Effectiveness of dilon and certain candidate synergists against DDT resistant house flies. *J. econ. Ent.* 48, 146-147.
- MURRAY H. L. (1964) Chlorinated biphenyl and related compounds. *Kirk-Othmer Encyclopedia of Chemical Technology*, 2nd edn. Vol. 5, 289-297.
- JENSEN S., NIELSEN H. and WIDMARK G. (1968) *The OECD Study of Analysis of PCB*. Stockholm, December, mimeo.
- JENSEN S. and WIDMARK G. (1967) *Organochlorine Residues, OECD Preliminary Study*. Stockholm, September, mimeo.
- KAWANO Y., BEVEMER A., BECKMAN H. and ENRO F. (1969) Studies on the effect of sulfuric-fuming nitric acid treatment on the analytical characteristics of toxaphene. *J. Ass. Official Analyt. Chem.* 52, 167-172.
- KOFMAN J. H., DE BRAUW M. C., TEN NOEVER and DE VOS R. H. (1969) Chlorinated biphenyls in fish, mussels and birds from the Rhine River and the Netherlands coastal area. *Nature, Lond.* 221, 1126-9.
- KUPFER D. (1967) Effects of some pesticides and related compounds on steroid function and metabolism. *Residue Rev.* 19, 11-30.
- LEUTHOLDEN F. P., SCHUB Z. K. R., FUHRMANN T. W. and LIANG T. T. (1969) Biological interaction between planktivores and insecticides. *J. econ. Ent.* 62, 761-765.
- MIRK J. W., ALLEN J. J. and KURTIN B. L. (1952) Chloracne from an unusual exposure to Arochlor. *J. Am. med. Ass.* 154, 1417-1418.
- MINISANTO COMPANY, *Technical Bulletin O/PL-306*, St. Louis, Missouri.
- RICHIE, W. L. (1969) Chief, section of Chemistry. Personal communication from the Patuxent Wildlife Research Center, Laurel, Maryland.
- REYNOLDS L. M. (1969) Polychlorobiphenyls (PCB's) and their interference with pesticide residue analysis. *Bull. Env. Contam. Toxicol.* 4, 128-143.
- RISBROUGH R. W., HUGGETT R. J., GRIFFIN J. J. and GOLDBERG E. D. (1968 a) Pesticides: Transatlantic movements in the Northeast Trades. *Science* 159, 1233-1234.
- RISBROUGH R. W., RICHIE P., HERMAN S. G., BEAKALL D. B. and KIRVIN M. N. (1968 b) Polychlorinated biphenyls in the global ecosystem. *Nature, Lond.* 220, 1098-1102.
- RISBROUGH R. W., RICHIE P. and OLCOTT H. S. (1969) Current progress in the determination of the polychlorinated biphenyls. *Bull. Env. Contam. Toxicol.* 4, 192-201.
- ROBINSON J., RICHARDSON A., CRABTREE A. N., COULSON J. C. and POTTS G. R. (1967) Organochlorine residues in marine organisms. *Nature, Lond.* 214, 1307-1311.
- ROBURN J. (1965) A simple concentration cell technique for determining small amounts of halide ions and its use in the determination of residues of organochlorine pesticides in extracts of animal and plant tissue. *Analyst* 90, 467-475.
- SULLIVAN W. N. and HORNSTEIN I. (1953) A comparison of evaporation rates and toxicity to house flies of lindane and lindane-chlorinated polyphenyl deposits. *J. econ. Ent.* 46, 882-894.
- SULLIVAN W. N., HORNSTEIN I. and YROMANS A. H. (1955) Improved deposits for controlling insects outdoors. *J. econ. Ent.* 48, 153-155.
- VEITH G. D. and LEE G. F. (1969) *Chlorinated Organic Contaminants in the Milwaukee River*. Progress Report to the Wisconsin Department of Natural Resources, Madison, October, mimeo.
- WIDMARK G. (1967) Pesticide Residues. *J. Ass. Official Analyt. Chem.* 50, 1069.