

## OCCUPATIONAL CANCER

OF THE URINARY BLADDER IN DYESTUFFS OPERATIVES AND OF THE LUNG IN  
ASBESTOS TEXTILE WORKERS AND IRON-ORE MINERSGEORGLANA M. BONSER, M.D., J. S. FAULDS, M.D., AND M. J. STEWART, LL.D.  
*University of Leeds and Cumberland Infirmary, Carlisle, England*

The study of occupational cancer has proved to be a valuable method of learning some aspects of the biology of cancer in its wider sense. This form of cancer is an unwitting experiment in man, but until it can be reproduced in animals at will, doubt must always exist as to the exact nature of the carcinogen involved and of its mode of action. The time lag between the observation that cancer occurs in those engaged in certain occupations and the successful induction of the disease in animals may be very long. For example, Percival Pott<sup>12</sup> described cancer of the scrotum in chimney sweeps in 1773, but it was not until 1922 that Passey<sup>13</sup> induced cancer of the skin of mice by means of an ether extract of soot. Similarly, in 1893, Rehn<sup>14</sup> described cancer of the bladder occurring in workers engaged in the manufacture of aniline dyes, but it was not until 1937 that Hueper and his co-workers<sup>15</sup> were able to induce bladder cancer in dogs by means of ingestion of the dye intermediates. So far no attempt has succeeded in reproducing the lung cancers of the Schneeberg and Joachimstal miners (Haerting and Hesse, 1879;<sup>7</sup> Šíkl, 1930<sup>16</sup>), though this disease has been known for centuries. The induction of a few local sarcomas in rats following injection of cobalt metal (Heath, 1954<sup>8</sup>) might be a stimulus to further experimental work.

We are well advised to be ready to recognize new forms of hazard in any industrial process, but it is even more important to advise industry of the dangers that are likely to be inherent in the manufacture or use of suspected or proved carcinogens. This was certainly done when the new plants for effecting the catalytic cracking of oils were set up (Symposium on a Cancer Control Program, 1951).<sup>21</sup>

As such a variety of occupational cancers is now recognized, this presentation is confined to making a classification of the most important and to discussion of those with which the authors are most familiar (Table 1). It seems hardly necessary to say that, before it can be claimed that any particular form of cancer has an industrial origin, it must be shown that the incidence is significantly higher in exposed workmen than it is in the general population. This is not always easy, as morbidity statistics of cancer in most countries are not available, hospital statistics being quite fallaciously high. Secondly, in sites where there is a high natural incidence, any occupational risk tends to be masked.

Received for publication August 31, 1954; accepted, September 23, 1954.

Dr. Bonser is Reader in Cancer Research, University of Leeds, and Morbid Anatomist to St. James Hospital, Leeds; Dr. Faulds is Pathologist, Cumberland Royal Infirmary, Carlisle; and Dr. Stewart is Emeritus Professor of Pathology, University of Leeds.

TABLE 1

LIST OF SOME SUBSTANCES CAUSING CANCER, INDUSTRY CONCERNED, AND SITE OF CANCER

| Substance               | Industry                         | Site of Cancer            | Proof      |              |
|-------------------------|----------------------------------|---------------------------|------------|--------------|
|                         |                                  |                           | Industrial | Experimental |
| Minerals                |                                  |                           |            |              |
| Arsenic                 | Sheepdip manufacture and usage   | Skin, lung                | Doubtful   | 0            |
| Chromium (ore)          | Electroplating, tanning, dyeing  | Respiratory, usually lung | Doubtful   | 0            |
| Nickel                  | Chemical manufacturers           | Respiratory               | Doubtful   | 0            |
| Silica                  | Stone blasting                   | Lung                      | Doubtful   | 0            |
| Asbestos                | Miners and users                 | Lung                      | +          | 0            |
| Iron (hematite)         | Miners                           | Lung                      | +          | 0            |
| Beryllium               | Fluorescent lamps                | Lung, bones               | 0          | +            |
| Unknown                 | Pitchblende miners               | Lung                      | +          | 0            |
| Oils                    |                                  |                           |            |              |
| Lubricating             | Mule spinners                    | Skin                      | +          | +            |
| Petroleum               | Refiners                         | Skin                      | +          | +            |
| Pitches                 |                                  |                           |            |              |
| Soot                    | Chimney sweeps                   | Skin                      | +          | +            |
| Tar                     | Road workers                     | Skin                      | +          | +            |
| Rays                    |                                  |                           |            |              |
| Actinic                 | Farmers, sailors                 | Skin, exposed sites       | +          | +            |
| Röntgen                 | Scientific personnel             | Skin, exposed sites       | +          | +            |
| Luminous paint (radium) | Dial painters                    | Skin, exposed sites       | +          | +            |
| Organic chemicals       |                                  |                           |            |              |
| Dye intermediates       | Chemical manufacturers and users | Bladder                   | +          | +            |

A very significant fact has been observed by many workers and has been emphasized by Kennaway (1950).<sup>14</sup> The "amount" or incidence of cancer in any given population does not vary greatly, as regards either sex or race, but the proportions of the different "forms" or sites of cancer may vary greatly. For example, there is a high rate of carcinoma in organs peculiar to the female, in contrast with the high rates in lung and stomach in the male. Similarly, primary carcinoma of the liver is high in the Javanese in Sumatra, but insignificant in European populations. "There appears to be a general law that when in a given population the incidence of cancer in one particular organ is markedly increased compared with another population, there is then a compensating decrease in the

incidence of cancer in a number of the other organs" (Cramer, 1936).<sup>6</sup> A constant watch must be kept on "amounts" of cancer if proportions are being altered by industrial forms of the disease.

#### OCCUPATIONAL CANCER IN THE DYESTUFFS INDUSTRY

Why should 1 in 10 or even more of men engaged in the manufacture and usage of certain dye intermediates (aromatic amines) develop tumors of the bladder? These workers are exposed by inhalation, ingestion and absorption of these compounds through the skin and yet the tumors are located in the bladder. This striking observation has fired the imagination of many workers and we think we are justified in stating that we are beginning to understand the reasons. The knowledge thus gained has altered our conception of the range and mode of action of chemical carcinogens in an interesting way.

An extremely good survey of tumors of the urinary bladder in workmen engaged in the manufacture and use of certain dyestuff intermediates in the British chemical industry has recently been published by Case and his associates (Table 2). They have shown that although there is a small spontaneous incidence of bladder tumors in the general population (2½ times greater in males than in females and rare under 55 years of age) the risk is 30 times greater in chemical workers and the age of onset is on an average 15 years earlier. Manufacturers, users and purifiers are affected in descending order of severity, the noxious substances being 2-naphthylamine, benzidine and possibly auramine and magenta. There is no evidence that aniline is incriminated. As the induction period is constant at 15 to 20 years, the age at onset of the disease is dependent upon the age at entry of a man into the industry. This is linked with the fact that it is the length and not the severity of exposure that matters, individual susceptibility being a marked feature. There is no special tendency for industrial tumors to be

TABLE 2  
OCCURRENCE OF BLADDER TUMORS IN MANUFACTURERS AND USERS OF DYESTUFF  
INTERMEDIATES (CASE ET AL.)  
Population at Risk = 4622 Men

|                                                  |                                                        |
|--------------------------------------------------|--------------------------------------------------------|
| Expected deaths in comparable general population | 3-5                                                    |
| Actual deaths in industry                        | 127                                                    |
| Actual cases in industry                         | 262                                                    |
| Risk in industry                                 | 30 times that of general population                    |
| Chemical exposure                                | Benzidine<br>Beta-naphthylamine<br>Alpha-naphthylamine |
| Age of onset                                     | 15 years earlier than in general population            |
| Induction period                                 | 15-20 years                                            |
| Degree of malignancy                             | Comparable to tumors of bladder in general population  |
| Survival after diagnosis                         | 20 per cent survived 10 years                          |
| Exposure                                         | Length, not severity, affects induction                |
| Individual susceptibility                        | A marked feature of unknown cause                      |

more or less malignant than spontaneous ones. This is a killing disease, only 20 per cent of all cases having survived more than 10 years from the first recognition of the disease.

In 1935, certain alterations in plant and technic were made and these have reduced the risk just significantly, but another 243 cases (over and above the 262 traced cases in a population of 4622 men) may be expected among British workers engaged in the manufacture of chemicals during the next few years. In 1945 and 1950, more drastic plant alterations were made and we can expect these to bear fruit in about 1965.

Meanwhile, we need to be ever vigilant in recognizing as occupational tumors any that may occur in other industries in which the dyestuff intermediates are in use. Case and Hosker (1954)<sup>2</sup> have recently demonstrated an occupational bladder-tumor risk in the rubber industry in England and Wales. Attention was drawn to this by the finding of a high incidence of bladder tumors in the hospital records of a County Borough which is an important center of the rubber industry. It is probable that the risk was introduced into the industry in 1928, when a certain antioxidant began to be used. The latter was known to have caused bladder tumors among the men in the chemical industry who manufactured it, and it contained about 2.5 per cent of free naphthylamine, both alpha and beta isomers. The treatment of the rubber "mix" containing the antioxidant volatilized quantities of naphthylamines. It needed a most laborious and careful search of records of many kinds to establish this risk on a sound statistical basis and it is satisfactory to report that the manufacturers of the antioxidant immediately ceased its manufacture and the users discontinued its use and destroyed old stocks.

Of great interest to the experimental worker is the search for the reasons for the localization of the tumors in the urinary tract, or perhaps the wider conception of localization along the routes of excretion would be preferable. If 2-naphthylamine is administered by feeding to dogs, all the animals will eventually develop tumors of the bladder, though the number and location will vary in individuals (Hueper and Wolfe, 1937;<sup>10</sup> Bonser and her co-workers, 1951<sup>11</sup>). If 2-naphthylamine is fed to mice, hepatomas or benign cholangiomas will occur. If fed to rats or rabbits, occasional benign papillomas of the bladder will be seen. If benzidine is fed to dogs, bladder tumors occur after many years (Spitz, personal communication), but if injected into rats, hepatomas and acoustic gland tumors will be found (Spitz, Maguigan and Dobriner, 1950).<sup>12</sup> If injected into mice, hepatomas only will occur (Bonser and her associates, unpublished observation) (Table 3). Such localization of tumors suggests that 3 important factors must be at work: (a) species variations in biologic response; (b) excretion of the original chemical or its metabolites along a specific pathway; and (c) concentration of these substances in certain sites.

A study<sup>1</sup> of the metabolism of 2-naphthylamine in various species showed that a conversion of the amine to 2-amino-1-naphthol conjugates occurred and that the capacity to form tumors was roughly proportional to the concentration of these substances in the urine. In the dog, the ratio of urinary to serum content of



TABLE 4  
INCIDENCE OF CANCER IN ASBESTOS-TEXTILE WORKERS AT POSTMORTEM EXAMINATION

|                                              | Un-<br>complicated<br>Asbestosis |    | Tuber-<br>culosis and<br>Asbestosis |   | Lung Cancer<br>and<br>Asbestosis |     | Peritoneal<br>Cancer and<br>Asbestosis |   |
|----------------------------------------------|----------------------------------|----|-------------------------------------|---|----------------------------------|-----|----------------------------------------|---|
|                                              | M                                | F  | M                                   | F | M                                | F   | M                                      | F |
| Number of cases investigated postmortem (72) | 22                               | 18 | 12*                                 | 3 | 12*                              | 2   | 1                                      | 3 |
| Incidence of lung cancer (per cent)          |                                  |    |                                     |   | 26.1                             | 8.7 |                                        |   |
| Average age at death                         | 54(21)†                          |    | 47(12)                              |   | 56(11)                           |     |                                        |   |
| Average age at starting work                 | 31(21)                           |    | 28(12)                              |   | 27(10)                           |     |                                        |   |
| Average period of exposure (years)           |                                  |    |                                     |   |                                  |     |                                        |   |
| Total                                        | 23(21)                           |    | 18(12)                              |   | 26(10)                           |     |                                        |   |
| Prior to 1931                                | 14(10)                           |    | 14(7)                               |   | 12(10)                           |     |                                        |   |
| In cases entering after 1931                 | 5(2)                             |    | 7(5)                                |   |                                  |     |                                        |   |

\* One man had asbestosis, tuberculosis and lung cancer.

† Figures in parentheses represent the number of cases for which the information was available.

and 26 females. In Table 4 the data in regard to the association of asbestosis and cancer are given. Cancer arising primarily in pleurae, lungs or bronchi was present in 26.1 per cent of males and 8.7 per cent of females. This significantly higher incidence in males may be accounted for by the fact that 7 women died before the cancer age, 4 following parturition or abortion and 3 from tuberculosis. There was no significant difference in average age at death, average period of exposure or average age of starting work between males with uncomplicated asbestosis and those with the complication of lung cancer. This is interesting, as in the dyestuff industry death was accelerated by the supervention of bladder cancer. It was noted that the degree of fibrosis was rather less in the lungs with cancer than in those without. Tuberculosis significantly lowered the age at death.

A fair assessment of the period of exposure is difficult in such a series of cases. Taking into account the known fact that after a carcinogen has ceased to be applied to a tissue a latent period is required before the development of the malignant change, it was decided to use the period from the first entry into the factory until death as the period of exposure. This disregards absences from the factory for any reason. It also presupposes that once the fibers have gained access to the lung the carcinogenic process is set in motion, becoming fully developed at the end of the latent period.

Two interesting features deserve mention. The morbid anatomy and histology of the 13 lung cancers of which full notes were available was unusually varied. Four were large tumors situated near the lung hilum, disseminating in characteristic fashion and of adenocarcinomatous or oat-cell type. Seven were situated in the periphery of the lung, usually near the base, 3 disseminating by blood and lymph streams and the other 4 remaining confined to the lung substance. Six of these were adenocarcinomatous or of oat-cell type, 1 being spindle cell and possibly a rhabdomyosarcoma. Small peripheral cancers were described by Raeburn and Spencer (1953)<sup>14</sup> as the likely precursors of the large hilar masses

usually seen at postmortem examination. The peripheral cancers in this series may, therefore, represent a stage prior to hilar involvement. Two tumors were pleural in distribution, 1 disseminating and 1 not. Both were adenocarcinomas, the former having spindle-cell areas.

Attention must also be drawn to the 4 peritoneal cancers, 1 in a male and 3 in females. This is a very high number in such a small series of cases. All these tumors were anaplastic carcinomas, the primary site not being found in any organ. The possibility of an origin from the sequestered bronchial epithelium present in the diaphragmatic areas of hyaloseritis is suggested, but without any clear proof.

No adequate comparable postmortem material from persons working under factory conditions and not exposed to asbestos dust is available with which to compare the incidence of lung cancer in asbestos workers. The incidence here cited is very high and by comparison with the incidence in iron-ore miners must be highly significant. The occurrence of occupational lung cancer in the exposed female is noteworthy.

In connection with this particular form of industrial hazard, it must be emphasized that since the introduction of regulations for the control of asbestos dust in 1931, the amount of exposure of the workers to dust has been enormously reduced and we shall therefore expect to see a disappearance of those severe degrees of pulmonary fibrosis that have been described heretofore and possibly a disappearance of associated cancer. All the present cases that had asbestosis and cancer had been exposed prior to 1931.

## 2. Cancer of the Lung in Iron-Ore Miners

One of the authors (JSF) has had a unique opportunity to study this disease (Stewart and Faulds, 1934).<sup>13</sup> He has personally conducted postmortem examinations on 192 Cumberland hematite miners during the last 20 years and in addition has performed postmortem examinations on 2378 Cumberland males above the age of 20 years, unselected in regard to cancer or lung disease. The difference in incidence of lung cancer in these 2 groups is statistically significant (Table 5). The age of onset is not significantly delayed, most of the miners having entered the mines under the age of 30 years. All the miners' lungs contained much silica and iron at death, but there was no significant difference in the content of either mineral in the lungs with and without cancer. There was, however, an interesting difference in the degree of fibrosis (Table 6) and this has greater significance when a similar occurrence in relation to asbestosis is remembered. All the tumors were classified as oat-cell or squamous cancers.

TABLE 5  
CANCER OF THE LUNG IN IRON-ORE MINERS

|                                | Number | Lung Cancers | Percentage Incidence |
|--------------------------------|--------|--------------|----------------------|
| Iron-ore miners                | 192    | 17           | 8.85 = 9             |
| Cumberland males over 20 years | 2378   | 44           | 1.85 = 2             |

Brone  
Fibro-  
Fibro-  
No fit

Fig  
avail-

Th  
It is -  
silicic  
mont  
which  
million  
than  
exam  
avail  
the lu  
asbest  
chemi  
nature

The  
number  
of cases  
any p  
found  
Now  
in reg  
study  
lation  
ratio  
times

TABLE 6  
MINERAL CONTENT, DURATION OF OCCUPATION AND PULMONARY FIBROSIS IN  
IRON-ORE MINERS

|                                 | Number in Group | Average Age at Death | Average Duration of Occupation (Years) | Average Percentage Mineral Content to Dry Lung |      | Degree of Fibrosis |
|---------------------------------|-----------------|----------------------|----------------------------------------|------------------------------------------------|------|--------------------|
|                                 |                 |                      |                                        | Silica                                         | Iron |                    |
| Bronchial carcinoma             | 15              | 61                   | 32 (11)                                | 1.11                                           | 5.3  | +                  |
| Fibrosis alone                  | 59              | 60                   | 34 (20)                                | 1.15                                           | 6.3  | +++                |
| Fibrosis and tuberculosis       | 75              | 57                   | 28 (48)                                | 1.14                                           | 5.4  | ++                 |
| No fibrosis and no tuberculosis | 28              | 62                   | 28 (10)                                | 0.59                                           | 4.04 | 0                  |

Figures in parentheses represent the number of cases in which the information was available.

The common factor in these 2 types of occupational lung cancer is silica, and it is suggested that this may be the carcinogenic agent. The presence of silica (as silicic acid) causes pulmonary fibrosis, which precedes the initiation of the malignant process. The fibrosis is usually of lesser degree, however, in those lungs in which cancer supervenes. On published evidence the case for an etiologic relationship between nodular silicosis *per se* and lung cancer is much less strong than in silicosis associated with siderosis and in asbestosis. Hueper (1952)\* examined the literature very thoroughly and came to the conclusion that "the available evidence supporting a causal relation between silicosis and cancer of the lung is unimpressive." It seems possible, therefore, that the fibrous nature of asbestos or the presence of hematite dust may exert a modifying effect on the chemical action of the silica or on the fibrotic process itself. Further study of the nature of the carcinogenic process in pneumoconioses is thus obviously needed.

#### SUMMARY

The value to the cancer problem as a whole of the study of occupational cancer in the field and by experiment, together with the time lag between these 2 aspects, is discussed. A short consideration of the total amount of cancer in any population follows, and a classification of the types of recognized occupational cancer.

More detailed consideration of industrial bladder and lung cancer follows. In regard to the former, the extent of the hazard, the nature of the chemicals involved and their metabolism in various species are discussed. In regard to the latter, the occurrence of lung cancer in a group of 72 male and female asbestos textile workers and in 192 hematite miners in Cumberland, England, is described.

#### REFERENCES

1. BONSER, G. M., CLAYSON, D. B., AND JULL, J. W.: An experimental inquiry into the causes of industrial bladder cancer. *Lancet*, 2: 286-287, 1951.
2. BONSER, G. M., CLAYSON, D. B., JULL, J. W., AND PYRAH, L. N.: The carcinogenic properties of 2-amino-1-naphthol hydrochloride and its parent amine 2-naphthylamine. *Brit. J. Cancer*, 6: 412-424, 1952.

his series  
tors were  
cinomas,

and 3 in  
All these  
d in any  
thelium  
without

ing under  
which to  
ace here  
ers must  
exposed

be em-  
tos dust  
mously  
e severe  
ossibly a  
osis d

disease  
exami-  
and in  
d males  
The dif-  
nificant  
having  
d much  
tent of  
ver, an  
greater  
red. All

Incidence

# 1

# 2

3. CASE, R. A. M., AND HOSKER, M. E.: Tumour of the urinary bladder as an occupational disease in the rubber industry in England and Wales. *Brit. J. Prevent. & Soc. Med.*, 8: 39-50, 1954.
4. CASE, R. A. M., HOSKER, M. E., McDONALD, D. B., AND PEARSON, J. T.: Tumours of the urinary bladder in workmen engaged in the manufacture and use of certain dyestuff intermediates in the British chemical industry. Part I. The role of aniline, benzidine, alpha-naphthylamine and beta-naphthylamine. *Brit. J. Indust. Med.*, 11: 75-104, 1954.
5. CLAYSON, D. B.: A working hypothesis for the mode of carcinogenesis of aromatic amines. *Brit. J. Cancer*, 7: 460-471, 1953.
6. CRAMER, W.: The importance of statistical investigations in the campaign against cancer. Report of the 2nd Internat. Congress against Cancer, 1: 441-459, 1936.
7. HAERTING, F. H., AND HESSE, W.: Der Lungenkrebs, die Bergkrankheit in den Schneeberger Gruben. *Vrtijchr. f. gerichtl. Med.*, 30: 206-309; 31: 102-134; 313-337, 1879.
8. HEATH, J. C.: Cobalt as a carcinogen. *Nature*, 173: 522, 1954.
9. HUEPER, W. C.: Occupational and environmental pulmonary cancer with special reference to pneumoconiosis. Proceedings of the 7th Saranac Symposium of Pneumoconiosis, 1952, in press.
10. HUEPER, W. C., AND WOLFE, H. D.: Experimental production of aniline tumors in dogs. *Am. J. Path.*, 13: 656-673, 1937.
11. KENNAWAY, E. L.: The data relating to cancer in the publications of the General Registrar's Office. *Brit. J. Cancer*, 4: 153-172, 1950.
12. PASSET, R. D.: Experimental soot cancer. *Brit. Med. J.*, 2: 539-540, 1922.
13. POTT, P.: Chirurgische Observations Relative to the Cataract, the Polypus of the Nose, the Cancer of the Scrotum, the Different Kinds of Ruptures, and the Mortification of the Toes and Feet. London: T. J. Carnegie, Hower, Clark and Pollins, 1775, p. 65.
14. RAEBURN, C., AND SPENCER, H.: A study of the origin and development of lung cancer. *Thorax*, 8: 1-10, 1953.
15. REHN, L.: Blasengeschwülste bei Fuchsin-Arbeitern. *Arch. f. klin. Chir.*, 50: 588-600, 1895.
16. ŠIKL, H.: Über der Lungenkrebs der Bergleute in Joachimstal (Tschechowiowakei). *Ztschr. f. Krebsforsch.*, 32: 609-613, 1930.
17. STEWART, M. J.: Diagnosis of pulmonary asbestosis at necropsy. *Brit. Med. J.*, 2: 509, 1923.
18. STEWART, M. J., AND FAULDS, J. S.: The pulmonary fibrosis of haematite miners. *J. Path. & Bact.*, 39: 233-253, 1934.
19. STEWART, M. J., AND HADDOW, A. C.: Demonstration of the peculiar bodies of pulmonary asbestosis ("asbestosis bodies") in material obtained by lung puncture and in the sputum. *J. Path. & Bact.*, 32: 172, 1929.
20. STEWART, M. J., TATTERSALL, N., AND HADDOW, A. C.: On the occurrence of clumps of asbestosis bodies in the sputum of asbestos workers. *J. Path. & Bact.*, 35: 737-741, 1932.
21. Symposium on a Cancer Control Program for High-Boiling Catalytically Cracked Oils. *Arch. Indust. Hyg. & Occup. Med.*, 4: 297, 1951.
22. SPITZ, S., MAGUIGAN, W. H., AND DOBRINER, K.: The carcinogenic action of benzidine. *Cancer*, 3: 789-804, 1950.