Plankton Samples Show Pollutants in Marine Organisms

JAN 7 1972

By GERALD LEACH
Knight-Ridder-Washington Post Service

LONDON — Plankton samples throughout the Atlantic Ocean have been found to contain such unexpectedly high levels of industrial pollutants called PCBS that marine biologists are pressing for an immediate international conference.

The discovery, which comes as a surprise, was made by the first survey ever carried out to measure PCBS (polychlorinated biphenyls) in the open oceans.

The survey by the Wood's Hole Oceanographic Institute in Massachusetts, was completed last month after a year spent collecting and analyzing hundreds of samples of plankton and fish from an area stretching from the Antarctic to the arctic and from the Americas to Africa and Europe.

-:-

THE SURVEY found that levels of PCBS — chemical cousins to the pesticide DDT — are roughly 10 to 100 times higher in plankton from the open ocean than from coastal waters, where one would expect greater contamination from river pollution and dumping of industrial wastes. (PCBS are mainly used in paints and varnishes and as electrical insulators: world production is about 50,000 tons a year.)

Actual levels found in plankton — the tiny plants and animals on which all marine life depends for food — ranged from 0.1 to 1.5 parts per million. These are extraordinarily high figures when compared to those for plankton in coastal waters — usually around .01 to .05 ppm. They are also much higher than levels of DDT contamination in the open oceans — usually put at .01 ppm. for plankton.

-:-

ANOTHER SURPRISE finding is that the plankton contained roughly 10 times as much PCBS as did other marine organisms — particularly fish. This contradicts the well-worn ideas that pollutants are progressively concetrated as they pass up the "food chain" — the process in which little fish eat plankton, bigger fish eat the little fish, and so on.

George Harvey, the survey's leader said that, in many food chains, the investigators found that PCBS contamination decreased the higher up the chain they went. All the findings suggest that fish and plankton get most of their PCBS (and by implication other pollutants) directly from the water, not from lowlier animals.

-:-

THIS FINDING in turn has several important implications for marine pollution. One is that PCBS must be reaching the plankton mainly by "wash out" from the atmosphere — suggesting that the biggest primary source is not river pollution or dumping at sea (both easily controllable), but the burning on land of such things as old paint-covered wood.

Harvey estimates that PCBS last more than 100 times as long as DDT (itself highly persistent) — in other words, at least 500 years.

For this reason the ban imposed last March by Monsanto Ltd, the leading manufacturer, on sales of PCBS for anything but totally "captive" use (inside sealed electrical equipment) won't be very effective in lowering ocean contamination by PCBS.

MONS 069887