

FILE NAME: Synkoloid (SYN)

DATE: 1975 SYN014

DOC#: SYN014

DOCUMENT DESCRIPTION: Newspaper article - Asbestos Minerals Detected in
Several Spackling Mixes

relations have been shown for both "viva" and "Duffy" b."

Although they have not yet produced an antigen-antibody reaction for Duffy negative, the scientists believe there is a substance there, and that this substance provides immunity against vivax malaria.

For example, vivax malaria does not exist in West Africa, where 90 per cent of the population is Duffy negative. Most blacks in the United States also are resistant to vivax malaria, and 68 per cent are Duffy negative.

(Because of these facts, the scientists were able to transfer their labora-



—U.S. Medicine photos

Mary McGinniss

Low Standards for Products

of lasers, ranging from low-powered ones producing no harmful effects to those that could cause biological damage from reflected as well as direct exposure. Those which could produce damage must carry warning labels.

Lasers used for medical treatment must, in addition, have a means of measuring the radiation intended to irradiate the human body.

FDA said most of the required standards either already are in use or can be incorporated into existing products at little expense.

Under the regulations, lasers must have protective housing.

Lasers also must have a system for indicating when they are ready for use.

Inverted microscope sequence taken at NIH shows (left to right, from top) invasion of red blood cell by malaria parasite. After attachment by the parasite (arrow) there is marked distortion of the cell, followed by slow invasion.

tory results using monkey malaria to vivax malaria.)

Further, screening showed that all Caucasians tested were Duffy positive.

"This is highly suggestive that the double Duffy negative is the determinant of resistance," Dr. Miller said.

This leads to the possibility—still in the future, however—of developing a neutralizing antibody to the Duffy positive antigen that would block invasion by the malaria parasite.

Such a preventive approach could offer new hope for populations unable to rid themselves of the disease as the United States in effect has through

medication and through environmental controls.

For example, in West Africa, where falciparum malaria—the most virulent of the three common forms, vivax, falciparum and malariae—is prevalent, nearly 100 per cent of pre-school children have the disease. Another form of malaria, Ovale, exists only in West Africa.

If the genetic factor involved in these forms of malaria could be found, a simple vaccine might be developed to stop the growth cycle.

One merozoite, once it penetrates a cell in the laboratory, for example, multiplies into 10 in 24 hours.

Asbestos Minerals Detected In Several Spackling Mixes

RESEARCH TRIANGLE PARK, N.C.—Appreciable amounts of chrysotile or other asbestos minerals were found in samples of spackling and patching compounds analyzed by grantees of the National Institute of Environmental Health Sciences.

The investigators also found substantial amounts of quartz, talc, and other minerals.

Furthermore, air samples obtained during the use of asbestos-containing spackle compounds were analyzed and showed concentrations frequently in excess of the current occupational standard for permissible asbestos exposure levels.

Fiber counts measured during mixing were found to be 7-12 times greater than the current occupational standard. Detectable fiber concentrations were found in adjacent rooms during mixing, and fibers were still suspended in the room air at least 15 minutes after mixing had ceased.

"These findings suggest the possibility of significant asbestos exposure

during home construction and repair," Dr. David P. Rall, NIEHS director, said.

"Additional work by the scientists suggests that members of the entire household or other occupants of a building may also inhale asbestos fibers. This could occur during mixing, sanding, or cleaning up of debris," he said.

Dr. Rall said the investigators, all of the Mount Sinai School of Medicine, New York, have accumulated other evidence indicating that far less than the intense exposure of the work environment is capable of producing asbestos-related disease.

The investigators analyzed 15 representative samples of consumer spackling and patching compounds and 10 of industrial drywall taping compounds for asbestos mineral contents.

Five of the 15 consumer spackling and patching samples contained appreciable amounts of chrysotile or other asbestos minerals, as did 9 of the 10 industrial products.

U.S. Medicine, p. 4, Sept. 1, 1975

Dave Rall - 919-549-8411 ext 301

PLAINTIFF'S
EXHIBIT
SY-5