

Polychlorinated Biphenyls in the Global Ecosystem

by

R. W. RISEBROUGH
P. RIECHE

Institute of Marine Resources,
University of California, Berkeley 94720

D. B. PEAKALL

Division of Ecology and Systematics,
Cornell University, Ithaca,
New York 14850

S. G. HERMAN

Department of Zoology,
University of California, Davis 95616

M. N. KIRVEN

San Diego Natural History Museum,
PO Box 1390, San Diego, California 92112

Polychlorinated biphenyls are widely dispersed in the global ecosystem, and are powerful inducers of hepatic enzymes which degrade oestradiol. Together with other chlorinated biocides, such as DDT, they could account for a large part of the aberration in calcium metabolism which has been observed in many species of birds since the Second World War.

DECLINING populations of raptorial and fish-eating birds in Great Britain¹ and North America² have produced thin-shelled eggs since the period after the Second World War. A widespread change in the chemical environment which affected the calcium physiology of these species evidently occurred at that time. The chlorinated hydrocarbons, which came into general use in the 1940s, may now be the most abundant synthetic pollutants present in the global environment³. Thin eggshells have been found only in species which accumulate high concentrations of these compounds: relatively uncontaminated populations of these species continue to produce normal eggs^{4,5}.

Calcium metabolism in birds is intimately related to reproductive metabolism and is to a large extent regulated by steroids such as oestrogen and vitamin D⁶. The deposition of medullary bone, the chief source of calcium during egg and eggshell formation, is controlled by the steroid sex hormones⁷, and hens deficient in vitamin D lay eggs with lower eggshell weights⁸. Steroids are hydroxylated and thereby degraded *in vivo* and *in vitro* by hepatic enzymes induced by exogenous, lipid-soluble substances, including the chlorinated hydrocarbons⁹⁻¹⁴. The relatively small amounts of chlorinated hydrocarbons required to produce this effect^{9,11,12,15}, and the discovery that small amounts of some of the DDT compounds are oestrogenic^{16,17}, have made irrelevant much of the parts per million approach to pollutant ecology based on toxicity data alone.

In both Great Britain¹¹⁻¹³ and North America¹⁸ it was the decline of the peregrine falcon which initiated concern about the extent of the harmful effects of environmental contamination. In the United States the eastern population was extinct before competent observers were aware of a general, widespread decline¹⁹. Breeding peregrines persist in apparently normal numbers in British Columbia²⁰ and in the Arctic^{21,22}.

In 1967 we collected an unhatched, abandoned egg of a peregrine falcon in south-western North America, where a small remnant population remains (unpublished work of M. N. K., R. W. R., and S. G. H.). Analysis of this egg (Table 1) showed that it contained almost 5 mg of *p,p'*-DDE (dichloro-2,2-bis(p-chlorophenyl)-ethylene). Unknown peaks present in the chromatograms of the extract of the egg were unidentified until polychlorinated biphenyls (PCB) were detected in European wildlife²³⁻²⁵. Positive confirmation of the identification was accomplished by mass spectrometry in Sweden²⁶. The retention times of the unknown peaks in the peregrine extracts proved to be identical with those of several PCB compounds on DC-200 and QF-1 columns² and on a mixed SE-30:QF-1 column. Other species of birds and fish were subsequently analysed for PCB. The chlorine content of several extracts and standards was determined with a Dohman microcoulometric detector and a method of quantification of the PCB compounds was devised based on peak heights produced in the electron capture detector related to standard *p,p'*-DDE². The DDT compounds are destroyed by nitration²⁷ and *p,p'*-DDT (1,1,1-trichloro-

2,2-bis(p-chlorophenyl)ethane, DDT) (1,1-dichloro-2,2-bis(p-chlorophenyl)ethane) and toxaphene are dehydrochlorinated by saponification with alcoholic KOH. PCB is not degraded by either procedure.

Table 1. CHLORINATED HYDROCARBONS IN NORTH AMERICAN PEREGRINE FALCONS

Sample	Dieldrin	Total DDT*	Percentage DDT	PCB	DDT/PCB
1. Unhatched egg† Baja California (wet)	0-11	102	97	10-2	10
2. Second year migrant from Arctic					
Breat muscle (wet)	NM	90	94	24	3-3
Breat muscle (dry)	NM	206	84	31	4-7
Brain (wet)	NM	85	94	31	3-3
Carcase (wet)	0-17	70	93	10-7	3-3
Carcase (lipid)	62-5	5,000		1,420	
3. Immature, California‡					
Breat muscle (wet)	NM	14-4	90	0-4	1-5
Liver (wet)	NM	7-7	92	4-5	1-7
Brain (wet)	0-01	2-8	90	1-5	1-9
Brain (lipid)	0-50	35		10-3	
Carcase (wet)	0-11	20-2	92	10-8	1-9
Carcase (lipid)	1-0	300		100	
4. Adult, California§					
Breat muscle (wet)	NM	127	87	98	1-3
Liver (wet)	NM	77	80	81	1-4
Brain (wet)	0-31	40-5	88	31-4	1-4
Brain (lipid)	8-7	595		415	
Carcase (wet)	1-7	85	87	85	1-3
Carcase (lipid)	60	2,600		1,960	
5. Immature migrant from Arctic					
Breat muscle (wet)	NM	2-3	81	0-16	14
Breat muscle (dry)	NM	7-8	92	0-54	10
Liver (wet)	NM	1-0	83	0-10	12
Brain (wet)	NM	0-43	83	0-037	10
Body fat (wet)	NM	50-3	82	8-2	18
Carcase (wet)	0-07	9-2	83	0-80	12
Carcase (lipid)	0-44	63-7		5-5	
6. Immature Arctic migrant**					
Breat muscle (wet)	NM	1-0	89	0-6	3-4
Breat muscle (dry)	NM	6-0		1-9	

Concentrations in parts per million wet weight, dry weight or lipid weight.

* DDT residues include: *p,p'*-DDT, *p,p'*-DDE, *p,p'*-DDD (*p,p'*-TDE), *p,p'*-DDMU, *p,p'*-DDT and *p,p'*-DDE. NM, Not measured.

† Chlorinated hydrocarbon contents of the egg were: 4,700 µg *p,p'*-DDE; 40 µg *p,p'*-DDE; 20 µg *p,p'*-DDT; 7-4 µg *p,p'*-DDD; 37 µg *p,p'*-DDMU; 5 µg dieldrin; 12 µg heptachlor epoxide. Concentrations were calculated by assuming a volume of 47-8 ml, the average value obtained by measuring eleven clutches of peregrine eggs from California in the Museum of Vertebrate Zoology, University of California, Berkeley, and by assuming a density of 1-0.

‡ Second year female, captured in October on the Texas coast during migration. Died in captivity shortly afterwards, no apparent cause of death. No body fat. Total body content of chlorinated hydrocarbons: 35 mg DDT, 10 mg PCB, 0-44 mg dieldrin, 0-18 mg heptachlor epoxide. Total body lipid, 7 g.

§ Plummage characteristics of both 3 and 4 were intermediate between those of typically resident California birds and the Arctic race *F. p. tundrus*, as described by White²⁸. No. 3 was an immature female of the year trapped in the southern San Francisco Bay area in the winter of 1966. Observed feeding in the area for a week before capture. Died suddenly after eating a dead gull. Moderate body fat. Total body chlorinated hydrocarbon content: 18 mg DDT, 6-8 mg PCB, 0-07 mg dieldrin, 0-09 mg heptachlor epoxide. Total body lipid, 43 g.

|| Adult female. Trapped in the southern San Francisco Bay area in the winter of 1966. Observed feeding in the area for a month before capture. Died shortly afterwards with no apparent cause of death and no body fat. Total body chlorinated hydrocarbon content: 62 mg DDT, 40 mg PCB, 1-0 mg dieldrin, 1-0 mg heptachlor epoxide. Total body lipid, 20 g.

** First year female trapped on the Texas coast in October. Lost at Pt Mugu, California, the following January and was shot by a sportsman 3 weeks later. Abundant body fat. Total body content: 7 mg DDT, 0-6 mg PCB, 0-06 mg dieldrin, 0-17 mg heptachlor epoxide. Total body lipid, 110 g.

** First year female trapped on the Texas coast in October. Died shortly afterwards of heat prostration.

In Table 1 are presented the results of analyses of peregrine falcons which died from a variety of causes

shortly
amount
a few
were p
except
trapped
lipid c
content
toxic b
times o
signifi
noted b
liver m
were b
but D
parable

Table
prey sp
Baja C
remains
and of
found at
eared g
(Loomis
respecti
Black

(Table 1
both D
populati
1968 no
a single
number
were r
municat
eggshell
cyste wh
of 34 p
mm (0.5
collected
with litt
sources o

In Cal
been red
The rema
normally
four you
surviving
uncontan

Species

Eared grebe (

Whole body

Black petrel (

Whole body

Least petrel (C

Whole body

Eggs

Fish bat (3)

Whole bod

Cravert's munt

Eggs (one c

Whole bod

Whole bod

Elegant tern (

Eggs

Heermann's g

Eggs

Content 1

* All sp

each prey sp

one northern

most petrels,

petrels, Haliae

Larus argentat

* Not me

† Both eg

DSW 029241

year of
04/1/00

In California, numbers of breeding peregrines have been reduced by at least 80 per cent in recent years. The remaining few pairs, however, seem to be reproducing normally and rear, when undisturbed, between two and four young each year. They are found, like most of the surviving pairs in Great Britain¹¹, in a relatively uncontaminated region and seem to be feeding on birds

In sea birds from the Pacific the ratio was usually between 5 and 10 (Table 3). It was not found in eggs of the *Adelie penguin*, *Leucis caeva* (Linn.), *Antarctica*.

1. **Hold Page**

Content in μg of whole bodies and eggs; concentrations in p.p.m. wet weight (W) or lipid weight (L).
 * All specimens were collected in the vicinity of four peregrine eyries in Baja California. Numbers in parentheses are the number of remains of each prey species which were found at the eyries. Also found were remains of one cormorant (*Phalacrocorax* sp.), one red phalarope (*Phalaropus fulicarius*), one northern phalarope (*Lobipes lobatus*) and one mourning dove (*Zenaidura macroura*), but local specimens of these species were not analysed. Eggs of frigate, petrel, elegant tern and Boermann's gulls were from different clutches. Bared grebe: *Podiceps caspius*; black petrel: *Loomisania melanis*; least tern: *Aleutophila melanotos*; Ash-bird: *Fusonyx vivax*; Craver's murrelet: *Endomyschura craveri*; elegant tern: *Thalasseus elegans*; Boermann's gull: *Larus boerhaavi*.
 † Not measured.
 ‡ Both eggs also contained 0.06 p.p.m. dieldrin (lipid weight) and 0.17 p.p.m. endrin (lipid weight).

† Not measured. Interfering peaks on chromatograms.
‡ Both eggs also contained 0.06 p.p.m. dieldrin (lipid weight) and 0.17 p.p.m. endrin (lipid weight).

shortly after being trapped for falconry. Significant amounts of PCB were present in Arctic peregrines only a few months old (Nos. 5 and 6), but higher residues were present in a second year Arctic bird (No. 2) and exceptionally high residues were present in an adult trapped in California (No. 4). In birds 2 and 4 the total lipid reserves were very low, and in both the brain concentrations of DDE and PCB were high, perhaps at toxic levels. Fat mobilization during reproduction or in times of starvation or stress could be expected to cause significant changes in the internal distribution of chlorinated hydrocarbons. Steroid hydroxylase activity in the liver might increase at this time. Dieldrin concentrations were lower than in the peregrines analysed in Britain, but DDE concentrations were approximately comparable¹¹.

Table 2 presents the results of analyses of peregrine prey species, remains of which were collected at eyries in Baja California, Mexico. With the exception of the remains of one mourning dove (*Zenaidura macroura*) and of three fish bats (*Pisonyx vivax*), prey material found at four eyries consisted of sea birds. Of these, the eared grebe (*Podiceps caspicus*) and the black petrel (*Loonchianus melanotos*) constituted 32 and 25 per cent, respectively, of the remains.

Black petrels, like other petrels and shearwaters (Table 3), contained especially high concentrations of both DDT and PCB. Reproductive success in this population of peregrines seems to be subnormal. In 1968 no pairs were observed to hatch or fledge more than a single young. In the past, two to four young, the number normally produced by healthy peregrines^{12,13}, were raised by each pair (L. W. Walker, personal communication). The thickness of fragments of a peregrine eggshell, with its membrane, collected in 1968 below an eyrie where one young hatched, was 0.24 mm, a decrease of 34 per cent from the mean thickness of 0.34 mm $\pm$ 0.015 mm (95 per cent confidence level) in twenty-three eggs collected in the area before 1947. The region is wilderness, with little or no human interference, and is remote from sources of pollution.

In California, numbers of breeding peregrines have been reduced by at least 80 per cent in recent years. The remaining few pairs, however, seem to be reproducing normally and well, when undisturbed, between two and four young each year. They are found, like most of the surviving pairs in Great Britain^{12,13}, in a relatively uncontaminated region and seem to be feeding on birds

which contain low concentrations of chlorinated hydrocarbons. A pair which fledged three young in 1968 fed chiefly on shearwaters and cormorants during the breeding season. Both prey groups are relatively uncontaminated (M. N. K., R. W. R. and S. G. H., in preparation). A clutch of eggs collected in this region weighed as much as the eggs obtained in pre-war years, whereas other peregrine eggs from California collected since the Second World War have been thin-shelled (ref. 2, and D. W. Anderson, D. Hickey and R. F. Christensen, in preparation). Despite official protection the surviving birds are still subjected to shooting by sportsmen and to harassment at the eyries. If these could be effectively reduced the population might yet survive.

In Table 3 are presented the results of analyses of marine and terrestrial birds and of three species of freshwater fishes for PCB and DDT content and concentration. From the ratio of total DDT to PCB, it is apparent that regional fallout patterns exist. In most of the birds from San Francisco Bay which have been analysed, including peregrine falcons (Table 1) and eggs of the western gull, the Caspian tern and the black-crowned night heron (Table 3), the ratio was between one and two. Another black-crowned night heron egg had a typically "ocean" profile, suggesting that the adult female had wintered along the coast. In most of the birds from the Farallon Islands, which are 27 miles west of the Golden Gate Bridge, this ratio was between 2 and 5. In the Gulf of Panama, where PCB contamination might come from the Canal Zone and industrial areas in Panama City, the ratio is between 1 and 2 (Table 3).

In the Gulf of California, a region relatively remote from the sources of either DDT or PCB contamination, the ratio was in most cases approximately 9 or 10. This was true in the egg of the peregrine falcon (Table 1), in all of the black petrels and least petrels analysed, in the Craver's murrelets (Table 2), in five of six osprey eggs and in six of seven western gull eggs (Table 3). Among the exceptions the fish bat, with low concentrations of both DDT and PCB, was the most divergent, with a ratio of 43 (Table 2). It is not clear whether this reflects differences between avian and mammalian physiology or a fundamental difference in feeding habits. The other exceptions include species which are present in the area only during the breeding season (Table 2).

In sea birds from the Pacific the ratio was usually between 5 and 10 (Table 3). PCB was not found in eggs of the Adeline penguin from Cape Crozier, Antarctica.

Table 2. PCB AND DDT IN PREY SPECIES OF PEREGRINE FALCONS IN THE GULF OF CALIFORNIA

Species*	N	DDT (#g)	DDT (p.p.m.)	Percentage DDE	PCB (#g)	PCB (p.p.m.)	DDT/PCB
Eared grebe (18)	8	—	0.25, 0.26, 12.1	97	NM†	—	—
Whole body							
Black petrel (10)	8	810.1 (685-1,344)	9.2 (W)	81	—	1.0 (W) (0.90-1.14)	9.2
Whole body							
Least petrel (6)	8	99 80 (23-37)	8.2 (W)	83 84	— 8.1 (1.9-5.0)	0.35 (W)	9.3 10
Whole body							
Fish bat (3)	7	25 (16-31)	0.71 (W)	62	0.58 (0.45-1.06)	0.02 (W)	43
Whole body							
Craver's murrelet (2)	2	230.1 (223-238)	89 (L)	80	—	4.5 (L)	8.7
Eggs (one clutch)							
Whole body, adult	1	37.1	0.31 (W)	75	—	0.030 (W)	7.0
Whole body, adult	1	295	2.4 (W)	85	—	0.26 (W)	9.2
Elegant tern (1)	1	15.5 (9.6-24.3)	5.0 (L)	90	—	1.5 (L) (0.8-2.6)	3.0
Eggs							
Heermann's gull (1)	3	195 (94-278)	48 (L)	95	—	8.1 (L) (3.5-11.3)	5.9
Eggs							

* Content in #g of whole bodies and eggs; concentrations in p.p.m. wet weight (W) or lipid weight (L).

† All specimens were collected in the vicinity of four peregrine eyries in Baja California. Numbers in parentheses are the number of remains of each prey species which were found at the eyries. Also found were remains of one cormorant (*Phalacrocorax* sp.), one red phalarope (*Phalaropus fulicarius*), one northern phalarope (*Lophopus lobatus*) and one mourning dove (*Zenaidura macroura*), but local specimens of these species were not analysed. Eggs of least petrel, elegant tern and Heermann's gull were from different clutches. Eared grebe: *Podiceps caspicus*; black petrel: *Loonchianus melanotos*; least petrel: *Holocystops microps*; fish bat: *Pisonyx vivax*; Craver's murrelet: *Endomychus craveri*; elegant tern: *Thalasseus elegans*; Heermann's gull: *Larus heermanni*.

‡ Not measured. Interfering peaks on chromatograms.

§ Both eggs also contained 0.08 p.p.m. dieldrin (lipid weight) and 0.17 p.p.m. endrin (lipid weight).

DSW 029243

Table 2. PCB AND DDT IN THE GLOBAL ECOSYSTEM

Species	N	Total DDT (μ g)	Percentage DDT (p.p.m.)	PCB (μ g)	DDT PCB (p.p.m.)
White crappie (1)	1	—	1-83 (W)	—	0-004 (W)
Black crappie (2)	1	—	2-10 (W)	—	0-001
Bluegill (3)	1	—	5-5 (W)	—	0-005
Adelie penguin (4)	5	0-74 (0-59-1-04)	0-128 (L)	74	< 0-041 μ g/egg
Western grebe (5)	1	—	26-4 (W)	—	0-006 (W)
Breast muscle	1	—	—	—	270
Fulmar (6)	1	—	0-41 (W)	76	0-004 (W)
A	1	—	3-4 (W)	89	0-34 (W)
B	1	10-475	17-5 (W)	86	4-5 (W)
C	1	2,000	3-0 (W)	93	0-42 (W)
Pink-footed shearwater (7)	1	—	—	—	277
Sooty shearwater (8)	1	—	12-3 (W)	94	1-2 (W)
A	1	—	10-3 (W)	86	0-9 (W)
B	1	1,265	2-3 (W)	85	NM
C	1	—	32-0 (W)	92	3-1 (W)
Slender-billed shearwater (9)	1	—	—	—	—
Ashy petrel (10)	8	2,158 (1,041-3,488)	54-3 (W)	95	840 (594-1192)
Brown pelican (11)	4	89 (18-183)	11-5 (L)	81 (47-78)	23-1 (18-30)
Eggs, Panama	2	53 (47-60)	10-0 (L)	81	10-4 (8-0-11-7)
Eggs, Baja California	8	9-8, 8-7, 30-0	—	84	8-4, 6-7, 84-0
Frigate-bird (12) eggs, Panama	4	20-8 (16-4-24-5)	8-2 (L)	80	12-2 (6-5-19-9)
Brown booby (13) eggs, Panama	4	825	—	91	113
Braill's cormorant (14) eggs	17	124 (125-130)	—	90	82 (44-75)
Pink cormorant (15) eggs	2	4,340	—	70	—
Common teal (16)	1	—	10-9 (W)	—	0-91
White-tailed kila (17)	4	11-1 (9-8-12-0)	0-76 (W)	77	6-4 (5-5-7-8)
Eggs, clutch A	1	5-1	0-34 (W)	82	0-81
Unhatched egg B	1	6-3	0-35 (W)	73	3-0
Unhatched egg C	1	8-3/egg	—	82	4-3/egg
Clutch D	2	10-2/egg	9-0 (L)	87	4-0/egg
Clutch E	3	21-0/egg	—	80	3-8/egg
Clutch F	4	8,500	25-2 (W)	90	6-3
Cooper's hawk (18)	1	160	2-0 (W)	94	17-5
Golden eagle (19) egg	1	127	55 (L)	85	17-5 (7-3, 3-0, 103)
Osprey (20), Baja California	6	80-264	—	85-4	24, 7-0, 19
Merlin (21)	1	435	2-0 (W)	94	0-39 (W)
American kestrel (22)	1	5-1	0-044	90	2-7
Whole body, adult	1	2-4/egg	0-30 (W)	93	1-0/egg
Eggs, two clutches	6	—	—	—	0-031 0-06 (W)
Black-crowned night heron (23)	1	541	—	89	330
Egg	1	809	—	99	23
Western gull (24)	4	803 (632-1,125)	—	85	805 (580-1,310)
Eggs, San Francisco Bay, three clutches	10	412 $\pm$ 102	—	89	188 $\pm$ 55
Eggs, Farallon Is., ten clutches	7	885 $\pm$ 290	—	97	45 $\pm$ 80
Eggs, Baja Calif., seven clutches	2	665 (594-732)	—	89	114 (91-137)
Forster's tern (25) eggs	2	1,209 (1,216-1,222)	—	89	805 (660-950)
Caspian tern (26)	8	1,430 (991-2,480)	—	88	1,010 (550-1,600)
Eggs, San Diego Bay	1	—	0-78 (W)	70	—
Red phalarope (27)	6	1,945 (932-3,021)	161 (L)	90	558 (364-1,010)
Common nurre (28) eggs	1	—	5-8 (W)	98	—
Cassin's auklet (29)	1	—	0-75	90	0-16
Ancient murrelet (30)	1	—	2-7	97	0-15
Black-throated auklet (31)	1	—	0-19 (W)	77, 93	0-26
Mourning dove (32)	2	12-0, 33-1	—	—	—
Barn owl (33)	8	27-7/egg	1-25 (W)	95	10-4/egg
Eggs, one clutch	2	143/egg	6-6 (W)	96	14-4/egg
Eggs, one clutch	2	21-2, 448	0-18, 3-3 (W)	77, 93	2-5, 28-1
Meadowlark (34)	2	—	—	—	—

Content in μ g and concentrations in p.p.m. wet weight (W) or lipid weight (L).

Unless otherwise specified, analyses were of whole bodies. Endrin and dieldrin were identified on the basis of retention times on both QF-1 and DC-200 columns.

1. *Pomoxis annularis*, 244 g. Clear Lake, Lake Co., Calif., May 1968.
2. *Pomoxis nigromaculatus*, 212 g. Clear Lake, Lake Co., Calif., May 1968.
3. *Lepomis macrochirus*, 229 g. Clear Lake, Lake Co., Calif., May 1968.
4. *Thynnus aethiops*, Cape Crozier, Antarctica, October 1967.
5. *Archimophorus occidentalis*, Clear Lake, Lake Co., Calif., May 1968.
6. *Fulmarus glacialis*, A and B: Monterey Bay, Calif., November 1, 1966. C: Point Reyes, Calif., December 1967. Fulmars breed in Alaska.
7. *Fulmarus glacialis*, Breeds in Chile. Collected in May 1968, in the Gulf of California.
8. *Fulmarus glacialis*, Breeds in New Zealand and Chile. A and B: Monterey Bay, November 1, 1966. C: Gulf of California, May 1968. NM, Not measured, interfering peaks.
9. *Fulmarus glacialis*, Breeds in Australia. Monterey Bay, December 1966.
10. *Oceanodroma homochroa*, Farallon Islands, Calif., May 1968.
11. *Pelecanus occidentalis*, Panama eggs were collected on Isla Pacheca and Isla Pachequilla, Gulf of Panama, February 1968. One egg contained 0-06 p.p.m. dieldrin and 0-86 p.p.m. endrin, another contained 0-16 p.p.m. dieldrin and 0-07 p.p.m. endrin (lipid weight). Baja California eggs were collected at Bahia de los Angeles, March 1968. One egg contained 0-24 p.p.m. dieldrin and 1-18 p.p.m. endrin (lipid weight).
12. *Fregata magnificens*, Isla Pacheca and Isla Pachequilla, Panama, February 1968.
13. *Sula leucogaster*, Isla Pacheca and Isla Pachequilla, Panama, February 1968. Two eggs were analysed for dieldrin and endrin. Dieldrin: 0-06, 0-18 p.p.m.; endrin: 0-06 and 0-01 p.p.m. (lipid weight).
14. *Phalacrocorax auritus*, Farallon Islands, May 1967.
15. *Phalacrocorax auritus*, San Mateo Co., Calif.
16. *Anas platyrhynchos*, San Diego, April 1968. Adult male.
17. *Blennius leucurus*, A: Contra Costa Co., Calif., April 1968. B and C: Contra Costa Co., May 1968, two and three young raised, respectively. D: Abandoned, Contra Costa Co., May 1967. E and F: Destroyed nests, Contra Costa Co., March 1968.

DSW 029244

Table 3 (continued)

18. <i>Accipiter cooperii</i> . Ballboa Park, San Diego, February 1968. First year female, died of trichomoniasis.	
19. <i>Aquila chrysaetos</i> . San Luis Obispo Co., Calif., April 1968. Unhatched egg in nest where one young was raised. Egg also contained 4.7 µg of dieldrin, 1.0 µg of heptachlor epoxide, but no endrin.	
20. <i>Pandion haliaetus</i> . Gulf of California, March 1968. One egg also contained 0.10 p.p.m. dieldrin and 0.26 p.p.m. endrin (li).	
21. <i>Falco columbarius</i> . Immature, Utah, December 1967.	
22. <i>Falco sparverius</i> . Adult, was killed on road, Mendocino Co., Calif., December 1967. Eggs from Davis, California, 1966.	
23. <i>Nucifraga velox</i> . Eggs from different clutches, San Francisco Bay, May 1967.	
24. <i>Larus occidentalis</i> . Standard errors, 95 per cent confidence limits.	
25. <i>Sterna forsteri</i> . San Diego Bay, May 1967.	
26. <i>Hudobrookea caspia</i> , 1967.	
27. <i>Phalacrocorax auritus</i> . Monterey Bay, November 1, 1966.	
28. <i>Uria lomvia</i> . Farallon Islands, May 1967.	
29. <i>Myiophobus alpestris</i> . Farallon Islands, April 1966.	
30. <i>Synthlibramphus antiquus</i> . Monterey Bay, November 1, 1966.	
31. <i>Certhia minckleyi</i> . Monterey Bay, November 1, 1966.	
32. <i>Geothlypis trichas</i> . San Diego, July 1968.	
33. <i>Tyrus alba</i> . Clutch A from Contra Costa Co., Calif., March 1968. Clutch B from Yolo Co., Calif., April 1968.	
34. <i>Sturnella neglecta</i> . Davis, Calif., December 1967.	

In these, however, the amount of DDT was very low; with a DDT:PCB ratio of 18, no PCB would have been detected. A larger amount of fat material from Antarctic organisms, which would contain more DDT, would therefore have to be analysed before concluding that PCB has not yet reached the Antarctic.

Individuals of species resident in industrial areas have, as expected, higher PCB levels than individuals of the same species from more remote regions. Analysis of ten eggs from ten clutches of the western gull from the Farallon Islands showed an average PCB content of 130 ± 55 µg (95 per cent confidence level). PCB content of four eggs from three clutches from an island in San Francisco Bay averaged 805 µg with a range from 580 to 1,310 (Table 3). Seven eggs from seven clutches in the Gulf of California contained 45 ± 30 µg of PCB. The DDT content of the eggs from the Farallons was not significantly different from that of the Baja California eggs.

Birds from the Gulf of California also contain dieldrin and endrin (Tables 2 and 3), but the number of analyses is as yet insufficient to compare their relative abundance with that of DDT and PCB. Despite the inability of the Shell Chemical Company to find any chlorinated hydrocarbons in the Gulf of California and at the mouth of the Colorado River¹², which drains into the Gulf, it is likely that some do enter from the Colorado River and from agricultural areas in western Mexico. A significant fraction, however, must come from the atmosphere¹³, and air transport best explains the presence of PCB in remote areas. In extracts from the Gulf of California one PCB compound was present in relatively small concentrations. This compound is readily degraded by ultraviolet light irradiation in laboratory experiments, and may therefore be selectively degraded in the atmosphere. Although PCB is not soluble in water, it has a low but finite vapour pressure¹⁴. Incineration of materials containing PCB would greatly increase the rate of entry into the atmosphere. PCB is used in the manufacture of many industrial products, so the high amounts found in San Francisco Bay, Puget Sound¹⁵ and San Diego Bay presumably result from direct discharge of industrial wastes into these waters and from local fallout.

The presence of PCB in the sea land birds and fresh-water fish analysed indicates that it is also distributed among continental ecosystems in North America. Peregrines could therefore acquire PCB, as well as the other chlorinated hydrocarbons, over all their global range. No PCB residue data are available for prey species from the Atlantic, but compounds which seem to be PCB have been isolated from seals¹⁶. A second year male peregrine spent the winter of 1967-68 on Isla Pacheca, Panama, where the pelican, booby and frigate-bird eggs of Table 3 were collected. The Cooper's hawk which was analysed, a species which also preys upon birds, contained high residues of both DDT and PCB (Table 3). Cooper's hawks have declined in eastern North America and in some regions have produced thin-shelled eggs¹⁷. Several

species of raptors do not accumulate high amounts of the organo-chlorine compounds (Table 3), a result of the very low residue levels usually found in their prey. American kestrels and barn owls are common residents of California cities. The white-tailed kite, which was near extinction 40 years ago in California, is now abundant in areas where insecticide use is intense, yet because it preys primarily upon the short-lived and herbivorous vole, *Microtus*, the species accumulates very little DDT or PCB (Table 3).

Previous work¹⁸ has shown that both DDT and dieldrin induce hepatic enzymes in the pigeon which, in an *in vitro* preparation, increase the metabolism of progesterone and testosterone. This work has now been extended to study the metabolism of oestradiol by enzymes induced by *p,p'*-DDE, technical DDT (Dupont) and PCB (Aroclor 1262). The experimental procedure previously described¹⁸ was followed except that the chlorinated hydrocarbons were injected intramuscularly rather than given orally, and in the separation of oestradiol and its metabolites the solvent system used was the upper layer of a mixture of benzene:heptane:methanol:water 7:3:8:2. The strips were monitored with an autographic strip scanner. The profiles obtained for the various inducing agents are shown in Fig. 1. It will be noted that the profiles of the metabolites obtained after enzyme induction using DDE and DDT are identical, but that a different metabolite is produced by the enzyme induced by PCB. The amount of metabolites formed was calculated from the radioactivity of the peaks. The results obtained are given in Table 4.

Table 4. INCREASE OF OESTRADIOL METABOLISM BY PIGEON LIVER HOMOGENATES FROM BIRDS TREATED WITH VARIOUS CHLORINATED HYDROCARBONS

	Amount of polar metabolites formed in mmoles
Control	29.3 ± 0.5
DDE (40 mg/kg)	76.2 ± 13.1
DDT (40 mg/kg)	93.1 ± 11.2
PCB (20 mg/kg)	180.0 ± 10.5

Each figure is the average with standard deviation of a group of four birds. In all cases 500 mmoles of oestradiol-6,7-T (500 mCi/mmoles) was present in the incubation mixture. Incubation time 30 min, weight of microsomal fraction used 800 mg. All chlorinated hydrocarbons were injected intramuscularly into the pectoral muscle 7 days before death.

Body concentrations of 40 p.p.m. of *p,p'*-DDE significantly increased the rate of oestradiol degradation by the induced enzymes in the experimental conditions. On a weight basis the PCB preparation had an oestradiol degrading potential approximately five times that of *p,p'*-DDE or technical DDT. Both DDE and PCB, which are apparently the most abundant of the chlorinated hydrocarbon pollutants in the global ecosystem, have therefore the capacity to produce sublethal physiological effects in birds.

Studies on the activity of induced enzymes in the rat at various times after a single injection of DDT or dieldrin have been made by Ghazal *et al.*¹⁹. He found that it took 70 days for the activity to fall to half its maximum value in the case of DDT and 15 days in the case of dieldrin. Experiments with dieldrin on pigeons

DDT:PCB

47.8

600

1,200

> 10

270

5

10

2.7

2.2

10

12

13

8.5

1.35

(0.97-3.2)

0.8, 4.0

1.1, 1.5, 0.4

1.7

2.9

2.1

12

1.8

4.0

1.5

2.0

2.4

4.3

4.0

4.5

5.5, 10.1, 1.3

11.2, 9.7, 11.1

12.7

1.4

2.8

1.6

80

1.0

8.0

0.1, 10.3, 8.0

18.3, 10.6

10.7, 10.5

0.8

1.7

1.4

6

5.5

34

3

K

3.7

10

0.4, 15.9

1 add DC-800

Alaska.

B. NM, Not

eg contained

allifornia eggs

aldrin: 0.06,

otively. D:




Fig. 1. Chromatographic separation of oestradiol and its metabolites. The larger peak is unaltered oestradiol.

show that only a quarter of maximum activity remains after a month. Thus there is evidence that the effects of these induced enzymes can persist over a long period of time although more studies are needed to determine the steroid degrading potentials of livers of those species contaminated with chlorinated hydrocarbons. The profile of the metabolites should be some indication of the history of exposure to chlorinated hydrocarbons.

The reductions in eggshell thickness and eggshell weight increase the chance of egg breakage¹¹⁻¹⁴, and water retention, which affects hatching success¹⁵⁻¹⁷, might be impaired. The environment in which birds now exist is therefore no longer the same as that in which they evolved; it is unlikely that any species has the genetic capacity to meet the selection pressures resulting from the abrupt environmental change which has produced the thin eggshells. The peregrine falcon is a species long highly revered and respected. G. H. Thayer¹⁸ has described it as "the embodiment of noble rapacity and lonely freedom". An irony therefore exists in the fact that the peregrine may be the first species to be extirpated by global contamination.

Tsunamis on the Moon?

by

W. G. VAN DORN

Scripps Institution of Oceanography,
University of California,
San Diego, California

The question whether the multiple ring-like mountain structures surrounding most of the lunar craters larger than 120 miles in diameter might be "frozen tsunamis", set in motion by the shock waves caused by impacting

The spacing of the five annular mountain rings around Mare Orientale fits a dispersion curve for gravity waves on a 50 km "liquid" layer overlying a rigid basement.

This work was supported by the US National Science Foundation, the Bureau of Sport Fisheries and Wildlife and the US Public Health Service. Pure *p,p'*-DDE was supplied by the Federal Drug Administration, and samples of 'Aroclor' by the Monsanto Chemical Company.

Received August 26; revised November 14, 1968.

- ¹ Ratcliffe, D. A., *Nature*, **215**, 208 (1967).
- ² Hickey, J. J., and Anderson, D. W., *Science*, **182**, 271 (1968).
- ³ Riechbrough, R. W., in *Chemical Fallout, Fetal Rochester Conference on Toxicity* (in the press, 1968).
- ⁴ Slinkin, K., *Calcium in Reproductive Physiology* (Rothschild, New York, 1967).
- ⁵ Slinkin, K., *Biol. Rev.*, **34**, 321 (1961).
- ⁶ Clavert, J., and Benoit, J., *Chim. Soc. Biol., Paris*, **134**, 500 (1962).
- ⁷ Kova, P., and Potter, T. R., *Anal. Rec.*, **69**, 377 (1964).
- ⁸ Hart, E. B., Steenbock, H., Lepkovsky, S., Kletzian, R. W. F., Halpin, J. G., and Johnson, O. N., *J. Biol. Chem.*, **64**, 679 (1955).
- ⁹ Welch, R. M., Levin, W., and Conney, A. H., *Chemical Fallout, Fetal Rochester Conference on Toxicity* (in the press).
- ¹⁰ Hart, L. G., Schuller, R. W., and Fouts, J. R., *Toxicol. Appl. Pharmacol.*, **8**, 371 (1965).
- ¹¹ Hart, L. G., and Fouts, J. R., *Neuropharmacology's Arch. Exp. Path. Pharmacol.*, **248**, 486 (1965).
- ¹² Conney, A. H., Welch, R. M., Kuntzman, R., and Burns, J. J., *Clin. Pharmacol. Therap.*, **8**, 2 (1966).
- ¹³ Conney, A. H., *Pharmacol. Rev.*, **19**, 317 (1967).
- ¹⁴ Kupfer, D., *Neuropharmacol.*, **19**, 11 (1967).
- ¹⁵ Gerbath, G., and Schwabe, U., *Arch. Exp. Pathol. Pharmacol.*, **244**, 469 (1964).
- ¹⁶ Ollivet, J. W., *J. Agric. Food Chem.*, **16**, 295 (1968).
- ¹⁷ Schwabe, U., *Arzneimittel-Forsch.*, **14**, 1265 (1964).
- ¹⁸ Peakall, D. B., *Nature*, **216**, 505 (1967).
- ¹⁹ Welch, R. M., Levin, W., and Conney, A. H., *Toxicol. Appl. Pharmacol.* (in the press).
- ²⁰ Bliman, J., Cecil, H. C., Harris, S. J., and Yike, G. F., *Science*, **188**, 371 (1968).
- ²¹ Ratcliffe, D. A., *Bird. Birds*, **31**, 23 (1958).
- ²² Ratcliffe, D. A., *Bird Study*, **10**, 56 (1963).
- ²³ Ratcliffe, D. A., *Bird Study*, **12**, 66 (1965).
- ²⁴ Ratcliffe, D. A., *Bird Study*, **14**, 238 (1967).
- ²⁵ Hickey, J. J., *Peregrine Falcon Populations: Their Biology and Decline* (University of Wisconsin Press, in the press).
- ²⁶ Boebe, F. L., *Condor*, **62**, 145 (1960).
- ²⁷ Cade, T. J., White, C. M., and Haugh, J. R., *Condor*, **70**, 170 (1968).
- ²⁸ Enderson, J. H., and Berger, J. D., *Condor*, **70**, 149 (1968).
- ²⁹ Jensen, B., *New Sci.*, **22**, 1612 (1964).
- ³⁰ Whitmark, G., *J. Assoc. Off. Anal. Chem.*, **50**, 1009 (1967).
- ³¹ Holmer, D. C., Shimmons, J. H., and Talton, J. O., *Nature*, **216**, 227 (1967).
- ³² Erro, F., Bevenue, A., and Deckman, H., *Bull. Environ. Cont. Toxicol.*, **372** (1967).
- ³³ White, C. M., *Auk*, **85**, 179 (1968).
- ³⁴ Jefferies, D. J., and Prest, I., *Bird. Birds*, **39**, 49 (1966).
- ³⁵ Aldrin, Dieldrin, Endrin (A Status Report) (The Shell Chemical Company, 1967).
- ³⁶ Riechbrough, R. W., Huggot, R. J., Griffin, J. J., and Goldberg, E. J., *Science*, **159**, 1233 (1968).
- ³⁷ Tech. Bull. PL-306 (Monsanto Chemical Company).
- ³⁸ Holden, A. V., and Marsden, K., *Nature*, **216**, 1274 (1967).
- ³⁹ Ghazal, A., Korasaky, W., Porus, J., Vohlend, H. W., and Klempeu, I., *Arch. Exp. Pathol. Pharmacol.*, **249**, 1 (1964).
- ⁴⁰ Jones, B. E., and Leopold, A. B., *J. Wildl. Management*, **37**, 221 (1967).
- ⁴¹ Mayhew, W. W., *J. Wildl. Management*, **19**, 34 (1965).
- ⁴² Thayer, G. H., *Bird Lays*, **6**, 47 (1904).

have
of a
large
its
to
rebu
conce
fricti
two
calcu
Mare
are
at
The
Mount
plan
most
bene
elevat
supple
pouch
basin,
the
rebu
explos
region
km in
the m
of the
into a
the on
character
in the
crosses
For sur
rigid ha
related
motions
tanous
ring-spa
wave hy

Table 1.

Ring No
Elevation
(km ±1)
Radius (l
e Obse
Com

* Courtesy

Mountain
The w
inviscid li
sively (see
Van Dorn
polar coord
placement
tanous) d
f by

where $E(k)$
transform
 $2\pi/\lambda$ is th
length, $\omega =$
The group
number by