

Copies to: [REDACTED]

E. I. du Pont de Nemours and Company
Haskell Laboratory for Toxicology and Industrial Medicine

HASKELL LABORATORY REPORT NO. 93-71

MR NO. [REDACTED]

Material Tested: [REDACTED]

Haskell No.: 6869

Material Submitted by: [REDACTED] Organic Chemicals
Department, Jackson Laboratory

Other Codes: [REDACTED]

ACUTE ORAL TEST

Procedure: The test material, as an aqueous suspension, was administered by intragastric intubation to young adult ChR-CD male rats in single doses. Survivors were sacrificed 14 days later.

Results:

Suspension (%)	Dose (mg/kg)	Mortality†	Clinical Signs	ALD
40	17,000‡	D - 2 d.	<u>Lethal Dose:</u> Prostration on day of dosing; incoordination, weakness, hypothermia, pallor, prostration, and weight loss on day after dosing	17,000 mg/kg
40	11,000‡	S - 14 d.		
40	7,500	S - 14 d.		
40	5,000	S - 14 d.		
40	3,400	S - 14 d.	<u>Nonlethal Doses:</u> Pallor and hypothermia on day of dosing and weakness on day after dosing at 7500 mg/kg and above; incoordination on day of dosing at 5000 mg/kg and above; weight loss for 1-3 days	
12	2,250	S - 14 d.		
12	670	S - 14 d.		

† D - () d. = Found dead () days after dosing; S - () d. = Sacrificed () days after dosing
‡ Given in divided doses

Company Sanitized. Does not contain TSCA CBI

Summary: [REDACTED] has low toxicity when administered by intragastric intubation to young adult ChR-CD male rats in single doses; its Approximate Lethal Dose (ALD) is 17,000 mg/kg of body weight.

Report by: Laurel S. Wasileski
Laurel S. Wasileski
Oral Toxicology Section

Approved by: John A. Zapp, Jr.
John A. Zapp, Jr.
Director

NB 942, p. 78
LSW:pgh
Date: April 6, 1971

Company Sanitized. Does not contain TSCA CBI