METHOD FOR ANALYSIS OF BIPHENYL FOR
DIBENZOFURAN AND DIBENZO-P-DIOXIN

PRINCIPLE

Biphenyl is dissolved in a suitable solvent and analyzed for dibenzofuran and dibenzo-p-dioxin by gas chromatography/mass spectrometry.

EXPERIMENTAL

1. Biphenyl dissolved in Nanograde® methylene chloride.

2. Solution analyzed using gas chromatograph/mass spectometer/computer system.

3. Concentrations of dibenzofuran and dibenzo-p-dioxin analyzed using identical conditions.

4. Concentrations of components, if found, are calculated as follows:

$$\text{Concentration (ppm)} = \frac{\text{area sample}}{\text{area standard}} \times \frac{\text{concentration standard (mg/ml)}}{\text{biphenyl concentration (g/ml)}}$$

0256751

PCB-ARCH0747420



0256752

PCB-ARCH0747421



0256753

PCB-ARCH0747422



0256754

PCB-ARCH0747423



0256755

PCB-ARCH0747424



0256756

PCB-ARCH0747425


** SELECTED ION CHROMATOGRAM **
5 MICRO-L 3.1 GM
ISO150 10% S-405
184 0
168 0
7
8
9
10

0256757