

107(n)
FILE COPY

EPIDEMIOLOGICAL STUDY
OF
VINYL CHLORIDE WORKERS

Final Report

May 3, 1974

Submitted to
The Manufacturing Chemists Association
1825 Connecticut Avenue N. W.
Washington, D. C. 20009

by
Tabershaw/Cooper Associates, Inc.
2180 Milvia Street
Berkeley, California 94704

DOW 1051

ACKNOWLEDGMENT

The research which forms the basis for this report was supported by the voluntary contributions of the following named chemical companies engaged in the synthesis and/or the polymerization of vinyl chloride, and administered in their behalf by the Manufacturing Chemists Association

Air Products and Chemicals, Inc.

Allied Chemical Corporation

Borden Chemical Division of Borden, Inc.

Continental Oil Company

Diamond Shamrock Company

The Dow Chemical Company

Ethyl Corporation

Exxon Chemical Company

Firestone Plastics Company

B. F. Goodrich Chemical Company

The Goodyear Tire and Rubber Company

Monsanto Company

Olin Corporation

PPG Industries, Inc.

Shell Chemical Company

Stauffer Chemical Company

Tenneco Chemicals, Inc.

Union Carbide Corporation

UNIROYAL, Inc.

TABLE OF CONTENTS

SECTION	Page
I. Summary	1
II. Introduction	2
III. Selection of Study Plants	3
IV. Data Collection	4
V. The Measurement of Exposure	5
VI. Follow-up of Study Population	6
VII. Description of Study Population	7
VIII. Calculation of Risk of Death	9
IX. Analysis	11
X. Discussion	13
XI. Possible Biases in the Calculation of Risk	15
REFERENCES	17

LIST OF TABLES

1. Follow-up Status of 8384 Vinyl Chloride Workers
2. Distribution of Months in Exposed Employment by Birth Year, for 7128 Vinyl Chloride Workers with Completed Follow-up
3. Distribution of Months in Exposed Employment by Year in Which Exposure Began, for 7128 Vinyl Chloride Workers with Completed Follow-up
4. Distribution by Months of Exposed Employment and Time Weighted Average Exposure Score, for 7128 Vinyl Chloride Workers with Complete Follow-up
5. Person Years of Observation by Year of Birth and Age for Vinyl Chloride Workers
6. Observed Deaths/Expected Deaths and Standardized Mortality Ratios in Vinyl Chloride Workers

DOW 1053

7. Observed Deaths/Expected Deaths and Standardized Mortality Ratios in Vinyl Chloride Workers, by Estimated Level of Exposure
8. Observed Deaths/Expected Deaths and Standardized Mortality Ratios in Vinyl Chloride Workers by Duration of Exposed Employment
9. Observed Deaths/Expected Deaths and Standardized Mortality Ratios in Vinyl Chloride Workers with Exposure Indices below 1.5, by Duration of Exposed Employment
10. Observed Deaths/Expected Deaths and Standardized Mortality Ratios in Vinyl Chloride Workers with Exposure Indices of 1.5 or Greater, by Duration of Exposed Employment
11. Distribution of Deaths by Exposure Category and by Whether Cause of Death Was Determined, and Correction Factors Used for SMR Calculations
12. Results of Follow-up in Several Occupational Groups
13. Angiosarcoma Deaths in VCM Epidemiology Study Population
14. Other Malignancies (190-199, I.C.D.) in VCM Epidemiology Study Population
15. Malignant Neoplasms of Buccal Cavity and Pharynx (140-148, I.C.D.) in VCM Epidemiology Study Population

FIGURE

1. Flow Chart of Follow-up Procedure

ATTACHMENTS

- A. Plant Contact Form
- B. Plant Background Form
- C. Job History Form
- D. Follow-up Letter

EPIDEMIOLOGICAL STUDY OF VINYL CHLORIDE WORKERS

Final Report - May 2, 1974

I. SUMMARY

Eight thousand three hundred and eighty four men who had had at least one year of occupational exposure to vinyl chloride before December 31, 1972, were followed in a historical prospective mortality study. The major findings of the study are:

1. The overall mortality of the study population was approximately 75 percent of what would be expected in a comparable population of U. S. males.
2. No cause of death showed a statistically significant excess over what would be expected in a comparable U. S. male population.
3. No new deaths identified as angiosarcoma of the liver were found.
4. All previously known Angiosarcoma deaths in the study population during the study period were found as part of the routine study procedure.
5. Cancers of the liver (primarily angiosarcoma), respiratory system, brain, and cancers of unknown primary site, as well as lymphosarcoma, occurred more often than expected in those members of the study population with the greatest exposure. Even though the excesses were not statistically significant, the findings warrant further study.
6. Other cancers occurred at lower levels than in the general male population, with the exception of cancers of the buccal cavity and pharynx. There was an excess of these

cancers, which however was inversely related to exposure.

The explanation for this finding is not apparent.

II. INTRODUCTION

The study was begun by Tabershaw/Cooper, Associates, Inc. on June 15, 1973, under contract from the Manufacturing Chemists Association. The original eight month period of the contract was later extended by two months, in consideration of logistical problems arising from the number of plants which participated in the study.

The objectives of the study are:

- (1) To compare the mortality of individuals who have worked in vinyl chloride plants with that of the general population;
- (2) To compare mortality patterns within the population of vinyl chloride workers, based upon estimated occupational exposure; and
- (3) To compare mortality among vinyl chloride workers with the mortality of other groups which TCA, Inc. has studied.

The study took the form of a historical prospective mortality study. The study population consisted of individuals who had worked for at least one year in a job involving exposure to vinyl chloride before December 31, 1972, and included retired and terminated as well as active workers. For each such worker the date of birth and an employment history were obtained, and the vital status of the worker as of December 31, 1972, was ascertained. For those found to have died, those death certificates that were available were obtained and the cause of death determined. The observed mortality was compared with that of the United States male population.

III. SELECTION OF STUDY PLANTS

The Manufacturing Chemists Association identified 43 United States plants belonging to 19 companies, which either produced vinyl chloride or used it in the production of polyvinyl chloride. The name and address of a responsible liaison person at each company and plant was provided to TCA, Inc. (Attachment A).

A brief questionnaire was sent to each plant to obtain an estimate of the exposed population and to find out how far back in time personnel records could be obtained, and to determine whether job histories were available for retired and terminated workers (Attachment B).

All the plants responded. On the basis of the replies, four plants were dropped from the study because they had begun operation after 1968. It was felt that exposures in such plants would be too recent and too brief to provide useful data. Study of one other plant was deferred because it had stopped production of vinyl chloride seven years ago, and appeared to present difficulties in identifying an exposed population.

In addition, three more plants were excluded from analysis because, when data analysis began, it was found that information from these plants was defective in one of several respects (e.g., no exposure data, no job histories available for terminated employees, etc.).

The present report therefore deals with 35 domestic plants involved in either VCM or PVC production, which provided a study population of 8384 individuals.

IV. DATA COLLECTION

In each plant, data were collected for each worker stated by the plant management to have been employed for at least one year in a job involving exposure to vinyl chloride. In some plants, particularly those producing the monomer, this determination could be made on the basis of job titles. Usually, however, exposure was a function of both job title and the location of the job in the plant, so that the assessment of exposure had to be made on a case-by-case basis by plant officials.

Data were collected for as far back in time as complete records were kept. In most cases this covered the entire history of the plant. In others, records were kept for a fixed period such as a decade. In a few, records were kept for different periods, depending on whether the worker had died on the job or had left employment.

In the latter case it was necessary to exercise care in defining the period for which study data were to be collected. For example, if a plant kept records for five years on those who left employment, but kept them indefinitely for those who died on the job, no termination could be included in the study unless it occurred within the last five years. Data from the earlier period would be biased for the purpose of this study, since a worker had to die to be in the records for the earlier period. The policy therefore was to determine how long each kind of record was kept (quit, discharged, retired and alive, retired and deceased, died while employed) and to define the study period as the shortest of these.

One problem which arose in the PVC plants resulted from the fact that a worker could transfer within the plant from exposed to non-exposed jobs. Therefore, even though the number of workers in exposed jobs at a given

time may have been a small proportion of the total work force, it was necessary to search every worker's record to find those who might in the past have had over a year's experience in exposed jobs. Several of the PVC plants undertook to conduct this search themselves.

As a result, plant data were collected in two ways. In some plants, a TCA, Inc., team visited the establishment, copied the records designated by management as representing exposed workers, and obtained sufficiently detailed exposure estimates to produce a job history. In others, the plants were provided by TCA, Inc. with standard forms which they filled out for workers whom they identified as exposed. Three basic forms were used, depending on whether the worker was known to be alive, known to be dead, or was of unknown status. In all three cases a work history with estimated exposure was recorded, the differences being in the ancillary data required. Attachment C shows such a form for workers known to be dead.

Approximately 5000 of the study population came from plants visited by TCA, Inc., while the remainder came from plants which had filled out the forms themselves.

V. THE MEASUREMENT OF EXPOSURE

The job history forms (Attachment C) were originally designed in the expectation that vinyl chloride exposure on each job could be quantified in parts per million. This proved generally to be impossible. However, industrial hygiene and safety personnel in each plant were able to identify certain jobs and locations as involving the highest exposures in the plant, and to classify other exposures as medium or low relative to the "high" represented by the jobs with the greatest exposure. Therefore, the attempt

to obtain objective exposure data was abandoned, and each exposed job in the worker's history was scored 1, 2, or 3 to indicate low, medium or high estimated exposure.

This gross classification has two major failings, as a result of the subjective nature of the estimates. The first is that the scores represent estimated relative exposure within a given plant. It is therefore possible that, in objective terms, a "high" score in one plant corresponds to a "medium" or even "low" score in another. The second is that exposures in most plants have tended to decrease over time, so that a worker with long service may have had jobs in the remote past which involved "low" exposure relative to other jobs at that time, but which might be "high" in comparison with current exposures in the same job. This subjective classification is therefore of questionable validity in characterizing the exposure of a given worker. For epidemiological purposes, however, those who have high scores can reasonably be expected, on the average, to have had the greatest exposure, while those with low scores will have had the least, even though the true exposure in each group may vary considerably from person to person.

The estimated exposure history of each worker was summarized by calculating An Exposure Index (EI). This was done by multiplying the number of months on each job by the exposure score, totalling these overall exposed jobs, and dividing by the total number of months of exposure.

VI. FOLLOW-UP OF STUDY POPULATION

A follow-up procedure was instituted for those who had left employment and whose vital status could not be determined at the local plant. A portion of the terminations was traced by mail, using a form letter sent to the last

known address. Attachment D shows such a letter. If the letter was returned as undeliverable, the procedure shown in Figure 1 was used to obtain a current address.

The remainder of the terminations was traced through a retail credit bureau. The latter was also done for a portion of those who could not be found by the mail follow-up.

Approximately 5500 of the study population came from plants in which direct mail follow up was employed. In the remainder of the population follow-ups was by retail credit bureau. There was no significant difference between the success rates of the two methodologies.

Table 1 shows the vital status of the study population as of December 31, 1972. Follow-up is 85 percent complete, although follow-up data were still being received at the time the files were closed for this report. Of the 352 workers known to be dead, death certificates have been received for all but 24.

The mortality calculations in the succeeding sections of this report are based only on those workers who were successfully traced, which is equivalent to assuming that mortality among those not found is the same as among those who were found. In order to examine the reasonableness of this assumption, the differences between those found and not found are examined briefly as part of the following section.

VII. DESCRIPTION OF STUDY POPULATION

Table 2 shows the distribution of those successfully traced, by year of birth and number of months of exposure. The median birth year was 1931 (compared with 1920 for those not found) and the median duration of employment in an exposed job was 80 months (44 months in those not found). The

shortest exposures occur in both the oldest men, most of whose working life was over when vinyl chloride came into general use, and the youngest men, whose date of hire was closer to the closing date of the study.

Although nearly half of the study population had less than 60 months of exposed employment, there were nevertheless 855 workers with 20 years or more exposure, and 1641 with 15 years or more.

Table 3 shows the distribution by duration of exposure and the year in which exposure began. The median year in which exposure began was 1962 (versus 1953 for those not found). Notice that almost half the study population first entered exposed employment in the decade 1960-69. This reflects the fact that a great many plants first began vinyl chloride operations in that decade and means that these workers have not been followed for an substantial length of time after their exposure occurred.

Table 4 shows the relationship between duration of exposure and Exposure Index (EI). The median EI is 1.44 (1.75 for those not found). There does not appear to be a close relationship between the EI and the duration of exposure, that is, workers with a higher EI do not differ substantially in duration of exposure from those with a lower EI. One implication is that in assessing the relationship between mortality and exposure, both duration and level of exposure should be examined separately, as well as in combination.

The data shown above for those not found indicates that they were born (and began their exposure) about 10 years before the group on which follow-up was complete, and had about half the duration of employment with a slightly higher EI. Typically, their employment in an exposed job ended before 1960. The bulk of these workers came from older plants from

which data were collected late in the study.

Although there appear to be nothing very unusual about this group in terms of work history and exposure, it is nevertheless true that their exposures took place further back in time than that of the group successfully traced. It is therefore possible that their mortality, after a substantial latent period, might show a somewhat different pattern of mortality from the traced group.

VIII. CALCULATION OF RISK OF DEATH

The usual method of describing the risk of death in a study of this kind is to express the number of deaths which actually occurred as a percentage of the number which would have been expected in a comparable male population observed over the same age and time intervals. This statistic is called the Standardized Mortality Ratio (SMR).

In order to calculate the number of expected deaths it is first necessary to calculate the number of years for which each worker was observed, and to classify these years of observation by the age intervals into which they fell and the birth date of the worker.

Table 5 shows such a calculation for the 7128 vinyl chloride workers on whom follow-up was complete. This group represents 77,846 person-years of observation. For example, the table shows that for workers born in the years 1925-29, there were 2725 person-years of observation in the age interval 30-34, and an easy calculation shows that these person-years were lived in a period whose midpoint was about 1960. Therefore, the number of deaths to be expected in this cell of the table is the number of deaths occurring in 1960 among 2725 males aged 30-34 in a comparable non-exposed population.

In the present study the U. S. male population was used as the standard of comparison, and the age-specific U. S. male death rates for the appropriate ages and years were applied to the corresponding cells of Table 5 to obtain the expected deaths.¹ This was done for all causes of death together, and for each of the 35 causes for which detailed mortality rates are published on a national basis (omitting such causes as breast cancer, congenital malformations, and diseases of infancy).

Table 6 shows observed and expected deaths, and the SMR, for each of these causes for the total study group. In calculating the SMR's for specific causes, the 24 deaths for which no certificates were found were assumed to have the same cause distribution as those for which certificates were available.

In the standard population, each SMR would be equal to 100. Therefore, the statistical significance of the deviation of each SMR in the study population from the expected value of 100 was tested. A single asterisk indicates those SMR's which differed significantly from 100 at the 5 percent level, that is, which had a probability of .05 or less of occurring by chance. A double asterisk indicates those which were significant at the 1 percent level. SMR's based on fewer than 5 observed cases were not tested for significance.² The practical interpretation of these statistical statements will be examined below.

¹ Each diagonal in Table 5 corresponds to experience centered around a given year. For the first three diagonals (from left to right) 1950 rates were used; for the next three, 1955, 1959 and 1965 rates were used; for the last two, 1967 rates were used.

² Large-sample tests of significance were derived from: Chiang, C. L., "Standard error of the age-adjusted death rate," Vital Statistics Special Reports, 47 (1961) pp 275-285.

Table 7 shows the same SMR's for workers with an Exposure Index below 1.5 versus those at 1.5 or above. The dividing point of 1.5 represents a level halfway between "low" and "medium."

Table 8 shows similar results for workers with less than 5 years exposure versus those with 5 years or more.

In order to examine the possible interaction between duration and level of exposure, the study population was divided into 4 groups on the basis of both EI (low vs high) and duration of exposure (short vs long) using the same dichotomization as Tables 7 and 8.

Table 9 shows the results for short versus long exposure in the low EI group, and Table 10 shows the same comparison in the high EI group.

In each of the above tables, deaths for which certificates had not been received were assumed to be distributed as a uniform percentage of all causes. The cause specific SMR's were therefore adjusted upward by a percentage which varied in each subgroup. Table II shows the distribution of uncertified deaths in each group, and the resulting percentage by which the cause specific SMR's were increased.

IX. ANALYSIS

The overall mortality of the study population is statistically significantly lower than that of the U. S. male population. There were 352 observed deaths compared with 467 expected, for an SMR of 75.

Table 6 shows that no specific cause of death was statistically significantly greater than expected. Several, particularly heart disease, accidents and "other diseases" not detailed in the tables, were significantly below their expected values.

When the study population is divided according to length and duration of exposure (Tables 7 and 8) and combinations of these measurements (Tables 9 and 10) three major patterns emerge.

For malignant neoplasms as a whole, the SMR increases with increasing exposure, whether measured by level, duration, or both. In the high exposure group with 5 years or more exposure (Table 10) there are 36 observed cases and 26.11 expected.

For cardiovascular - renal diseases as a group, there are also increases in the SMR with increasing exposure, but the number of observed cases remain less than expected, the differences being statistically significant in all groups except the high exposure, long duration group in Table 11.

For all other causes, there are no consistent relationships with exposure.

Within the malignant neoplasms, the largest (although not statistically) significant SMR is in cancers of the buccal cavity and pharynx, with 5 observed, 2.84 expected, and an SMR of 189. However, Tables 7 to 10 show that all these cases have Exposure Indexes below 1.5, and 4 out of the 5 have less than 5 years exposure.

Cancer of the digestive system shows no excess in the study population as a whole. However, in those workers with Exposure Indexes of 1.5 or higher, there are 12 observed cases where 9.14 are expected (table 7). In the subgroup of the above workers with 5 years or more exposure, there are 11 observed cases and 7.47 expected.

Respiratory cancer shows a slight excess in the total group, and a similar pattern for different exposure categories, with 13 observed versus 10.28 expected when the Exposure Index is 1.5 or higher, and 12 observed versus 8.50 expected when, in addition, the duration of exposure is 5 years

or more.

Malignant neoplasms of other and unspecified sites show an excess in the total group, and an increase with both level and duration of exposure (Tables 7 and 8). The relationship with exposure is more pronounced, since those with exposures of less than 5 years have fewer cases than expected.

The lymphosarcomas, although occurring at about the expected rate when the whole group is considered, are concentrated almost entirely in the high exposure long duration group. In that category there are 4 cases observed and 1.84 expected.

Cancers of the genital and urinary organs, and leukemia, have fewer cases than expected. The number of cases is too small to examine any trends.

X. DISCUSSION

The favorable overall mortality of the study population is a phenomenon commonly observed in working populations. Studies of several occupational groups have shown that, even if there is an occupational hazard which greatly increases the risk of death from a particular cause, the overall mortality may well be favorable. Table 12 shows the SMR's obtained in a number of recent studies of other occupational groups. Although for technical reasons these SMR's are not strictly comparable, the overall mortality in the present study is clearly favorable.

SMR's which are higher than expected may be worthy of attention even if they are not statistically significant. This is especially true in the present study since the number of deaths from many causes is quite small, and even a relatively high SMR may not reach statistical significance.

If, in addition, a particular cause shows a consistent pattern of increase with exposure or estimated exposure, the findings are particularly

interesting.

By these criteria, mortality from digestive cancer, respiratory cancer, cancer of other and unspecified sites, and lymphosarcoma, appear to be related to exposure as estimated in this study.

In view of the association between vinyl chloride exposure and angiosarcoma of the liver, the digestive cancers were examined further to see what contribution angiosarcoma made to the observed mortality pattern.

Of the 19 digestive cancers, 7 were liver cancers, of which two were angiosarcomas according to the death certificate. However, among angiosarcoma deaths in vinyl chloride workers identified by other investigators, there were 6 which occurred in the present study population during the study period. They were all found in the course of the study, but 4 were listed on the death certificate as due to causes other than angiosarcoma. Table 13 shows these deaths with the death certificate cause.

If there had been no angiosarcomas, Table 13 shows that the total number of digestive cancers in the high exposure group would have dropped to 8, and the number with high exposure and 5 years or more exposure would have dropped to 2, so that the mortality pattern in this cause group is due entirely to angiosarcoma.

The other cause group worth further investigation is cancer of other and unspecified sites, both because it is a heterogeneous category and because it seems, unlike the other cancers, to be more related to duration than to level of exposures.

Table 14 shows a list of the specific causes included in this category, which is essentially brain cancer and generalized cancer with primary site

unknown. About 40 percent of the observed deaths were due to brain cancer. In the general male population, about 22 percent of this category is due to brain cancer, so that not only is the mortality from this cause excessive, but brain cancer is overrepresented.

The possibility exists based on the lack of specificity of some of the listed causes that some of the brain cancers were not primary, but metastases from another unidentified site such as the lung.

The cancers of the buccal cavity and pharynx are difficult to explain because of their occurrence in the low exposure - short exposure group. It is possible that this is a chance occurrence, or that exposures to other substances were involved.

XI. POSSIBLE BIASES IN THE CALCULATION OF RISK

There are three areas in which bias could have entered the calculation of the above SMR's. The first is the choice of the U. S. male population as a standard. The second is the absence of 15 percent of the population who were not found. The third is the discovery, as the study ended, of a group of 1500 workers whose exposures occurred up to 35 years ago, and who are not included in the study group.

The choice of the U. S. male population was not entirely appropriate. The study population was concentrated almost entirely in the Eastern half of the United States, whose normal mortality is slightly higher than that of the whole country. This implies that the correct expected values should have been slightly higher than the ones which were used, and the corresponding SMR's slightly lower. This has no practical significance except for those malignancies in which the SMR's were above 100. However, regional differences in mortality from those causes is such that these SMR's would have been at most 1.5 to 3 percent lower, with no substantive difference in the results.

The workers who could not be traced may conceivably contain a disproportionate number of deaths, which could increase the overall SMR. It appears unlikely that the distribution of causes would be substantially different. However, without actual data in hand, the possibility cannot be ruled out that either some new causes of death might become significant, or that some of the relationships found in this study might be diluted.

The same comments apply even more to the group of 1500 workers discovered too late to be included in the study group, since they would provide data in precisely the area (very long exposure and long latency) in which the present study is relatively weak.

References:

Archer, VE, Wagoner, JK, Lundin, FE, Jr: Cancer mortality among uranium mill workers. J Occup Med 15:11-14, 1973.

Enterline, PE, DeCoulle, P, Henderson, V: Mortality in relation to occupational exposure in the asbestos industry. J Occup Med 14:897-903, 1972.

Enterline, PE, Kendrick, MA: Asbestos-dust exposures at various levels and mortality. Arch Environ Health 15:181-186, 1967.

Hammond, EC, Selikoff, IJ: Relation of cigarette smoking to risk of death of asbestos-associated disease among insulation workers in the United States. Symposium on Biological effects of Asbestos, Lyon, October 2-5, 1972, paper #46.

Lloyd, JW, Ciocco, A: Long-term mortality study of steelworkers; methodology. J Occup Med 11:229-310, 1969.

Lundin, FE, Jr., Lloyd, JW, Smith, EM, Archer, VE, Holaday, DE: Mortality of uranium miners in relation to radiation exposure, hard-rock mining and cigarette smoking--1950 through September 1967. Health Physics 16:571-578, 1969.

Redmond, CK, Ciocco, A, Lloyd, JW, Rush, HW: Long-term mortality study of steelworkers--VI. Mortality from Malignant Neoplasms among coke oven workers. J Occup Med 14:621-629, 1972.

Tabershaw, IR, Cooper, WC, Balzer, JL: A Labor-management occupational health service in a construction industry. Arch Environ Health 21:784-788, 1970.

Table 1

Follow-up Status of 8384 Vinyl Chloride Workers

Total		Alive	Dead	Unknown	Death Certificates	
					Rec'd.	Not Rec'd.
Number	8384	6776	352	1256	328	24
Percent	100.0	80.8	4.2	15.0	92.7	7.3

April 15, 1974

Table 2

Distribution of Months in Exposed Employment by Birth Year,
for 7128 Vinyl Chloride Workers with Completed Follow-up

Year of birth	Total	Months of exposure							Unknown
		<60	60-119	120-179	180-239	240-299	300-359	360-419	
1875-79	1	1							
1880-84	5		2	2	1				
1885-89	7	1	3	3					
1890-94	21	3	7	9	2				
1895-99	38	10	4	9	12	2			1
1900-04	89	9	13	21	22	17	6		1
1905-09	214	33	27	35	37	43	34	3	2
1910-14	404	79	62	58	72	73	43	15	2
1915-19	648	145	96	80	120	105	84	13	5
1920-24	897	229	191	98	143	160	64	8	4
1925-29	940	276	214	118	169	139	20		4
1930-34	877	287	252	128	181	26			3
1935-39	986	454	378	120	27				7
1940-44	1169	664	481	16					8
1945-49	700	632	66						2
1950-54	132	132							
Total	7128	2955	1796	697	786	565	25	39	39

April 15, 1974

not
1073

Table 3

Distribution of Months in Exposed Employment by Year
in Which Exposure Began, for 7128 Vinyl Chloride Workers with Completed Follow-up

Year Exp. Started	Total	Months of Exposure							Unknown
		<60	60-119	120-179	180-239	240-299	300-359	360-419	
1930-39	35	2	4	1	4	6	13	5	
1940-49	1048	135	93	119	151	277	237	34	2
1950-59	1962	389	257	383	631	282			20
1960-69	3368	1714	1442	195					17
1970-71	715	715							
Total	7128	2955	1796	698	786	565	250	39	39

April 15, 1974

1001
1074

Table 4

Distribution by Months of Exposed Employment and
Exposure Index
for 7128 Vinyl Chloride Workers with Complete Follow-up

Exposure Index	Total	Months of exposure							Unknown
		<60	60-119	120-179	180-239	240-299	300-359	360-419	
1.0-1.4	4032	1715	1125	402	406	274	92	18	
1.5+	3057	1240	671	295	380	291	159	21	
Unknown	39								39
Total	7128	2955	1796	697	786	565	251	39	39

May 2, 1974

1004
1075

Table 5

Person Years of Observation by Year of Birth and
Age for Vinyl Chloride Workers

Age	Totals	Year of Birth															
		1875	1880	1885	1890	1895	1900	1905	1910	1915	1920	1925	1930	1935	1940	1945	1950
		1879	1884	1889	1894	1899	1904	1909	1914	1919	1924	1929	1934	1939	1944	1949	1954
Total	77846	13	92	124	317	705	1662	4055	6710	11038	14305	13807	9408	7239	6174	2023	174
15-19	134										1	11	12	5	53	21	31
20-24	4070								39	180	572	440	475	920	1301	143	
25-29	12076								32	335	1090	1882	1642	2173	4221	701	
30-34	13861							14	264	1054	2183	2725	2863	3778	980		
35-39	13550						3	221	697	1685	2742	3688	3706	808			
40-44	12031					5	70	524	1028	2044	3502	4113	745				
45-49	9527					56	241	710	1226	2631	3847	816					
50-54	6368				35	135	334	762	1574	2768	760						
55-59	3632			22	88	155	363	886	1636	482							
60-64	1684		14	35	77	142	341	822	253								
65-69	632	2	25	31	62	126	270	116									
70-74	202	5	23	20	43	71	40										
75-79	58	5	15	11	12	15											
80-84	16	1	10	5													
85-89	5		5														

April 15, 1974

Doc 1076

TABLE 6

OBSERVED DEATHS/EXPECTED DEATHS AND STANDARDIZED MORTALITY RATIOS IN VINYL CHLORIDE WORKERS

Cause of death with I.C.D. number	obs/exp	SMR ¹
All causes	352/467.28	75**
Tuberculosis (001-019)	1/5.71	19
Tuberculosis of respiratory system (001-008)	1/5.34	20
Malignant neoplasms (140-205)	79/77.16	110
Malignant neoplasms, buccal cavity and pharynx (140-148)	5/2.84	189
Malignant neoplasms, digestive organs and peritoneum (150-159)	19/21.67	94
Malignant neoplasms, respiratory system (160-164)	25/23.93	112
Malignant neoplasms, genital organs (170-179)	3/3.55	91
Malignant neoplasms, urinary organs (180-181)	1/3.60	30
Malignant neoplasms, other and unspecified sites (190-199)	17/11.75	155
Leukemia and aleukemia (204)	3/3.77	85
Lymphosarcoma, lymphatic and hematopoietic tissues (200-203, 205)	6/6.06	106
Diabetes mellitus (260)	7/6.31	120
Major cardiovascular and renal diseases (330-334, 400-468, 592-594)	155/207.46	80**
Vascular lesions affecting CNS (330-334)	13/24.48	57**
Rheumatic fever & chronic rheumatic heart dis. (400-402, 410-416)	5/6.87	78
Arteriosclerotic heart disease (420)	121/137.33	95
Nonrheumatic endocarditis (421, 422)	1/6.58	16
Hypertensive heart disease (440-443)	3/9.33	35
Other hypertensive disease (444-447)	3/2.60	124
Chronic & unspecified nephritis & renal sclerosis (592-594)	0/4.27	0
Influenza and pneumonia (480-493)	7/9.96	54*
Ulcer of stomach and duodenum (540, 541)	2/3.83	56
Appendicitis (550-553)	0/0.66	0
Hernia and intestinal obstruction (560, 561, 570)	1/1.51	71
Gastritis, duodenitis, enteritis and colitis (543, 571, 572)	1/1.31	82
Cirrhosis of liver (581)	3/15.60	21
Hyperplasia of prostate (610)	0/0.39	0
Symptoms, senility and ill-defined conditions (780-795)	1/7.34	15
All other diseases (residual)	21/45.78	49**
Motor vehicle accidents (810-835)	17/32.70	56**
Other accidents (800-802, 840-962)	18/30.64	63**
Suicide (963, 970-979)	16/16.83	102
Homicide (964, 980-985)	1/11.98	9
Number of workers	7128	
Person-years	77846	

¹SMR's adjusted for deaths with cause unknown. See Table 11.

*Significant at 5% level.

**Significant at 1% level.

May 2, 1974

1004
1077

TABLE 7

OBSERVED DEATHS/EXPECTED DEATHS AND STANDARDIZED MORTALITY RATIOS IN VINYL CHLORIDE WORKERS,
BY ESTIMATED LEVEL OF EXPOSURE

Cause of death with I.C.D. number	EI <1.5		EI ≥1.5	
	obs/exp	SMR ¹	obs/exp	SMR ¹
All causes	188/270.33	70**	157/195.68	80**
Tuberculosis (001-019)	0/3.38	0	0/2.33	0
Tuberculosis of respiratory system (001-008)	0/3.16	0	0/2.18	0
Malignant neoplasms (140-205)	37/44.28	90	41/32.67	134
Malignant neoplasms, buccal cavity and pharynx (140-148)	5/1.62	330	0/1.21	0
Malignant neoplasms, digestive organs and peritoneum (150-159)	7/12.50	60*	12/9.14	141
Malignant neoplasms, respiratory system (160-164)	11/13.56	86	13/10.28	135
Malignant neoplasms, genital organs (170-179)	2/2.30	93	1/1.43	75
Malignant neoplasms, urinary organs (180-181)	1/2.07	51	0/1.52	0
Malignant neoplasms, other and unspecified sites (190-199)	9/6.57	146	8/4.52	190
Leukemia and aleukemia (204)	1/2.18	49	2/1.57	136
Lymphosarcoma, lymphatic and hematopoietic tissues (200-203, 205)	1/3.48	31	5/2.54	212
Diabetes mellitus (260)	5/3.65	146	2/2.65	81
Major cardiovascular and renal diseases (330-334, 400-468, 592-594)	84/120.11	75**	69/86.99	85*
Vascular lesions affecting CNS (330-334)	7/14.42	52**	6/10.06	64
Rheumatic fever & chronic rheumatic heart dis. (400-402, 410-416)	3/3.98	80	2/2.85	75
Arteriosclerotic heart disease (420)	68/78.94	92	51/58.05	95
Nonrheumatic endocarditis (421, 422)	0/4.20	0	1/2.89	38
Hypertensive heart disease (440-443)	1/5.48	19	2/3.86	56
Other hypertensive disease (444-447)	1/1.52	70	2/1.07	201
Chronic & unspecified nephritis & renal sclerosis (592-594)	0/2.50	0	0/1.77	0
Influenza and pneumonia (480-493)	5/5.80	92	0/4.13	0
Ulcer of stomach and duodenum (540, 541)	1/2.21	48	1/1.60	68
Appendicitis (550-553)	0/0.39	0	0/0.27	0
Hernia and intestinal obstruction (560, 561, 570)	0/0.88	0	1/0.63	171
Gastritis, duodenitis, enteritis and colitis (543, 571, 572)	0/0.76	0	1/0.55	196
Cirrhosis of liver (581)	2/8.90	23	1/6.64	16
Hyperplasia of prostate (610)	0/0.25	0	0/0.14	0
Symptoms, senility and ill-defined conditions (780-795)	0/4.22	0	1/3.09	34
All other diseases (residual)	14/21.90	68*	6/15.89	41**
Motor vehicle accidents (810-835)	8/19.08	45**	9/13.46	72
Other accidents (800-802, 840-962)	11/17.83	66*	6/12.67	50**
Suicide (963, 970-979)	9/9.73	98	7/7.02	107
Homicide (964, 980-985)	0/6.98	0	1/4.94	21
Number of workers	4032		3057	
Person-years	45354		32108	

¹SMR's adjusted for deaths with cause unknown. See Table 11.

*Significant at 5% level.

**Significant at 1% level.

May 2, 1974

1001 1078

TABLE 8

OBSERVED DEATHS/EXPECTED DEATHS AND STANDARDIZED MORTALITY RATIOS IN VINYL CHLORIDE WORKERS
BY DURATION OF EXPOSED EMPLOYMENT

Cause of death with I.C.D. number	<60 months		≥60 months	
	obs/exp	SMR ¹	obs/exp	SMR ¹
All causes	94/140.53	67**	251/329.30	76**
Tuberculosis (001-019)	0/2.23	0	0/3.55	0
Tuberculosis of respiratory system (001-008)	0/2.07	0	0/3.33	0
Malignant neoplasms (140-205)	13/19.96	78	65/57.61	116
Malignant neoplasms, buccal cavity and pharynx (140-148)	4/0.70	688	1/2.16	47
Malignant neoplasms, digestive organs and peritoneum (150-159)	2/5.26	46	17/16.56	106
Malignant neoplasms, respiratory system (160-164)	3/5.52	65	21/18.51	116
Malignant neoplasms, genital organs (170-179)	0/0.99	0	3/2.76	112
Malignant neoplasms, urinary organs (180-181)	0/0.83	0	1/2.79	37
Malignant neoplasms, other and unspecified sites (190-199)	2/3.40	71	15/8.23	187
Leukemia and aleukemia (204)	1/1.25	96	2/2.53	81
Lymphosarcoma, lymphatic and hematopoietic tissues (200-203, 205)	1/2.01	60	5/4.07	126
Diabetes mellitus (260)	2/1.80	134	5/4.54	113
Major cardiovascular and renal diseases (330-334, 400-468, 592-594)	28/51.45	65**	125/157.39	81**
Vascular lesions affecting CNS (330-334)	4/5.97	81	9/18.71	49**
Rheumatic fever & chronic rheumatic heart dis. (400-402, 410-416)	2/2.39	101	3/4.53	68
Arteriosclerotic heart disease (420)	21/32.54	78*	98/125.39	96
Nonrheumatic endocarditis (421, 422)	0/1.79	0	1/5.35	20
Hypertensive heart disease (440-443)	1/2.42	49	2/7.01	30
Other hypertensive disease (444-447)	0/0.79	0	3/1.82	170
Chronic & unspecified nephritis & renal sclerosis (592-594)	0/1.55	0	0/2.76	0
Influenza and pneumonia (480-493)	3/2.93	122	2/7.09	29
Ulcer of stomach and duodenum (540, 541)	1/1.07	112	1/2.78	37
Appendicitis (550-553)	0/0.24	0	0/0.43	0
Hernia and intestinal obstruction (560, 561, 570)	0/0.42	0	1/1.10	93
Gastritis, duodenitis, enteritis and colitis (543, 571, 572)	1/0.40	301	0/0.92	0
Cirrhosis of liver (581)	1/4.49	26	2/11.18	18
Hyperplasia of prostate (610)	0/0.07	0	0/0.33	0
Symptoms, senility and ill-defined conditions (780-795)	1/2.28	52	0/5.09	0
All other diseases (residual)	4/11.17	40	16/26.34	63*
Motor vehicle accidents (810-835)	10/16.14	75	7/16.65	43**
Other accidents (800-802, 840-962)	7/12.94	65*	10/17.82	58*
Suicide (963, 970-979)	6/6.34	114	10/10.28	100
Homicide (964, 980-985)	1/5.80	20	0/1.20	0
Number of workers	2955		4134	
Person-years	34201		43240	

¹SMR's adjusted for deaths with cause unknown. See Table 11.

*Significant at 5% level.

**Significant at 1% level.

May 2, 1974

1079

TABLE 9

OBSERVED DEATHS/EXPECTED DEATHS AND STANDARDIZED MORTALITY RATIOS IN VINYL CHLORIDE WORKERS
WITH EXPOSURE INDICES BELOW 1.5, BY DURATION OF EXPOSED EMPLOYMENT

Cause of death with I.C.D. number	≤60 months exposure		≥60 months exposure	
	obs/exp	SMR ¹	obs/exp	SMR ¹
All causes	56/89.23	63**	132/181.28	73**
Tuberculosis (001-019)	0/1.41	0	0/1.97	0
Tuberculosis of respiratory system (001-008)	0/1.31	0	0/1.85	0
Malignant neoplasms (140-205)	8/12.86	73	29/31.46	95
Malignant neoplasms, buccal cavity and pharynx (140-148)	4/0.45	1036	1/1.17	88
Malignant neoplasms, digestive organs and peritoneum (150-159)	1/3.43	34	6/9.08	68
Malignant neoplasms, respiratory system (160-164)	2/3.58	6*	9/10.00	93
Malignant neoplasms, genital organs (170-179)	0/0.62	0	2/1.62	127
Malignant neoplasms, urinary organs (180-181)	0/0.55	0	1/1.53	67
Malignant neoplasms, other and unspecified sites (190-199)	1/0.97	120	8/4.43	187
Leukemia and aleukemia (204)	0/0.79	0	1/1.40	73
Lymphosarcoma, lymphatic and hematopoietic tissues (200-203,205)	0/1.26	0	1/2.23	46
Diabetes mellitus (260)	2/1.15	203	3/2.50	124
Major cardiovascular and renal diseases (330-334, 400-468, 592-594)	21/33.38	73**	63/86.82	75**
Vascular lesions affecting CNS (330-334)	2/3.93	59	5/10.51	49*
Rheumatic fever & chronic rheumatic heart dis. (400-402, 410-416)	2/1.50	155	1/2.49	41
Arteriosclerotic heart disease (420)	16/21.14	88	52/57.87	93
Nonrheumatic endocarditis (421, 422)	0/1.18	0	0/3.02	0
Hypertensive heart disease (440-443)	1/1.58	74	0/3.50	0
Other hypertensive disease (444-447)	0/0.50	0	1/1.02	101
Chronic & unspecified nephritis & renal sclerosis (592-594)	0/0.97	0	0/1.53	0
Influenza and pneumonia (480-493)	3/1.86	188	2/3.95	53
Ulcer of stomach and duodenum (540, 541)	0/0.68	0	1/1.53	67
Appendicitis (550-553)	0/0.15	0	0/0.24	0
Hernia and intestinal obstruction (560, 561, 570)	0/0.27	0	0/0.61	0
Gastritis, duodenitis, enteritis and colitis (543, 571, 572)	0/0.26	0	0/0.51	0
Cirrhosis of liver (581)	1/2.80	42	1/6.09	16
Hyperplasia of prostate (610)	0/0.07	0	0/0.20	0
Symptoms, senility and ill-defined conditions (780-795)	0/1.43	0	0/2.79	0
All other diseases (residual)	4/7.45	63	10/12.92	79
Motor vehicle accidents (810-835)	3/9.87	35	5/9.22	56
Other accidents (800-802, 840-962)	3/7.98	44	8/9.85	83
Suicide (963, 970-979)	3/4.08	86	6/5.66	109
Homicide (964, 980-985)	0/3.55	0	0/3.43	0
Number of workers	1715		2317	
Person-years	21418		23920	

¹SMR's adjusted for deaths with cause unknown. See table 11.

*Significant at 5% level.

**Significant at 1% level.

May 2, 1974

TABLE 10

OBSERVED DEATHS/EXPECTED DEATHS AND STANDARDIZED MORTALITY RATIOS IN VINYL CHLORIDE WORKERS
WITH EXPOSURE INDICES OF 1.5 OR GREATER, BY DURATION OF EXPOSED EMPLOYMENT

Cause of death with I.C.D. number	<60 months exposure		≥60 months exposure	
	obs/exp	SMR ¹	obs/exp	SMR ¹
All causes	38/47.93	79	119/147.81	81*
Tuberculosis (001-019)	0/0.76	0	0/1.57	0
Tuberculosis of respiratory system (001-008)	0/0.71	0	0/1.48	0
Malignant neoplasms (140-205)	5/6.57	96	36/26.11	141
Malignant neoplasms, buccal cavity and pharynx (140-148)	0/0.23	0	0/0.99	0
Malignant neoplasms, digestive organs and peritoneum (150-159)	1/1.67	76	11/7.47	151
Malignant neoplasms, respiratory system (160-164)	1/1.79	71	12/8.50	144
Malignant neoplasms, genital organs (170-179)	0/0.29	0	1/1.41	73
Malignant neoplasms, urinary organs (180-181)	0/0.26	0	0/1.26	0
Malignant neoplasms, other and unspecified sites (190-199)	1/1.18	107	7/3.51	204
Leukemia and aleukemia (204)	1/0.44	288	1/1.13	90
Lymphosarcoma, lymphatic and hematopoietic tissues (200-203, 205)	1/0.71	178	4/1.84	222
Diabetes mellitus (260)	0/0.61	0	2/2.04	100
Major cardiovascular and renal diseases (330-334, 400-468, 592-594)	7/16.54	54**	62/70.46	90
Vascular lesions affecting CNS (330-334)	2/1.87	135	4/8.19	50
Rheumatic fever & chronic rheumatic heart dis. (400-402, 410-416)	0/0.82	0	2/2.04	100
Arteriosclerotic heart disease (420)	5/10.41	61*	46/47.65	98
Nonrheumatic endocarditis (421, 422)	0/0.57	0	1/2.32	44
Hypertensive heart disease (440-443)	0/0.76	0	2/3.10	66
Other hypertensive disease (444-447)	0/0.27	0	2/0.81	253
Chronic & unspecified nephritis & renal sclerosis (592-594)	0/0.54	0	0/1.23	0
Influenza and pneumonia (480-493)	0/0.99	0	0/3.13	0
Ulcer of stomach and duodenum (540, 541)	1/0.35	362	0/1.25	0
Appendicitis (550-553)	0/0.08	0	0/0.19	0
Hernia and intestinal obstruction (560, 561, 570)	0/0.14	0	1/0.49	209
Gastritis, duodenitis, enteritis and colitis (543, 571, 572)	1/0.14	904	0/0.41	0
Cirrhosis of liver (581)	0/1.56	0	1/5.08	20
Hyperplasia of prostate (610)	0/0.01	0	0/0.13	0
Symptoms, senility and ill-defined conditions (780-795)	1/0.80	158	0/2.30	0
All other diseases (residual)	0/4.02	0	6/11.88	51*
Motor vehicle accidents (810-835)	7/6.05	146	2/7.43	28
Other accidents (800-802, 840-962)	4/4.73	107	2/7.96	26
Suicide (963, 970-979)	3/2.40	158	4/4.62	88
Homicide (964, 980-985)	1/2.18	58	0/2.76	0
Number of workers	1240		1817	
Person-years	12828		19305	

¹SMR's adjusted for deaths with cause unknown. See Table 11.

*Significant at 5% level.

**Significant at 1% level.

May 2, 1974

Table 11

Distribution of Deaths by Exposure Category
and by Whether Cause of Death Was Determined,
and Correction Factors Used for SMR Calculations

	Total*	Exposure Categories							
		Exposure Index		Total Exposure Time in Months		Exposure Index and Total Exposure in Months			
		<1.5	≥1.5	<60	≥60	<1.5 & <60	<1.5 & ≥60	≥1.5 & <60	≥1.5 & ≥60
All causes	352	188	157	94	251	56	132	38	119
Cause known	328	176	146	78	244	48	128	30	116
Cause unknown	24	12	11	16	7	8	4	8	3
Percent increase in SMR's	7.3	6.8	7.5	20.5	2.8	16.6	3.1	26.6	2.5

*Deaths in exposure categories do not add up to deaths in total category because 7 deaths had unknown exposure data, and one of those was of unknown cause.

May 2, 1974

Table 12

Results of Follow-up in Several Occupational Groups

<u>Study Population</u>	<u>Number Observed</u>	<u>Years of Observation</u>	<u>SMR All causes</u>	<u>Reference</u>
Asbestos building product workers	12,402	1951-1963	89	Enterline & Kendrick, 1967
Asbestos friction material workers	7,510	"	89	Enterline & Kendrick, 1967
Asbestos textile plant workers	1,843	"	121	Enterline & Kendrick, 1967
Cotton textile workers	6,281	"	88	Enterline & Kendrick, 1967
White underground uranium miners	3,414	1950-1967	159	Lundin et al, 1969
Steelworkers	59,072	1953-1961	82	Lloyd & Ciocco, 1969
Asbestos insulation workers > 20 years	250	1946-1965	140	Tabershaw et al, 1970
Coke oven workers	4,661	1951-1966	98	Richmond et al, 1972
Asbestos maintenance service, etc.	438	1941-1969	124	Enterline et al, 1972
Asbestos insulation workers > 20 years	370	1963-1971	198	Hammond & Selikoff, 1972
Asbestos insulation workers	17,800	1967-1971	136	Hammond & Selikoff, 1972
Uranium mill workers	662	1950-1967	99	Archer et al, 1973
Lead smelter workers	2,352	1946-1970	107	TCA, Inc., unpublished
Lead battery plant workers	4,680	1946-1970	99	TCA, Inc., unpublished
Petroleum refinery workers	18,916	1962-1971	68	TCA, Inc., unpublished

May 3, 1974

Table 13

Angiosarcoma Deaths in VCM Epidemiology Study Population

Study No.	Age at Death	Mos. Exposed	Cause as Given	Autopsy
4250	54	203	Cirrhosis of liver (sev. weeks)	yes
4255	60	281	Bleeding from hepatoma (3 days) (Laennec's cirrhosis)	yes
5765	45	167	Angiosarcoma rt. lobe of liver (10 mos.)	yes
7303	38	174	Liver failure (1 mo.) Cancer of liver, primary (15 mos.)	-
7376	52	238	Cardiac tamponade and massive left hemothorax (mins.) Widely metastatic angiosarcoma of liver (4 mos.)	-
7385	43	214	Hepatic failure Primary carcinoma liver	yes

April 15, 1974

Table 14

Other Malignancies (190-197, I.C.D.) in VCM Epidemiology Study Population

Study No.	Age at Death	Mos. Exposed	Cause as Given	Autopsy
2744	38	186	Widespread metastatic melanoma (1 yr.) Malignant melanoma of back	yes
3700	48	50	Malignant melanoma with widespread metastases	no
2096	67	69	Brain tumor (carcinoma) (8 mos.)	-
3946	43	64	Meningitis and pneumonitis (5 wks.) Ependymoma, fourth ventricle-brain (2 mos. post-op craniotomy)	no
4433	54	81	Astrocytoma, malignant-left cerebral hemisphere (1 yr.)	yes
4800	61	51	Carcinoma of the brain Arteriosclerosis Pneumonia due to static congestion	no
7279	57	271	Brain tumor, glioblastoma multiforme (18 mos.)	-
7300	44	211	Brain tumor, malignant (2 mos.)	-
7371	58	269	Brain tumor, malignant	no
7306	53	216	Thyroid carcinoma with metastases	no
5769	52	239	Carcinomatosis Carcinoma vertebral body	yes

Table 14
Other Malignancies (cont.)

Study No.	Age at Death	Mos. Exposed	Cause as Given	Autopsy
5246	67	201	Liposarcoma with metastasis (3 mos.)	yes
2850	64	193	Carcinomatosis, primary region not known (Past history, rheumatoid arthritis, myocardial infarction)	no
4778	54	300	Cardiac arrest (1 hr.) Respiratory arrest and cerebral hypoxia (2 days) Widespread metastatic carcinoma (2 mos.)	-
4786	61	318	Generalized metastasis undifferentiated (7 mos.) Squamous cell carcinoma, primary site undetermined	yes
5783	69	166	Metastatic carcinoma of abdomen Critical site undetermined (Pulmonary emphysema)	-
7301	55	133	Carcinomatosis (4 mos.) Primary site undetermined	no

April 15, 1974

Don 1086

Table 15

Malignant Neoplasms of Buccal Cavity and Pharynx (140-148, I.C.D.)
in VCM Epidemiology Study Population

Study No.	Age at Death	Mos. Exposed	Cause as Given	Autopsy
3003	31	49	Carcinoma of lip with metastasis to lung and neck (5 yrs.)	no
3431	56	30	Pulmonary edema (50 min.) Pulmonary metastases (Carcinoma, epidermoid, tongue and mandible)	no
3001	57	71	Metastatic squamous cell ca. - primary lesion palate	yes
7365	54	40	Atelectasis due to metastasis to mediastinum (4 mos.) Multiple malignant metastases to brain, liver & mediastinum (1 yr.) Adeno-carcinoma of nasal pharynx (2 yrs.)	yes
3354	63	26	Massive hemorrhage into tracheo-tree Status post laryngo-pharyngectomy, left radial neck dissection Well-differentiated keratinizing squamous cell carcinoma of left pyriform sinus (mos.)	yes

April 15, 1974

1004
1087

Doc 1086

INITIAL
STATUS
UNKNOWN

FOLLOWUP

INITIAL
STATUS
KNOWN




FIGURE 1

Manufacturing Chemists Association VCM Epidemiology Study

Parent Corporation _____

Plant Designation _____

Address _____

Telephone () - ext. _____

TABERSHAW-COOPER ASSOCIATES' CONTACT FOR SUBJECT STUDY:

Name _____

Title _____

Mailing Address _____

Street Address (please identify with sufficient precision
to serve as a taxi-driver's instruction)

Telephone () _____

Signed _____

Name _____

Title _____

Company _____

Address _____

Telephone () ext. _____

DOW 1089

Tabershaw-Cooper Associates, Inc.
2180 Milvia St., Suite 104
Berkeley, California 94704

Manufacturing Chemists Association VCM Epidemiological Study

Plant No. _____

1. When did this plant begin work involving vinyl chloride? _____
2. What is the approximate size of the current work force,
exclusive of secretarial and office personnel? _____
3. Approximately how many of these are employed in jobs in-
volving some exposure to vinyl chloride? _____
4. Please answer the appropriate one of the following
questions about your retirement plan.
 - a. No retirement plan _____
 - b. Voluntary plan, begun in _____
 - c. Required of all employees, begun in _____
 - d. Now required, formerly voluntary.
Voluntary plan begun in _____
Required plan begun in _____

Remarks: _____

5. Please answer the applicable questions below about
retention of records. How long are personnel records
kept
 - a. for those who retired but were not covered by a
retirement plan? _____
 - b. for those who retired under a retirement plan and
have died? _____
 - c. for those who quit or were discharged? _____
 - d. for those who died while employed? _____

Tabershaw-Cooper Associates, Inc.

page 2

6. Do the personnel records which are retained contain a record of job titles and dates of job title changes for the individual concerned?
7. Please list the job titles in the current work force which involve some exposure to vinyl chloride.

ATTACHMENT B (cont.)

DOW 1091

VCM Epidemiological Study - Individual Work History

Form B - Workers known to have died (please attach copy of death certificate)

1. Date of birth _____ Status at time of death:
5. Employed _____
2. Date hired _____
6. Retired (give date) _____
3. Date of death _____
7. Terminated (give date) _____
4. Place of death _____
8. If death certificate is not available, please give name of worker
so that death certificate can be obtained. No contact will be made
with relatives or attending physician.
-

Job history with estimated exposure

Job no.	Title	Check estimated exposure in ppm			
		450	50-199	200-500	7500
1	_____	from	_____		
		to	_____		
2	_____	from	_____		
		to	_____		
3	_____	from	_____		
		to	_____		
4	_____	from	_____		
		to	_____		
5	_____	from	_____		
		to	_____		
6	_____	from	_____		
		to	_____		
7	_____	from	_____		
		to	_____		

DOW 1092