

File: K-1711-(100) [1976] K-2516-(100) [1976]
 K-1712- [1976] K-2520- [1976]
 K-1716- [1976] K-2521- [1976]
 K-1717- [1976] K-2522- [1976]
 K-2197- [1976] K-2525- [1976]
 K-2511- [1976] K-9387- [1976]
 K-2515- [1976] K-9388- [1976]

CHEMICAL REACTIVITY, BIOTRANSFORMATION, AND TOXICITY OF POLYCHLORINATED ALIPHATIC COMPOUNDS

Authors: G. Bonse
 D. Henschler
 Institute of Toxicology
 University of Würzburg
 Würzburg, Germany

Referee: Perry J. Gehring
 Toxicology Research Laboratory
 Dow Chemical Company
 Midland, Michigan

INTRODUCTION

Chlorinated aliphatic compounds are released, in increasing number and quantity, into the environment. They are used on a large-scale basis as industrial solvents, in textile cleaning shops, and as plastic monomers, pesticides, and drugs. Some of the technically important products have long been known as hepatotoxic agents, and single members of the family (like vinyl chloride and vinylidene chloride) have recently been claimed to be carcinogenic; at present, trichloroethylene is also suspected as a potential carcinogen. In addition, several polychlorinated compounds possess a marked persistence in biological and abiotic systems and may accumulate to health-threatening concentrations.

The means and degree of metabolic transformation of chlorinated compounds, which may vary considerably, are influenced to a great extent by the nature of the chlorine substitution of differently hybridized C-atoms. The tendency of the behavior of the compounds under consideration with respect to metabolic and nonmetabolic reactivity can be particularly well demonstrated with poly- and perchlorosubstitutions. This report describes some aspects of the interrelationship

among polychlorinated aliphatics, chlorine substitution, chemical reactivity, and metabolic fate in biological systems, as well as their toxic properties. The molecular context is emphasized, whereas details of enzyme mechanisms and pathophysiological aspects of toxic reactions are not dealt with.

GENERAL CONSIDERATIONS ON CHLORINE SUBSTITUTION IN ALIPHATIC HYDROCARBONS

In general, chlorine substitution provides a decrease in the electron density of the involved C-atoms by their electron-attracting inductive effect (-I effect) which dominates the mesomeric donor effect. In nuclear magnetic resonance spectroscopy, the frequency of the resonance of an atomic nucleus depends on the electron density of its environment. The diamagnetic protection by the electrons diminishes the effective magnetic field at the nucleus. Thus, the shift of the resonance absorption may serve as a criterion for the electronic influence of substituents.¹ Recent publications²⁻⁵ reveal the possibility of a linear correlation between ¹³C-NMR shifts and the total electron density of the corresponding C-atoms.


Thus, the position of the ^{13}C -NMR signals is a measure of the electron-attracting effect of chlorine substitutions in organic molecules. The electronic shifts indicated in Table I clearly illustrate that an increase in chlorine substitution induces a decrease in electron density at the substituted C-atom; this correlates with a shift of resonance signals to lower fields. Substitution of hydrogen by chlorine in alkenes results in a smaller shift of ^{13}C -signals than it does in alkanes. This is probably due to the interaction of the chlorine with the π -bond. The inductive effect may be dominant in chlorinated alkanes, but in alkenes another effect (magnetic anisotropy, resonance effect) may contribute to the shift.⁸

The mesomeric substitution effect encountered with chlorine substituents may also be demonstrated separately by microwave and radiowave spectroscopy.^{9,10} One criterion of the mesomeric effect is the bond extension C-Cl which will be submitted in part to some kind of "double-bond character" and thus to a contraction by the participation of zwitter ionic structures. Thus, for example, the bond extension C-Cl is 1.76 Å with alkanes, 1.69 Å with alkenes (e.g., vinyl chloride), and 1.63 Å with alkynes (e.g., chloroacetylene). The proportion of the double-bond character of the C-Cl bond, according to nuclear quadrupole measurements, amounts to 5 to 6% in the case of vinyl chloride.


An enforced electron-attracting effect in connection with steric factors resulting from polychloro- or perchlorosubstitution induces decisive changes in reactivity as compared to

unsubstituted hydrocarbons. In the series of alkanes, an increasing chlorine substitution with an elongation of the carbon chain results in an enhanced tendency toward destabilization, which leads to C-C fissions or eliminations of HCl or chlorine with consequent formation of alkenes. On the other hand, alkenes are stabilized against electrophilic agents by increasing chlorine substitution; chlorinated olefins are characterized by an elevated thermal stability. Alkynes, unlike alkenes, are considerably destabilized by chlorine substitution.


ALKANES



The destabilizing effect in chlorinated alkanes is observed with an increase in the number of chlorine and carbon atoms. The steric hindrance of accumulated chlorine substituents, as well as their electron-attracting effect, favors C-C fissions (a, structure below) and eliminations of Cl_2 or HCl with the formation of more stable olefinic compounds (b, structure below).



For example, thermal decomposition of decachlorobutane (1) renders, by C-C fission, tetrachloroethylene (2) and hexachloroethane (3). Under dechlorinating conditions, hexachlorobutadiene (4) may easily be obtained from decachlorobutane (1).¹¹



C-A Fission


Up to now, investigations on metabolic conver-

sions of chlorinated compounds have been focused on C_1 and C_2 homologues of C_3 and C_4 have been included. Research on chlorinated methane converted metabolically to CO_2 ,¹² chloroform to methane to CO .¹⁴

A deeper insight into the hepatotoxicity of carbon tetrachloride was gained in 1961 by Butler¹⁵ who proposed a C-Cl fission with a methyl radical and chlorine as probable subsequent mechanism. The methyl radical with oxygen leads to the formation of peroxides and hydroperoxides. The latter react with iron to induce the decomposition of the chains.¹⁶ At the same time, it was demonstrated that carbon



Carbon tetrachloride forms a complex with cytochrome P-450 in which a redox reaction takes place.¹⁷



The complex cytochrome P-450 decomposes in the presence of carbon tetrachloride as the substrate.

The work of Hepp¹⁸ has established that enzyme-mediated C-Cl fission of chlorinated compounds occurs in the organism. According to Dyke and Chenoweth,¹⁹ the reaction involves shares all redox function oxidases. It is known that benzo-a-pyrene NADPH²⁰ High rates of metabolism found with this enzyme in chlorinated ethane ($\text{CHCl}_2-\text{CH}_3$), (1,1-dichloroethane ($\text{CHCl}_2-\text{CH}_2\text{Cl}$), and 1,1,2-dichloroethane ($\text{CHCl}_2-\text{CHCl}_2$). Eth-

TABLE I


^{13}C -NMR Shifts of Chlorinated Aliphatic Hydrocarbons (ppm Referring to CS_2 , = 0 ppm)⁸⁻¹²

	C_1	C_2	C_1	C_2
CH_4	194.9	CH_3-CH_3	186.9	186.9
CH_2Cl	167.7	$\text{CH}_2\text{Cl}-\text{CH}_3$	152.9	174.1
CHCl_2	138.6	$\text{CHCl}_2-\text{CH}_3$	123.6	160.3
CHCl_3	115.6	CCl_3-CH_3	96.9	146.5
CCl_4	96.8	$\text{CCl}_3-\text{CCl}_3$	87.3	87.3
		$\text{CH}_2=\text{CH}_2$	70.0	70.0
		$\text{CHCl}=\text{CH}_2$	66.7	75.4
		$\text{CCl}_2=\text{CH}_2$	64.1	77.3
		trans- $\text{CHCl}=\text{CHCl}$	71.7	71.7
		cis- $\text{CHCl}=\text{CHCl}$	73.5	73.5
		$\text{CCl}_2=\text{CHCl}$	67.8	75.3
		$\text{CCl}_2=\text{CCl}_2$	72.1	72.1


ons. In the series of
ne substitution with an
n chain results in an
a destabilization, which
eliminations of HCl or
formation of alkenes. On
are stabilized against
easing chlorine substitu-
are characterized by an
Alkynes, unlike alkenes,
ed by chlorine substitu-

VES

in chlorinated alkanes is
se in the number of
. The steric hindrance of
utients, as well as their
avors C-C fissions (a,
ions of Cl₂ or HCl
re stable olefinic
low).




decomposition of de-
ers, by C-C fission,
and hexachloroethane
conditions, hexachloro-
ay be obtained from




ons on metabolic conver-

sions of chlorinated aliphatic hydrocarbons have
focused on C₁ and C₂ compounds; higher
homologues of C₃ and C₄ units have occasionally
been included. Regarding the C₁ series (the
chlorinated methanes), carbon tetrachloride is
converted metabolically to chloroform and
CO₂,¹² chloroform to CO₂,¹³ and dichloro-
methane to CO.¹⁴

A deeper insight into the mechanism of the
hepatotoxicity of carbon tetrachloride was provid-
ed in 1961 by Butler¹⁵ who suggested a homolytic
C-Cl fission with the formation of trichloro-
methyl and chlorine radicals. At present, the most
probable subsequent pathochemical reaction
mechanism is the interaction of the trichloro-
methyl radical with unsaturated fatty acid chains,
with the formation of chloroform and a fatty acid
radical. The latter reacts with oxygen and forms
peroxides and hydroperoxides which, in turn,
induce the decomposition of the fatty acid
chains.¹⁶ At the same time, it was recently
demonstrated that the hepatotoxic activity of



Carbon tetrachloride forms an enzyme-
substrate complex with cytochrome P₄₅₀ of liver
microsomes²⁰ in which the following sequence of
reactions takes place.²¹



The complex cytochrome P₄₅₀-|CCl₃θ may then
decompose in the presence of protons, releasing
chloroform as the stable final product.

The work of Heppel and Porterfield²² has well
established that enzymes catalyze the oxidative
C-Cl fission of chlorinated organic compounds in
the organism. According to Van Dyke²³ and Van
Dyke and Chenoweth,²⁴ the enzyme system
involved shares all relevant properties with mixed
function oxidases. It is inducible by phenobarbital
and benzo-a-pyrene and requires oxygen and
NADPH.²⁵ High rates of dechlorination have been
found with this enzyme system in the series of
chlorinated ethanes: 1,1-dichloroethanes
(CHCl₂-CH₃), 1,1,2-trichloroethane
(CHCl₂-CH₂Cl), and 1,1,2,2-tetrachloroethane
(CHCl₂-CHCl₂). Ethanes with a trichloromethyl


bromotrichloromethane (BrCCl₃) is considerably
greater than that of CCl₄ or CHCl₃.¹⁷ There is a
correlation between the cytotoxic effects produc-
ed in vitro and in vivo and the bond-dissociation
energies for the fission of the three methane
derivatives (CHCl₃, CCl₄, and BrCCl₃): H-CCl₃,
95.7 kcal/mol; Cl-CCl₃, 73 kcal/mol; Br-CCl₃,
54 kcal/mol.^{18,19} A low dissociation energy
corresponds to an increased tendency for a
homolytic fission to the trichloromethyl radical.
Correspondingly, BrCCl₃ is considerably more
effective than CCl₄ in producing a peroxidative
breakdown of the microsomal lipids in rat liver.¹⁷

In principle, chlorine-substituted alkanes tend
to have radical reaction mechanisms because
chlorine substituents are capable of delocalizing
the unpaired electron via empty d orbitals of low
energy. Consequently, the resonance energy of the
CCl₃ radical amounts to 8.3 kcal/mol (as compar-
ed to the CH₃ radical = 0). Possible resonance
structures of the trichloromethyl radical are shown
below.


(CCl₃-) group [for example, 1,1,1-trichloro-
ethane (CCl₃-CH₃), 1,1,1,2-tetrachloroethane
(CCl₃-CH₂Cl), and 1,1,1,2,2-pentachloroethane
(CCl₃-CHCl₂)] are characterized by a low ten-
dency towards oxidative dechlorination; the same
is true for tetrachloroethylene (CCl₂=CCl₂).
Accordingly, acute and chronic hepatotoxicity are
high in the former and low in the latter group of
chlorinated aliphatics.²⁶⁻²⁸

Elimination of Chlorine

The elimination of chlorine with formation of
more stable alkenes represents the second type of
metabolic conversion of chlorinated alkanes. This
is the main pathway and the first step in the
metabolic conversion of pentachloroethane (5)²⁹
and hexachloroethane (6),³⁰ leading to the
formation of trichloroethylene (7) or tetrachloro-
ethylene (2), respectively.




However, an analogous mechanism with 1,1,2,2-tetrachloroethane (8) which would yield 1,2-dichloroethylene (9) could not be demonstrated up to now. The predominant pathway in the metabolism of 1,1,2,2-tetrachloroethane (8) goes through the sequence dichloroacetaldehyde-dichloroacetic acid (10-11), while the product of nonenzymatic dehydrochlorination (b) is found only in minor proportion; this reaction is already found in neutral phosphate buffer.³¹ In addition, traces of oxidatively formed tetrachloroethylene (2) are found.




Elimination of HCl

The elimination of hydrogen chloride from polychlorinated alkanes, leading to olefinic structures, also represents a major pathway in the metabolism of *p,p'*-dichlorodiphenyltrichloroethane (DDT) (12), with *p,p'*-dichlorodiphenyldichloroethylene (DDE) (13) as the first metabolite.³² This conversion is additionally catalyzed by the enzyme "DDT-dehydrochlorinase" (E.C. 4.5.1.1.) in the presence of glutathione.³³




Dehydrochlorinations are also discussed as primary metabolic pathways of polychlorinated cyclic alkanes, for example as in the case of 1,2,3,4,5,6-hexachlorocyclohexane.^{34,35} The γ -isomer of $C_6H_6Cl_6$ (14) is converted, by successive dehydrochlorination, via a γ -pentachlorocyclohexene to 1,2,4-trichlorobenzene (15),

which is then conjugated after aromatic hydroxylation.³⁵




ALKENES




Considerations on Stability

Poly- and perchlorosubstitution at sp^2 -hybridized C-atoms in unsaturated systems increases the thermal and chemical stability of these molecules. The $-I$ effect dominates the $+M$ effect of the chlorine substituents, thus resulting in a "deprivation" of the electron density of the double-bond system. This provides, in combination with a steric protective effect of the bulky chlorine substituents, an increased stability against electrophilic attack.

A good example of this effect is the reaction of chlorinated ethylenes with ozone. The relative rates of ozonization in the series ethylene:vinylchloride:trichloroethylene:tetrachloroethylene are 2500:1180:3.6:1.³⁶




(16)



(17)

Liquid butadiene (16) ($bp_{760} -5.6^\circ C$) polymerizes when left open to air for months after some period of induction; light irradiation accelerates the reaction. On the other hand, hexachlorobutadiene (17) ($bp_{760} 213^\circ C$) is stable up to $500^\circ C$ and cannot be brought to polymerization by pressure up to 100 atm (17), in addition, is characterized by a remarkable chemical stability, similar to the higher homologous stereoisomeric perchlorohexatrienes.³⁷ They are fairly stable against strong mineral acids and aqueous alkali even at high temperatures, but are converted to highly chlorinated vinyl ethers by alcoholic alkali at $80^\circ C$.³⁸ At least the action of fuming nitric acid at 130° and additional treatment with concentrated sulfuric acid at 170° are necessary to

convert hexachlorobutadiene to maleic acid anhydride.³⁹




Cyclooctatetraene is representative of cyclooctatetraenes containing double bonds, and is stable in air and oxygen. After an extensive heating, it converts to resinous substances. The molecule is again stable after the addition of halogens (O_2 , $KMnO_4$, and CrO_3) with ring contraction to chlorocyclooctatetraene. Cyclooctatetraene (20) could be used for thermal and chemical stability. The pressure tube for cyclooctatetraene (19) is stable up to $180^\circ C$. With elemental bromine, the composition C_8H_8 is stable up to $180^\circ C$. Cyclooctatetraene (20) does not react with aqueous $KMnO_4$ and is stable up to $180^\circ C$. The remarkable inertness of cyclooctatetraene can be determined in the related voluminous chlorinated compounds, which render an electrophilic attack. The value of an extraordinary stability is shown by systems by poly- and perchloroalkenes (Table 2).

Metabolism

Chlorinated ethylenes are converted to predominantly maleic acid anhydride. A significant interrelationship is observed between the stability of the molecule with increasing number of chlorine atoms and the metabolic stability of the molecule with all chlorinated

after aromatic hydroxyl-




ES

Cl


stitution at sp^2 -hybrid-
systems increases the
ity of these molecules,
ne +M effect of the
resulting in a "depriva-
y of the double-bond
bination with a steric
y chlorine substitu-
against electrophilic

fect is the reaction of
ozone. The relative
series ethylene:vinyl-
trichloroethylene are



$\text{bp}_{760} -5.6^\circ\text{C}$) poly-
air for months after
ant irradiation acceler-
er hand, hexachloro-
 $^\circ\text{C}$) is stable up to
nt to polymerization
(17), in addition, is
e chemical stability,
ogous stereoisomeric
v are fairly stable
and aqueous alkali
ut are converted to
s by alcoholic alkali
on of fuming nitric
al treatment with
 70° are necessary to

convert hexachlorobutadiene (17) to dichloro-
maleic acid anhydride.³⁹




Cyclononotetrene (18) ($\text{bp}_{760} 141^\circ\text{C}$), a
representative of cyclic polyolefines with alter-
nating double bonds, is sensitive against light and
oxygen. After an extended stay, or acceleration by
heating, it converts to a mixture of dimers and
resinous substances. The enormous reactivity of
the molecule is again demonstrated by the rapid
addition of halogens.^{40,41} Attack of oxidants
(O_2 , KMnO_4 , and CrO_3) easily results in products
with ring contraction. The transition to penta-
chlorocyclooctatetrene (19) and octachlorocyclo-
octatetrene (20) coincides with an increase in
thermal and chemical stability. After heating in
the pressure tube for 6 hr at 250°C , pentachloro-
cyclooctatetrene (19) can be isolated unchanged.
With elemental bromine it reacts to a product of
the composition $\text{C}_8\text{H}_3\text{Cl}_4\text{Br}_2$.⁴² Octachlorocyclo-
octatetrene (20) does not react with ozone or
aqueous KMnO_4 and cannot be brominated
under varying conditions. It remains thermally
stable up to 180°C .^{43,44} The reason for this
remarkable inertness of the double bonds should
be determined in the protection by the accumu-
lated voluminous chlorine atoms which, per se,
render an electrophilic attack more difficult due to
their -I effect. The validity of the general principle
of an extraordinary stabilization of olefinic
systems by poly- and perchlorosubstitution may
be further demonstrated by some paired examples
(Table 2).

Metabolism

Chlorinated ethylenes are metabolically
converted to predominantly C_2 alcohols and acids.
A significant interrelationship has been demon-
strated between the enhanced stability associated
with increasing numbers of chlorine substitutions
and the metabolic behavior in biologic systems
with all chlorinated ethylenes in the isolated

perfused rat liver preparation.^{51,52} At a given
prehepatic concentration of the ethylenes, a
significant rise in the proportion metabolized was
found in the series tetra-, tri-, and cis-1,2-dichloro-
ethylene vinylidene chloride.

The metabolic changes of chlorinated ethylenes
are initiated with the oxidation to corresponding
oxiranes⁵³⁻⁵⁶ by monooxygenases. Up to now
this class of compounds has only found limited
interest. The principal reactions of oxiranes in
biological systems are summarized in Figure 1. The
main toxic effects, acute as well as chronic, have
been associated with electrophilic reactions with
essential cellular components (alkylation), whereas
the other pathways (reduction, hydrolysis, and
conjugation, both enzymatically and nonenzymati-
cally) and rearrangements of carbonylic com-
pounds are considered to be detoxication
mechanisms. The type of rearrangement in chlori-
nated ethylenes depends on the number and
position of chlorine substitution(s); it goes to
either chlorinated aldehydes or acyl chlorides.



Further metabolic steps with the rearrangement
products are oxidations or reductions of aldehydes
to carboxylic acids of ethanol derivatives or
hydrolyses of acyl chlorides to corresponding
acids.

With tetrachloroethylene (2), the metabolic
formation of trichloroacetic acid (23) can
plausibly be explained by the primary formation
of an oxirane (21) and subsequent rearrangement
to trichloroacetyl chloride (22) and its hydroly-
sis.^{52,53} The transition of the oxirane to
trichloroacetyl chloride (21 to 22) is, in analogy,
found with the tetrachloroethylene oxide (2)
which can be synthesized by photooxidation and
which rearranges in vitro in different solvents to
trichloroacetyl chloride.⁵⁷








TABLE 2

Comparison of the Stability of Unsubstituted and Perchlorosubstituted Cyclic Polyolefines

 (45) Bicyclo-(4,2,0) octatrien-(1,5,7); some hours stable at room temperature, rapid polymerization at open air	 (46) Stable at room temperature, no tendency to polymerization, rearranges to aromatic compounds at 80°C
 (47) 3,4-Dimethylencyclobutene; rapid polymerization in the presence of light and air	 (48) Stable in air up to 160°C

R&S 026858

Com


It

H

FIG
(alk


Trichloroethene is converted to trichloroethanol by metabolic reaction to trichloroethanol (23) or trichloroacetic acid (25) as substantiated by the catalytic hydrogenation of the enzyme monooxygenase.

After administration of ethylene to the enzyme activity remains metabolites in




(23)

TABLE 2 (continued)
Comparison of the Stability of Unsubstituted and Perchlorosubstituted Cyclic Polyolefines



Heptafulvene; polymerization in solution already at -80°C



Stable in air up to 340°C





FIGURE 1. Synopsis of metabolic conversions and biological consequences of toxication (alkylation) and detoxication (conjugations, hydrolysis, and rearrangement) of alkenes.

Trichloroethylene (7) is metabolically converted to trichloroethylene oxide (24) and further converted to chloral.^{53-55,58,59} In subsequent metabolic reactions, chloral (25) is in part reduced to trichloroethanol (26) or oxidized to trichloroacetic acid (23).⁶⁰ The real occurrence of the oxirane (24) as a metabolic intermediate could be substantiated by spectroscopic investigations with the catalytic hemoprotein P_{450} of the converting enzyme monooxygenase.⁶¹

After administration of ^{36}Cl -labeled trichloroethylene to experimental animals, the specific activity remains unchanged in the recovered metabolites trichloroethanol and trichloroacetic


acid: no exchange of ^{36}Cl with the chlorine pool of the organism is observed.⁵³ Again, this is indicative of an intramolecular chlorine migration in the course of the transition of trichloroethylene oxide to chloral (24 to 25). Surprisingly, at first glance, the trichloroethylene oxide (24) which can be synthesized from trichloroethylene by photochemical oxidation⁶² thermally rearranges (in vitro) mainly to dichloroacetyl chloride (27).⁶³ In vivo, however, no dichloroacetic acid is found as a metabolite.^{52,64} This remarkable difference has important practical implications. In principle, there are three possible ways of rearranging the oxirane of trichloroethylene (24).



Regarding the two rearrangements leading to dichloroacetyl chloride, the hydride shift (2) has low probability, according to investigations by McDonald and Schwab.^{6,5} If one postulates that α -ketocarbonium ions or ion pairs^{6-6,8} are transition forms of the rearrangement (a synchronous process cannot, however, positively be ruled out^{6,9}), the carbonium ion (which after C-O heterolysis is solely destabilized by one direct, liganded Cl atom) should be indicative of a favored rearrangement. Way (3) would involve a carbonium ion with two direct liganded Cl atoms, consequently giving a lower degree of probability. Experiments on the behavior of the oxirane are in accordance with this assumption; rearrangement to



dichloroacetyl chloride is accelerated by tertiary amines⁷⁰ which, after addition of methanol, render dichloroacetic acid methyl ester.⁷¹ Upon heating to 100 to 140°C for 4 hr in a glass tube, the oxirane is converted in high yields (88%) into dichloroacetyl chloride, and only a small proportion of chloral can be detected.^{6,3}

The rearrangement of trichloroethylene oxirane to chloral in vitro can only be elicited by Lewis acids such as $AlCl_3$ or $FeCl_3$.^{5,2} Thus, it has been suggested that in the living organism the "environment" at the site of formation of the oxirane (24) might possess electron acceptor properties and induce the rearrangement in the direction of chloral.^{5,1,5,2} The most plausible mechanism for the influence of




Lewis acids is an interaction with the oxirane (24) at the site where there is a steric opening of the molecule with the oxirane oxygen, with the



nearest chlorine atom, or with both, inducing the final rearrangement according to shift (3) to chloral (25).



Cis- and trans-1,2 (29) both form the (32) and dichloroacetyl chloride identified in perfusion preparation.^{5,1,5,2} The compound obtained by the primary



The isomeric 1,2-dichloroethane (33) is methyl chloride (36).^{6,4,7,2} If 1,2-dichloroethane with *m*-chloroperbenzoic acid (35) (the epoxide of the oxirane) can be formed, it seems to be extremely




R&S 026860

II
Cl


erated by tertiary
on of methanol,
nyl ester.⁷¹ Upon
hr in a glass tube,
yields (88%) into
y a small propor-
3
roethylene oxirane
ited by Lewis acids
it has been suggest-
e "environment" at
ne (24) might poss-
and induce the re-
f chloral.^{51,52}
for the influence of

both, inducing the
to shift (3) to




Cis- and trans-1,2-dichloroethylenes (28 and 29) both form the metabolites dichloroethanol (32) and dichloroacetic acid (11) which have been identified in perfusates of the isolated rat liver preparation.^{51,52} Their formation can be explained by the primary intermediate oxiranes (30

and 31) and subsequent rearrangement to dichloroacetaldehyde (10), which is then oxidized to dichloroacetic acid (11) or reduced to dichloroethanol (32). The thermal rearrangement of the oxiranes that occurs with chlorine migration to dichloroacetaldehyde (10) has been demonstrated.⁶⁶



The isomeric *1,1-dichloroethylene* (vinylidene chloride) (33) is metabolized to monochloroacetic acid (36).^{64,72} If 1,1-dichloroethylene is oxidized with *m*-chloroperbenzoic acid, only chloroacetyl chloride (35) (the expected rearrangement product of the oxirane) can be isolated. The oxirane itself seems to be extremely unstable and resistive to





synthesis under varying conditions.⁷³ However, from these findings it can be speculated that the oxirane (34) is formed as a metabolic intermediate and rearranges spontaneously to chloroacetyl chloride (35), which is subsequently hydrolyzed to monochloroacetic acid (36).



Monochloroacetic acid (36) is likewise formed as a metabolite of *monochloroethylene* (vinyl chloride) (37).⁷⁴⁻⁷⁶ One possible explanation of the formation of monochloroacetic acid (36) is the primary oxidation of monochloroethylene (37) to vinyl chloride oxirane (38), which rearranges to chloroacetaldehyde⁷⁴ (33) followed by further metabolic oxidation to chloroacetic acid (36). In analogy, monochloroacetaldehyde (39) is formed as a rearrangement product of the oxirane of vinyl chloride, which can be synthesized by chlorination of ethylene oxide (40) in the gaseous phase.^{77,78}

The further metabolic fate of vinyl chloride oxirane has been elucidated in part. A very small proportion of metabolized vinyl chloride is bound covalently to tissue components, as has been shown *in vitro*⁷⁹ and *in vivo*.⁸⁰ Major metabolites *in vivo* are 2-hydroxyethylcysteine⁸¹ (41), 2-hydroxyethyl-N-acetylcysteine⁸¹ (42), carboxymethylcysteine⁸² (43), and thiodiacetic acid^{81,82} (44). The mechanism of the formation of these conjugates can easily be explained by the previous rearrangement of the oxirane to chloroacetaldehyde (39) and oxidation to chloroacetic acid (36) which reacts with glutathione; carboxymethylglutathione is converted to carboxymethylcysteine (43), and the latter, by further oxidations, is converted to thiodiacetic acid (44). Yllner has identified carboxymethylcysteine (43) and thiodiacetic acid (44) as the main metabolites⁸³ in the urine of mice after dosing with chloroacetic acid. The formation of S-(2-chloroethyl)-cysteine and its N-acetyl conjugate⁷⁶ as products of simple addition reactions has been questioned;⁸¹ in fact, this hypothesis is inconsistent with an enzymatic mechanism postulated according to the results of experiments *in vivo* with enzyme inhibitors.^{74,80} The formation of a cyclic vinyl chloride peroxide, in equilibrium with an oxirane-singlet oxygen complex, might account for the observed formation of formaldehyde as a decomposition product and oxidation to carbon dioxide, which has been identified in part, *in vivo*, after metabolic incorporation into urea or methionine and serine.⁷⁶ Another interesting hypothesis is the addition of vinyl chloride to hydrogen sulfide under formation of bis-(2-chloroethyl)-sulfide, which then might account for the formation of thiodiacetic acid⁷⁶ (44); the alkylating intermediate could be suspected as one of the carcinogenic metabolites of vinyl chloride. However, further experiments, particularly in iso-

lated biological systems, are necessary to confirm or reject these assumptions.



Mutagenicity and Carcinogenicity

Three chlorinated ethylenes exert mutagenic effects: vinyl chloride,⁸⁴⁻⁸⁶ vinylidene chloride,^{86,87} and trichloroethylene⁸⁶ (see Figure 2). The common molecular feature of these compounds is the formation of unsymmetric oxiranes,^{86,88} the stability of which (as tested in non polar solvents) is far less^{63,73,77} than that of the others which form symmetric oxiranes^{57,66} and are not mutagenic. Oxiranes may react *in vivo* enzymatically or nonenzymatically with SH-compounds to form nonactive conjugates.⁸⁹ On the other hand, the highly electrophilic oxiranes may react directly with nucleophilic constituents of the animal cell as a first step in a genotoxic effect. If this mechanism is taken to be essential for the mutagenic effects (as has been elucidated in the case of vinyl chloride),⁸⁵ the interrelationship between symmetry/asymmetry and stability/instability of the oxiranes might offer an important rule⁸⁸ and possibly a useful criterion which may be used to measure chemical reactivity and predict biologic activity (see Figure 3). Vinyl chloride has been demonstrated to be carcinogenic




FIGURE 2. The compounds activating the different methyltry-







FIG. 3. of c. Fig.

R&S 026862

necessary to confirm



city
nes exert mutagenic
2, 84-86 vinylidene
ethylene⁸⁶ (see Figure
ar feature of these
ion of unsymmetric
of which (as tested in
3,73,77 than that of
metric oxiranes^{57,66}
anes may react in vivo
matically with SH-
ave conjugates.⁸⁹ On
electrophilic oxiranes
eophilic constituents
step in a genotoxic
taken to be essential
has been elucidated
ride),⁸⁵ the interre-
etry/asymmetry and
ranes might offer an
oly a useful criterion
are chemical reactivity
(see Figure 3). Vinyl
ted to be carcinogenic







FIGURE 2. Mutagenicity of chlorinated ethylenes in an in vitro test system. Incubation of the compounds in concentrations (aqueous phase) from 0.9 to 10.9 ml/l in a metabolic activating microsomal system. Results as percent of spontaneous (= 100%) mutation rate in different operons of the test germ *E. coli* K₁₂: gal⁺, galactose; arg⁺, arginine; MTR⁺, methyltryptophane; nad⁺, NAD.⁸⁴




tetrachloro-
ethylene




trichloro-
ethylene




cis-1,2-
dichloroethylene



1,1-dichloro-
ethylene



trans-1,2-
dichloroethylene



vinyl chloride

symmetric
rel. stable
not mutagenic

asymmetric
unstable
mutagenic


FIGURE 3. Molecular features of oxidative metabolic intermediates (oxiranes) of chlorinated ethylenes in relation to their mutagenic potential in vitro (see Figure 2).

in humans⁹⁰ and experimental animals.⁹¹ In high oral doses, trichloroethylene produces malignancies in mice.⁹² Vinylidene chloride also has been reported as an animal carcinogen.⁹³

ALKYNES



In general, chlorine substitution of sp-hybridized C-atoms in allenes and acetylenes results in a destabilization of the molecules. This has been demonstrated with perchloroallene (45), which is stable only below -50°C and starts dimerization from -30°C upward.^{94,95} Similar behavior is encountered with tribromoallene, perbromoallene, and trichloroallenes with electronegative substituents.⁹⁶⁻⁹⁹



(45)



(46)



(47)

Dichloroacetylene (46) is also extremely reactive (explosive) and decomposes immediately, in the presence of air, to phosgene and carbon monoxide.¹⁰⁰ The destabilizing effect of chlorine substitutions at sp-hybridized bonds has also been demonstrated (though in a less intensity) with perchlorobutyne¹⁰¹ (47); this thermally unstable compound forms (slowly above 25°C) a dimer, C_8Cl_8 .

The metabolic fate of chlorinated acetylenes and allenes has not yet been investigated. Dichloroacetylene which is formed from trichloroethylene, tetrachloroethane(s), or acetylene in some working environments produces a highly characteristic symptomatology of acute intoxication: irreversible damage to the cranial nerves (predominantly the nervus trigeminus) in humans¹⁰² as well as experimental animals¹⁰³⁻¹⁰⁵ and tubular nephrotoxicity in animals.^{104,105} Whether these toxic effects are due to the reactive dichloroacetylene or to metabolites is open to further experimentation.

REFERENCES

- Dailey, B. P. and Shoolery, J. N., The electron withdrawal power of substituent groups, *J. Am. Chem. Soc.*, **77**, 3977, 1955.
- Tokuhiro, T. and Fraenkel, G., Origin of linearity of carbon-13 shift with charge. Calculations for the azines, *J. Am. Chem. Soc.*, **91**, 5005, 1969.
- Bloor, J. E. and Breen, L., Valence shell calculations on polyatomic molecules. II. CNDO SCF calculations on monosubstituted benzenes, *J. Phys. Chem.*, **72**, 716, 1968.
- Kajimoto, O. and Fueno, T., Inverse substituent effect in the ¹³C chemical shifts of *trans*-1-substituted-1,3-butadienes. Direct evidence of bond polarization effected by substituents, *Tetrahedron Lett.*, p. 3329, 1972.
- Miyajima, G., Takahashi, K., and Nishimoto, K., Carbon-13 nuclear magnetic resonance spectroscopy. IX. Monosubstituted ethylenes, *Org. Magn. Reson.*, **6**, 413, 1974.
- Hawkes, G. E., Smith, R. A., and Roberts, J. D., Nuclear magnetic resonance spectroscopy. Carbon-13 chemical shifts of chlorinated organic compounds, *J. Org. Chem.*, **39**, 1276, 1974.
- Lauterbur, P. C., Some applications of ¹³C nuclear magnetic resonance spectra to organic chemistry, *Ann. N.Y. Acad. Sci.*, **70**, 841, 1958.
- Miyajima, G. and Takahashi, K., Carbon-13 nuclear magnetic resonance spectroscopy. III. Chloro-substituted ethanes and ethylenes, *J. Phys. Chem.*, **75**, 331, 1971.
- Goldstein, J. H., Quadrupole coupling and bond character in the vinyl halides, *J. Chem. Phys.*, **24**, 106, 1956.
- Meal, H. C., Zeeman quadrupole spectra of *p*-chloroaniline and *p*-chlorobenzylchloride, *J. Chem. Phys.*, **24**, 1011, 1956.
- Roedig, A., Über Perchlorierung und Chlorolyse in der Butan- und Pentan-Reihe, *Leibigs Ann. Chem.*, **574**, 122, 1951.
- Uehleke, H., Hellmer, K. H., and Tabarelli, S., Binding of ¹⁴C-carbon tetrachloride to microsomal proteins in vitro and formation of CHCl₃ by reduced liver microsomes, *Xenobiotica*, **3**, 1, 1973.
- Fry, B. J., Taylor, T., and Hathway, D. E., Pulmonary elimination of chloroform and its metabolite in man, *Arch. Int. Pharmacodyn. Ther.*, **196**, 98, 1972.
- Kubic, V. L. and A., *Metab. Dispos.*, **3**, 1.
- Butler, T. C., Redox reactions in vitro by tissues and cells, *Arch. Biochem. Biophys.*, **17**, 1, 1967.
- Recknagel, R. O., *Crit. Rev. Toxicol.*, **1**, 1, 1967.
- Koch, R. R., *Chem. Rev.*, **67**, 1, 1967.
- Kerr, J. A., *Bond Dissociation Energies*, 1967.
- Benson, S. W., *Bond Dissociation Energies*, 1967.
- McLean, A. E. M., *Chem. Rev.*, **67**, 1, 1967.
- Ullrich, V. and Se., *Drug Metab. Dispos.*, **1**, 1, 1967.
- Heppel, L. A. and J., *Biol. Chem.*, **17**, 1, 1967.
- Van Dyke, R. A., *Enzymes*, **1**, 1, 1967.
- Van Dyke, R. A., *Enzymes*, **2**, 1, 1967.
- Van Dyke, R. A., *Enzymes*, **3**, 1, 1967.
- Spencer, H. C., *Redox reactions in vitro*, *Biochem. Pharmacol.*, **1**, 1, 1967.
- Torkelson, T. R., *Organic chemistry of laboratory animals*, *Chem. Rev.*, **67**, 1, 1967.
- Carpenter, C. P., *Interpretation of results*, *Metabolism*, **1**, 1, 1967.
- Yllner, S., *Metabolism*, **1**, 1, 1967.
- Fowler, J. S. L., *Science*, **130**, 530, 1969.
- Yllner, S., *Metabolism*, **1**, 1, 1967.
- Hayes, W. J., *Quinoline with different degrees of substitution*, *Chem. Rev.*, **67**, 1, 1967.
- Agosin, M., *Present position*, *Chem. Rev.*, **67**, 1, 1967.
- Sternburg, J. and A., *Econ. Entomol.*, **1**, 1, 1967.
- Grover, P. L. and A., *Biochem. J.*, **1**, 1, 1967.
- Williamson, D. G., *Am. Chem. Soc.*, **1**, 1, 1967.
- Roedig, A., *Vollständiges Triacetylen, Liebigs Ann. Chem.*, **574**, 122, 1951.
- Roedig, A. and B., *aus Perchlorbutadien*, *Chem. Ber.*, **106**, 1363, 1973.
- Forsch, M. and A., *Chem. Ber.*, **106**, 1363, 1973.
- Keppe, W., *Schmelze Cyclooctatetraen*, *Chem. Ber.*, **106**, 1363, 1973.
- Huisgen, R., *Bochum, Am. Chem. Soc.*, **1**, 1, 1967.
- Roedig, A. and A., *Chem. Ber.*, **106**, 1363, 1973.
- Roedig, A., *Heft octatetraen an*, *Chem. Ber.*, **106**, 1363, 1973.
- Roedig, A., *Bochum, Polyhalogenide*, *Chem. Ber.*, **106**, 1363, 1973.
- Eisenhuth, L., *Chem. Ber.*, **106**, 1363, 1973.
- Roedig, A., *Bochum, Chem. Ber.*, **106**, 1363, 1973.

to) is also extremely decomposes immediately, to phosgene and carbon. Utilizing effect of chlorine sized bonds has also been in a less intensity) with (7); this thermally un- (slowly above 25°C) a

of chlorinated acetyl- not yet been investi- which is formed from chloroethane(s), or acetyl- environments produces symptomatology of reversible damage to dominantly the nervus as well as experi- and tubular nephro- 105 Whether these the reactive dichloro- ites is open to further

J. Am. Chem. Soc., 77,

utions for the azines, J. Am.

DO SCF calculations on

of trans-1-substituted-1,3- p. 3329, 1972.

ance spectroscopy. IX.

copy. Carbon-13 chemical

anic chemistry, Ann. N.Y.

III. Chloro-substituted

of. Phys., 24, 106, 1956.

Chem. Phys., 24, 1011,

Ann. Chem., 574, 122,

65 microsomal proteins in vitro

metabolite in man, Arch.

14. Kubic, V. L. and Anders, M. W., Metabolism of dihalomethanes to carbon monoxide. II. In vitro studies, *Drug Metab. Dispos.*, 3, 104, 1975.
15. Butler, T. C., Reduction of carbon tetrachloride in vivo and reduction of carbon tetrachloride and chloroform in vitro by tissues and tissue constituents, *J. Pharmacol. Exp. Ther.*, 134, 311, 1961.
16. Recknagel, R. O. and Glende, E. A., Jr., Carbon tetrachloride hepatotoxicity: An example of lethal cleavage, *C.R.C. Crit. Rev. Toxicol.*, 2, 263, 1974.
17. Koch, R. R., Glende, E. A., Jr., and Recknagel, R. O., Hepatotoxicity of bromotrichloromethane - Bond dissociation energy and lipoperoxidation, *Biochem. Pharmacol.*, 23, 2907, 1974.
18. Kerr, J. A., Bond dissociation energies by kinetic methods, *Chem. Rev.*, 66, 465, 1966.
19. Benson, S. W., Bond energies, *J. Chem. Educ.*, 42, 502, 1965.
20. McLean, A. E. M., Effect of halogenated hydrocarbons on microsomal cytochrome (P-450), *Biochem. J.*, 103, 13P, 1967.
21. Ullrich, V. and Schnabel, K. H., Formation and binding of carbanions by cytochrome P-450 of liver microsomes, *Drug Metab. Dispos.*, 1, 176, 1973.
22. Heppel, L. A. and Potterfield, V. T., Enzymatic dehalogenation of certain brominated and chlorinated compounds, *J. Biol. Chem.*, 176, 763, 1948.
23. Van Dyke, R. A., Metabolism of volatile anesthetics. III. Induction of microsomal dechlorinating and ether-cleaving enzymes, *J. Pharmacol. Exp. Ther.*, 154, 364, 1966.
24. Van Dyke, R. A. and Chenoweth, M. B., The metabolism of volatile anesthetics. II. In vitro metabolism of methoxyflurane and halothane in rat liver slices and cell fractions, *Biochem. Pharmacol.*, 14, 603, 1965.
25. Van Dyke, R. A. and Wineman, C. G., Enzymatic dechlorination. Dechlorination of chloroethanes and propanes in vitro, *Biochem. Pharmacol.*, 20, 463, 1971.
26. Spencer, H. C., Rowe, V. K., Adams, E. M., McCollister, D. D., and Irish, D. D., Vapor toxicity of ethylene dichloride determined by experiments on laboratory animals, *Arch. Ind. Hyg.*, 4, 482, 1951.
27. Torkelson, T. R., Oyen, F., McCollister, D. D., and Rowe, V. K., Toxicity of 1,1,1-trichloroethane as determined on laboratory animals and human subjects, *Am. Ind. Hyg. Assoc. J.*, 19, 353, 1959.
28. Carpenter, C. P., Smyth, H. F., Jr., and Pozzani, U. V., The assay of acute vapor toxicity and the grading and interpretation of results on 96 chemical compounds, *J. Ind. Hyg.*, 31, 343, 1949.
29. Yllner, S., Metabolism of pentachloroethane in the mouse, *Acta Pharmacol. Toxicol.*, 29, 481, 1971.
30. Fowler, J. S. L., Some hepatotoxic actions of hexachloroethane and its metabolites in sheep, *Br. J. Pharmacol.*, 35, 530, 1969.
31. Yllner, S., Metabolism of 1,1,2,2-tetrachloroethane-¹⁴C in the mouse, *Acta Pharmacol. Toxicol.*, 29, 499, 1971.
32. Hayes, W. J., Quinby, G. E., Walker, K. C., Elliot, J. W., and Upholt, W. M., Storage of DDT and DDE in people with different degrees of exposure to DDT, *AMA Arch. Ind. Health*, 18, 398, 1958.
33. Agosin, M., Present status of biochemical research on the insecticide resistance problem, *Bull. Who.*, 29(Suppl.), 69, 1963.
34. Sternburg, J. and Kearns, C. W., Pentachlorocyclohexene, an intermediate in the metabolism of lindane by house flies, *J. Econ. Entomol.*, 49, 548, 1956.
35. Grover, P. L. and Sims, P., The metabolism of γ -2,3,4,5,6-pentachlorocyclohex-1-ene and γ -hexachlorocyclohexane in rats, *Biochem. J.*, 96, 521, 1965.
36. Williamson, D. G. and Cvetanovic, R. J., Rates of reactions of ozone with chlorinated and conjugated olefines, *J. Am. Chem. Soc.*, 90, 4248, 1968.
37. Roedig, A., Voss, G., and Kuchinke, E., Über α - und β -Perchlor-hexatrien-(1,3,5) und deren Reduktion zum Triacetylen, *Liebigs Ann. Chem.*, 580, 24, 1953.
38. Roedig, A. and Bernemann, P., Zur Kenntnis von hochchlorierten Vinylathern. 1-Äthoxy-pentachlor-butadien-(1,3) aus Perchlor-butadien, *Liebigs Ann. Chem.*, 600, 1, 1956.
39. Försch, M. and Roedig, A., Notiz zur Darstellung von Dichlormaleinsäureanhydrid aus Hexachlorbutadien, *Chem. Ber.*, 106, 1363, 1973.
40. Reppe, W., Schlichting, O., Klager, K., and Toepel, T., Cyclisierende Polymerisation von Acetylen. I. Über Cyclooctatetraen, *Liebigs Ann. Chem.*, 560, 1, 1948.
41. Huisgen, R., Boche, G., and Huber, H., The halogenation of cyclooctatetraene via 8-halohomotropylum ions, *J. Am. Chem. Soc.*, 89, 3345, 1967.
42. Roedig, A. and Bonse, G., unpublished results.
43. Roedig, A., Helm, R., West, R., and Smith, M., Structures of the dimers of perchlorobutyne. Octachlorocyclo-octatetraene and octachlorobicyclo-[4.2.0]-octatriene-[2.4.7], *Tetrahedron Lett.*, p. 2137, 1969.
44. Roedig, A., Bonse, G., and Helm, R., Cyclooctatetraen-dihydropentalen-umlagerung von Perchlor-COT und einige Polyhalogenderivate des Bicyclo [3.3.0] octans, *Chem. Ber.*, 106, 2156, 1973.
45. Eisenhuth, L. and Hopf, H., Gas phase pyrolysis of trans- and cis-1,2-diethynylcyclobutane. A new entry in the C₄ H₂ energy surface, *J. Am. Chem. Soc.*, 96, 5667, 1974.
46. Roedig, A., Bonse, G., Helm, R., and Kohlhaupt, R., Konstitution und Reaktivität der dimeren Perchlorbutenine, *Chem. Ber.*, 104, 2378, 1971.

R&S 026865

47. Cava, M. P. and Mitchell, J. M., *Cyclobutadiene and Related Compounds*, Academic Press, New York, 1967, 157.
48. Mano, K., Kusuda, K., and Fujino, A., Dimers of perchloro-(3,4-dimethylenecyclobutene). II. Perchloro-4,3,4,7,8-tetramethylenecyclooctadiene, *Tetrahedron Lett.*, p. 1375, 1968.
49. Doering, W. V. E. and Wiley, D. W., Heptafulvene (Methylenecycloheptatriene), *Tetrahedron*, 11, 183, 1960.
50. Roedig, A., Forsch, M., Haveaux, B., and Scheutzw, D., Valenzisomerisierung von Perchlor-4-methylenbicyclo(3.2.0)-heptadien-(2.6) zum Perchlorheptafulven, *Tetrahedron Lett.*, p. 2613, 1972.
51. Bonse, G. and Urban, Th., Oxidative metabolism of chlorinated ethylenes, *Naunyn Schmiedebergs Arch. Pharmacol.*, 287, R101, 1975.
52. Bonse, G., Urban, Th., Reichert, D., and Henschler, D., Chemical reactivity, metabolic oxirane formation and biological reactivity of chlorinated ethylenes in the isolated perfused rat liver preparation, *Biochem. Pharmacol.*, 24, 1829, 1975.
53. Daniel, J. W., The metabolism of ^{34}Cl -labelled trichloroethylene and tetrachloroethylene in the rat, *Biochem. Pharmacol.*, 12, 795, 1963.
54. Leibman, K. C., Metabolism of "TRI" in liver microsomes. I. Characteristics of the reaction, *Mol. Pharmacol.*, 1, 239, 1965.
55. Byington, K. H. and Leibman, K. C., Metabolism of trichloroethylene (Tri) in liver microsomes. II. Identification of the reaction products as chloral hydrate, *Mol. Pharmacol.*, 1, 247, 1965.
56. Leibman, K. C. and Allister, W. J., Metabolism of trichloroethylene in liver microsomes. III. Induction of the enzymatic activity and its effect on excretion of metabolites, *J. Pharmacol.*, 157, 574, 1967.
57. Frankel, D. M., Johnson, C. E., and Pitt, H. M., Preparation and properties of tetrachloroethylene oxide, *J. Org. Chem.*, 22, 1119, 1957.
58. Ikeda, M. and Imamura, T., Biological half-life of trichloroethylene and tetrachloroethylene in human subjects, *Int. Arch. Arbeitsmed.*, 31, 209, 1973.
59. Cole, W. J., Mitchell, H. G., and Salamonsen, R. F., Isolation, characterization and quantitation of chloral hydrate as a transient metabolite of trichloroethylene in man using electron capture gas chromatography and mass fragmentography, *J. Pharm. Pharmacol.*, 27, 167, 1975.
60. Butler, T. C., Metabolic transformation of Tri, *J. Pharmacol. Exp. Ther.*, 97, 84, 1949.
61. Uehleke, H., Poplawski, S., Bonse, G., and Henschler, D., Spectral evidence for 2,2,3-trichlorooxirane formation during microsomal trichloroethylene oxidation, *Naunyn Schmiedebergs Arch. Pharmacol.*, 293(Suppl. R), 253, 1976.
62. Derkosch, J., personal communication, 1974; Poluektov, V. A. and Mekhryushev, Yu. Ya., Radiation-chemical oxidation of trichloroethylene, *Zh. Fiz. Khim.*, 45, 2335, 1971.
63. Osterreicher, F., Ph.D. thesis, University of Wien, 1967; Mekhryushev, Yu. Ya. and Poluektov, V. A., Kinetics of the thermal isomerisation of trichloroepoxyethane, *Russ. J. Phys. Chem.*, 47, 959, 1973.
64. Leibman, K. C. and Ortiz, E., Microsomal metabolism of chlorinated ethylenes, 6th Int. Congr. Pharmacol., Helsinki, Abstr. 257, Nr. 608, 1975.
65. McDonald, R. N. and Schwab, P. A., Molecular rearrangements. II. Chlorine migration in the epoxide-carbonyl rearrangement, *J. Am. Chem. Soc.*, 85, 4004, 1963.
66. Griesbaum, G., Kibar, R., and Pfeffer, B., Synthese und Stabilität von 2,3-Dichloroxiranen, *Liebigs Ann. Chem.*, p. 214, February 1975.
67. McDonald, R. N. and Steppel, R. N., Molecular rearrangements. XI. The synthesis and neat, thermal rearrangement of (+)-(1R,3R)-2-chloronorborene oxo-oxide, *J. Am. Chem. Soc.*, 92, 5664, 1970.
68. Nagendrappa, G. and Griesbaum, K., Rearrangement of 2,3-dichloro-2-norborene epoxide, *Chem. Ind.*, p. 902, 1973.
69. Kobrich, G., Grosser, J., and Werner, W., Zur Umlagerung von Li-Alkoxiden aus Dichlormethylithium und Carbonylverbindungen in α -Chloroxirane und α -Chloraldehyde, *Chem. Ber.*, 105, 2610, 1973.
70. Gaertner, G. W. and Ramey, D. E., Verfahren zur Oxydation von Trichloräthylen zu Dichloräthylchlorid, Pat. Shell Int. Research Maatschappij N.V. Den Haag (Niederlande), Schweiz. Patent 464879, 1968.
71. Poluektov, V. A. and Dobrov, I. V., Reaction of trichloroethylene oxide with alcohols, *Zh. Prikl. Khim.*, 44, 2759, 1971.
72. Reichert, D. and Bashti, N., Metabolism and disposition of 1,1-dichloroethylene in the isolated blood-perfused liver of the rat, *Naunyn Schmiedebergs Arch. Pharmacol.*, 293(Suppl. R), 255, 1976.
73. Bonse, G. and Henschler, D., unpublished results.
74. Heiner, R. E., Jr., Watanabe, P. G., and Gehring, P. J., Preliminary studies of the fate of inhaled vinyl chloride monomer (VCM) in rats, *Ann. N.Y. Acad. Sci.*, 246, 135, 1975.
75. Radwan, Z. and Henschler, D., Uptake and metabolism of vinyl chloride in the isolated perfused rat liver preparation, *Naunyn Schmiedebergs Arch. Pharmacol.*, 287, R100, 1975.
76. Green, T. and Hathway, D. E., The biological fate in rats of vinyl chloride in relation to its oncogenicity, *Chem. Biol. Interact.*, 11, 545, 1975.
77. Gross, H. and Freiberg, J., Zur Existenz von Chloräthylenoxid, *J. Prakt. Chem.*, 311, 506, 1969.
78. Zief, M. and Schramm, C. H., Chloroethylene oxide, *Chem. Ind.*, p. 660, 1964.
79. Kappus, H., B. chloride to mal...
80. Bolt, H. M., K. 153, 1976.
81. Watanabe, P. G. in rats, *Toxicol.*
82. Müller, G. and 1975.
83. Yllner, S., Meta...
84. Rannug, U., Jo... activation, *Am.*
85. Bartsch, H., Me... in *S. typhimuri*
86. Greim, H., Bo... carcinogenicity 2013, 1975.
87. Bartsch, H., Ma... 2-chlorobutadie...
88. Henschler, D., rules, *Proc. 3rd*
89. Oesch, F., Kaub... substrates and in... Report of a Wo... Report 75/001,
90. Maltoni, C. and *Environ. Res.*, 7,
91. Department of H...
92. Viola, P. L. et al.
93. Roedig, A., Mar... 1963.
94. Roedig, A., Bisc... 3-II-Pentachlorp...
95. Roedig, A., Detz...
96. Roedig, A. and 749, 153, 1971.
97. Roedig, A., Detz... 60, 1971.
98. Roedig, A. and Dimeren, *Chem.*
99. Ott, E., Über d... handhabenden M...
100. Roedig, A. and K...
101. Henschler, D., Br... Umgang mit Viny...
102. Reichert, D., Ew... *Toxicol.*, 13, 511.
103. Reichert, D., Br... *Naunyn Schmied...*
104. Siegel, J., Jones, continuous inhala...

- New York, 1967, 157.
ne). H. Perchloro-(3,4,7,8-
ron, 11, 183, 1960.
on Perchloro-4-methylenbi-
von Schmiedeberg's Arch.
ue oxirane formation and
Biochem. Pharmacol., 24,
ene in the rat, Biochem.
ction, Mol. Pharmacol., 1,
somes. II. Identification of
mes. III. Induction of the
proethylene oxide, J. Org.
ene in human subjects, Int.
ution of chloral hydrate as
chromatography and mass
dichlorooxirane formation
col., 293(Suppl. R), 253,
Ya., Radiation-chemical
V. A., Kinetics of the
Congr. Pharmacol.,
in the epoxide-carbonyl
n. Liebigs Ann. Chem., p.
at, thermal rearrangement
ade, Chem. Ind., p. 902,
chloromethylithium und
3.
oracetylchlorid, Pat Shell
Prikl. Khim., 44, 2759,
ated blood-perfused liver
of inhaled vinyl chloride
ated perfused rat liver
its oncogenicity, Chem.
-69.
79. Kappus, H., Bolt, H. M., Buchter, A., and Bolt, W., Rat liver microsomes catalyze covalent binding of ^{14}C -vin chloride to macromolecules, *Nature*, 257, 134, 1975.
 80. Bolt, H. M., Kappus, H., Buchter, A., and Bolt, W., Disposition of 1,2- ^{14}C vinyl chloride in rat, *Arch. Toxicol.*, 3, 153, 1976.
 81. Watanabe, P. G., McGowan, G. R., and Gehring, P. J., Fate of [^{14}C] vinyl chloride after single oral administration in rats, *Toxicol. Appl. Pharmacol.*, 36, 339, 1976.
 82. Müller, G. and Norpoth, K., Bestimmung zweier Urinmetabolite des Vinylchlorids, *Naturwissenschaften*, 62, 541, 1975.
 83. Yllner, S., Metabolism of chloroacetate-1- ^{14}C in the mouse, *Acta Pharm. Toxicol.*, 30, 69, 1971.
 84. Rannug, U., Johansson, A., Ramel, C., and Wachtmeister, C. A., The mutagenicity of vinyl chloride after metabolic activation, *Ambio*, 3, 194, 1974.
 85. Bartsch, H., Malaveille, C., and Montesano, R., Human, rat and mouse liver-mediated mutagenicity of vinyl chloride in *S. typhimurium* strains, *Int. J. Cancer*, 15, 429, 1975.
 86. Greim, H., Bonse, G., Radwan, Z., Reichert, D., and Henschler, D., Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation, *Biochem. Pharmacol.*, 24, 2013, 1975.
 87. Bartsch, H., Malaveille, C., Montesano, R., and Tomatis, L., Tissue-mediated mutagenicity of vinylidene chloride and 2-chlorobutadiene in *Salmonella typhimurium*, *Nature*, 255, 641, 1975.
 88. Henschler, D., Bonse, G., and Greim, H., Carcinogenic potential of chlorinated ethylenes - tentative molecular rules, Proc. 3rd WHO-I.A.R.C. Meet., Lyon, November 3, 1975.
 89. Oesch, F., Kaubisch, N., Jerina, D. M., and Daly, J. W., Hepatic epoxide hydraz. Structure-activity relationships for substrates and inhibitors, *Biochemistry*, 10, 4858, 1971.
 90. Report of a Working Group on Epidemiological Studies on Vinyl Chloride Exposed People, I.A.R.C. Intern. Tech. Report 75/001, Lyon, 1975.
 91. Maltoni, C. and Lefemine, G., Carcinogenicity bioassays of vinyl chloride. I. Research plan and early results, *Environ. Res.*, 7, 387, 1974.
 92. Department of Health, Education and Welfare, Washington, D.C., Memorandum, March 20, 1975.
 93. Viola, P. L. et al., cited in Reference 87.
 94. Roedig, A., Markl, G., and Heinrich, B., Perchlorpropadien-(1,2), ein hochreaktives Allen, *Angew. Chem.*, 75, 88, 1963.
 95. Roedig, A., Bischoff, F., Heinrich, B., and Markl, G., Perchlorallen als Zwischenstufe der Dehydrochlorierung von 3-H-Pentachlorpropen-(1) zu Perchlor-1,2-dimethylen-cyclobutan, *Liebigs Ann. Chem.*, 670, 8, 1963.
 96. Roedig, A., Detzer, N., and Friedrich, H. J., Tribromallen und Perbromallen, *Angew. Chem.*, 76, 379, 1964.
 97. Roedig, A. and Detzer, N., Thermisch instabile Allene. VIII. Trichlor-athoxycarbonyl-allen, *Liebigs Ann. Chem.*, 749, 153, 1971.
 98. Roedig, A., Detzer, N., and Bonse, G., Thermisch instabile Allene IX. Trichlorbromallen, *Liebigs Ann. Chem.*, 752, 60, 1971.
 99. Roedig, A. and Heinrich, B., Thermisch instabile Allene. IV. Einige Reaktionen des Perchlorallens und seines Dimeren, *Chem. Ber.*, 100, 3716, 1967.
 100. Ott, E., Über das Dichloracetylen. III. Mitteil: Darstellung und einige Vorlesungsversuche mit der gefahrlos zu handhabenden Molekülverbindung mit Ather, *Chem. Ber.*, 75, 1517, 1942.
 101. Roedig, A. and Kohlaupt, R., Dimerisierung von Perchlorbutenn, *Tetrahedron Lett.*, p. 1107, 1964.
 102. Henschler, D., Bräser, F., and Hopf, C. H., Polyneuritis cranialis durch Vergiftung mit chlorierten Acetylenen beim Umgang mit Vinylidenchlorid-Copolymeren, *Arch. Toxicol.*, 26, 62, 1970.
 103. Reichert, D., Ewald, D., and Henschler, D., Generation and inhalation toxicity of dichloroacetylen, *Food Cosmet. Toxicol.*, 13, 511, 1975.
 104. Reichert, D., Brinke, G., Bannasch, P., and Liebalde, G., Neuro- and nephrotoxic effects of dichloroacetylene, *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 274(Suppl.), R90, 1972.
 105. Siegel, J., Jones, R. A., Coon, R. A., and Lyon, J. P., Effects on experimental animals of acute, repeated and continuous inhalation exposures to dichloroacetylene mixtures, *Toxicol. Appl. Pharmacol.*, 18, 168, 1971.