

Comparative Toxicity Study of 2,4,5,2',4',5'-Hexachlorobiphenyl and a Polychlorinated Biphenyl Mixture in Rabbits

J. G. VOS AND ERICA NOTENBOOM-RAM

*Institute of Veterinary Pathology and Institute of Veterinary Pharmacology and Toxicology,
University of Utrecht, Bilthoven 172, Utrecht, The Netherlands*

Received December 6, 1971

Comparative Toxicity Study of 2,4,5,2',4',5'-Hexachlorobiphenyl and a Polychlorinated Biphenyl Mixture in Rabbits. Vos, J. G. and NOTENBOOM-RAM, ERICA (1972). *Toxicol. Appl. Pharmacol.* 23, 563-578. Three groups of rabbits were treated daily (5 times per week for 28 days) with 120 mg 2,4,5,2',4',5'-hexachlorobiphenyl, 120 mg polychlorinated biphenyl (PCB) mixture (Aroclor 1260[®]) and the solvent isopropanol, respectively. The dermal application resulted in early macroscopic skin lesions in the Aroclor group. The lesions in the 2,4,5,2',4',5'-hexachlorobiphenyl group appeared later on and were less severe. This difference was confirmed microscopically: hyperplasia and hyperkeratosis of the follicular and epidermal epithelium were more severe in the Aroclor[®] group. Fecal coproporphyrin levels were significantly increased in the experimental groups. Enhanced liver weights were found in both test groups. Liver injury, as judged by light microscopic lesions and elevated serum transaminase levels, was somewhat more severe in the hexachlorobiphenyl group when compared with the Aroclor[®] group, though the mean liver content was about the same (respectively, 239 and 236 ppm). Light microscopic findings included subcapsular necrosis, zonal necrosis, hydropic degeneration, as well as a peripheral and perinuclear shift of cell organelles, and focal cytoplasmic hyalin degeneration. In electron microscopy the shift was found to be due to a proliferation of smooth surfaced membranes of the endoplasmic reticulum (SER), resulting in a displacement of rough surfaced membranes (RER) and mitochondria. The hyalinized cytoplasm was recognized as tightly packed tubules of proliferated SER, that is considered as hypertrophic, hypoactive SER. The contribution of chlorinated dibenzofurans and pure PCB in the toxicity of crude PCB mixtures is discussed.

Chlorinated biphenyls (PCB) are industrial chemicals with many applications. Toxicity of these compounds for industrial workers has been known for years, especially the occupational disease known as chloracne (Jones and Alden, 1936; Jaritz, 1936; Meigs *et al.*, 1954). Recently, the consumption of rice bran oil accidentally contaminated by PCB resulted in many cases of chloracne among Japanese farmers (quoted by Crow, 1970); liver damage was also reported (quoted by Izumi, 1970). A critical report on the use, environmental contamination and the toxicity of PCB and other industrial halogenated hydrocarbons appeared recently

© 1972 by Academic Press, Inc.
All rights reserved.

563

Toxicol. Appl. Pharmacol.
v. 23 (1972)

NPC00026557

753894

(Zitko and Choi, 1971). There is an additional danger for wildlife from the widespread environmental contamination with PCB through food chains, and this is, ultimately, also a menace to human health.

The purpose of the present study is to compare the toxicity of Aroclor 1260[®] and a single isomer 2,4,5,2',4',5'-hexachlorobiphenyl. The commercial Aroclor sample was found to be the least toxic in a comparative study using three PCB mixtures originating from different firms (Vos and Koeman, 1970). Chlorinated dibenzofurans, considered to be responsible for the higher toxicity of two PCB mixtures, were found in the Aroclor 1260[®] sample (Vos *et al.*, 1970). To be more certain about the biological effect attributable to PCB itself, a pure compound, 2,4,5,2',4',5'-hexachlorobiphenyl, was used for comparison. This is also one step in the understanding of the toxicity of the different components of PCB mixtures.

Hepatic porphyria was also studied. Special attention was paid to PCB-induced liver damage, using light and electron microscopy for structural alterations and chromatography for the determination of liver residues.

METHODS

The technical PCB mixture, a viscous fluid with an average of 60% chlorine, was obtained from Monsanto in the United States (Aroclor 1260[®] Lot No. AB). 2,4,5,2',4',5'-Hexachlorobiphenyl, a crystalline compound, was synthesized according to the Ullman synthesis, using 2,4,5-trichloriodobenzene and copper powder at 225°C (Tas and de Vos, 1971). Because of the chemical reaction, it is unlikely that this particular isomer contains chlorinated dibenzofurans as impurity. The Aroclor[®] sample was analyzed for the presence of chlorinated dibenzofuran and found to be free (limit of detection: 1 ppm).

Adult female New Zealand rabbits, 3.5 mo old and weighing 2.5-2.9 kg, were distributed at random into 3 groups of 4 animals. The performance of the toxicity study was in general the same as described earlier (Vos and Beems, 1971). Because of the lower solubility of the hexachlorobiphenyl isomer when compared with the PCB mixture, 20 mg of the PCBs (instead of 120 mg in the prior study) were dissolved per milliliter of isopropanol. The PCB solutions were kept at body temperature in order to keep the PCB dissolved. Six milliliters of PCB solution (120 mg) or of isopropanol as a control was dropped daily, 5 times a week, on an area of 5 × 10 cm on the clipped and shaved backs. After a test period of 4 wk (20 applications of 120 mg PCB), the animals were killed.

The coproporphyrin and protoporphyrin analyses of feces from the cecum, a macroscopical examination of tissues in Wood's light as parameters for chlorinated porphyria, and the hematological examination were done according to a former study (Vos and Beems, 1971). At necropsy, body and organ weights were determined. The skin, liver and kidney were fixed in 10% buffered formalin. The spleen, thymus, aortic and mesenteric lymph node and appendix were fixed in Carnoy's fluid for 24 hr at 4°C and stained with methylgreen-pyronin according to Elias (1969). Paraplast[®] embedded sections were stained with hematoxylin and eosin. For detailed histology, whole sections were stained with Perls' iron stain and with Best's carmine. Moreover, cryostat sections of the livers were stained with Sudan black for the detection of lipids.

NPC00026558

753895

Paraplast[®] embedded sections and unstained cryostat sections of liver, kidney and intestine were studied in a fluorescence microscope for the detection of pigment and excess quantities of porphyrins, respectively.

For ultrastructural examination, small blocks of liver tissue were fixed in 0.1 M phosphate-HCl buffered 5% glutaraldehyde (pH 7.0) for 5 hr. The liver tissue was washed two times for 45 min in 0.1 M phosphate buffered 2% osmium tetroxide

TABLE 1

COPROPORPHYRIN AND PROTOPORPHYRIN CONTENTS ($\mu\text{g/g}$ DRY WEIGHT) OF FECES OF RABBITS TREATED WITH PCB FOR FOUR WEEKS*

Coproporphyrin			Protoporphyrin		
Hexachlorobiphenyl	Aroclor [®]	Control	Hexachlorobiphenyl	Aroclor [®]	Control
45.0	24.1	4.5	88.3	35.1	17.8
6.2	5.0	3.7	8.3	8.4	11.9
20.8	29.4	3.6	51.2	65.1	15.6
39.6	16.0	3.5	66.5	65.1	10.9
27.9 ^b	18.6 ^b	3.8	53.6	43.4	14.0

*Values are the contents of feces, collected from the cecum of the individual animals.
^bSignificantly different from controls, $p < 0.025$.

TABLE 2

ABSOLUTE AND RELATIVE LIVER WEIGHTS, MICROSCOPIC LIVER DAMAGE, LIVER RESIDUES AND TRANSAMINASE VALUES FOR RABBITS TREATED WITH PCB FOR FOUR WEEKS*

	Liver weight		Microscopic Liver damage ^b	Liver Content (ppm)	SGPT (U)	SGOT (U)
	g	g/100 g body weight				
Hexachlorobiphenyl	128	4.6	++	249	71	16
	132	4.2	++	140	56	34
	146	4.8	+++	261	70	30
	126	5.1	+++	104	92	21
	Mean: 133	4.7 ^c		239	72 ^d	25 ^d
Aroclor [®]	141	5.0	++	198	64	29
	129	4.0	++	248	32	10
	152	4.8	++	317	53	13
	133	4.2	+	182	34	12
	Mean: 139 ^c	4.5		236	46	16
Control	131	4.3	—	0.8	43	9
	110	3.5	—	0.6	22	10
	91	2.9	—	0.6	20	11
	83	2.8	—	0.7	31	11
	Mean: 104	3.4		0.7	29	10

*Values are given for the individual animals.

^b++ to +++ is a comparative estimate of the liver damage.

^cSignificantly different from controls, $p < 0.05$.

^dSignificantly different from controls, $p < 0.025$.

NPC00026559

753896




FIG. 1. Liver of a 2,4,5,2',4',5'-hexachlorobiphenyl treated rabbit. Subcapsular band of connective tissue (1) and area of hydropic degeneration (2). Hematoxylin and eosin. $\times 60$.




FIG. 2. Higher magnification of the subcapsular region of the liver seen in Fig. 1. Note the mesenchymal cells at 1, hydropic cells at 2 and proliferation of mesenchymal cells at 3. Hematoxylin and eosin. $\times 120$.

NPC00026560

753897




Fig. 3. Centrilobular area of an Arnelor[®] treated rabbit liver. A marked peripheral and perinuclear concentration of basophilic organelles can be seen. Hematoxylin and eosin, $\times 375$.




Fig. 4. Hepatocytes of a control animal. Clear areas (arrow) represent negative images of glycogen. Basophilic material representing cell organelles is scattered in the cytoplasm. Toluidine blue, $\times 940$.

NPC00026561




FIG. 5. Liver section of an Aroclor[®] treated rabbit. Toluidine blue. $\times 940$. (a) Showing perinuclear displacement of cell organelles; note also the foamy cytoplasm (arrow). (b) Hydropic cells showing vacuoles (1), that are separated from homogeneous hyalinized cytoplasm (2) here perivacuolar localization of densely staining organelles.




FIG. 6. Severely damaged hepatocytes of the same 2,4,5,2',4',5'-hexachlorobiphenyl-treated rabbit as shown in Figs. 1 and 2 showing hydropic changes (1), lipid droplets (2), and focal cytoplasmic hyalin degeneration (3). Note also the nuclear alterations such as the irregular outlines of the nuclei and the prominent nucleoli. Toluidine blue. $\times 940$.

NPC00026562

753899

72). Block staining was carried out with uranyl acetate. The tissue was then dehydrated rapidly in ethanol and embedded in Dow epoxy resin (DER^A) (Lockwood, 1964). Sections were cut with glass knives on a Reichert ultramicrotome. Thin (0.5–1.0 μ) sections for orientation and for light microscopic examination were stained with toluidine blue. Ultrathin sections poststained with lead citrate were examined by transmission microscopy (Philips EM 100^A) at 40 kV.

PCB concentrations in the livers were determined by gas chromatography, as reported earlier (Vos, *et al.*, 1971). The significance of difference between treated and control groups was determined on a one-tail significance level, using the Wilcoxon test for two unrelated samples (van der Waerden, 1957).

RESULTS

After 3 days of PCB treatment, the skin of rabbits in the Aroclor[®] group showed redness. After 6 days, definite redness and some thickening of the skin was noted in this group. After 2 wk of treatment, hyperkeratosis and formation of transverse scales were seen. At this time, some redness and slight hyperkeratosis, together with presence of hexachlorobiphenyl crystals on the skin, was seen in the 2,4,5,2',4',5'-hexachlorobiphenyl treated group, although a slight hyperkeratosis was present in the




Fig. 7. Control liver with hepatocytes showing diffusely scattered mitochondria and network of the endoplasmic reticulum of rough and smooth types. The clear areas may represent negative images of glycogen. The intercellular space between the bile canaliculi may be an artifact. Nucleus (N); bile canaliculus (BC); space of Disse (D); endothelial cell (E); and sinusoid (S). Uranyl acetate and lead citrate. $\times 4300$.

NPC00026563

753900

control (isopropanol) group. Differences in the regrowth of hair between the two groups were also noted. Regrowth was most reduced in the Aroclor[®] group as in the hexachlorobiphenyl treated animals. During the course of the experiment lesions in the Aroclor group were clearly progressive, while the lesions in the chlorobiphenyl group increased rather slowly.

All animals, except 1 rabbit in the hexachlorobiphenyl group, showed a weight gain during the experimental period of 4 wk. Mean weight gain in the hexachlorobiphenyl group was 142 g (ranging from - 215 to 365 g), in the Aroclor[®] group 325 g (70 to 640) and in the control group 491 g (from 430 to 550).



FIG. 8. Hepatocytes of the same liver of an Aroclor[®] treated rabbit as shown in Figs. 1 and 5a, the foamy cytoplasm. The cytoplasm is mostly occupied by smooth endoplasmic reticulum (SER). The perinuclear shift of mitochondria and RER can also be seen. N, nucleus. Uranyl acetate and citrate. $\times 4300$.

Coproporphyrin levels in the feces, collected at necropsy from the cecum of individual animals, were significantly increased in the hexachlorobiphenyl as well as in the Aroclor[®] treated rabbits (Table I). Protoporphyrin excretion was also increased, though not significantly. Macroscopically, fluorescence was found in the bile of 2 rabbits of the experimental groups. One animal of the Aroclor[®] group showed also fluorescence in the liver and bile. Examination of cryostat sections of liver, kidney, and small intestine with the fluorescence microscope showed fluorescence in the livers of 7 out of 8 PCB treated animals. This fluorescence was mostly located in the bile duct and sometimes diffuse.

NPC00026564

753901

periporally located hepatocytes. Some fluorescence was also noted in epithelial cells of the small intestine in 4 PCB treated animals. Severe abdominal edema was seen in an Aroclor[®] treated rabbit that showed also the highest liver content (317 ppm) and severe liver damage (Table 2).




Fig. 9. Higher magnification of a part of Fig. 8. The proliferation of SER consists of vesicular and tubular membranes. Clear areas of the cytoplasm, associated with the SER, most likely represent glycogen (arrow). Note the dilatation and degranulation of the rough endoplasmic reticulum (RER). M, mitochondria; N, nucleus. Uranyl acetate and lead citrate. $\times 13,300$.

No significant differences were found in absolute and relative organ weights, except in the liver, and in hematological findings, except for the serum transaminase levels. These values, together with the PCB concentrations in the livers and the comparative estimate of liver damage are given in Table 2. The SGPT and SGOT values were significantly increased in the hexachlorobiphenyl group. An increase was also noted in the Aroclor[®] group. Mean liver content of PCB was the same in the experimental

NPC00026565

753902

groups. The increase and severity of liver damage in the Aroclor* group as the increase in liver residue.

At examination of skin sections of the Aroclor* group, hyperplasia and keratosis of the epidermal and follicular epithelium were found. The lesions were yet so pronounced as in the prior study (Vos and Beems, 1971). Hexachlorocyclopentadiene induced skin damage, especially hyperplasia in the epidermal and follicular epithelium, as well as follicular plugging, was clearly less when compared with the Aroclor* treated rabbits. A slight hyperkeratosis was found in some control animals. To be due to the treatment with isopropanol (6 ml/day); it was not found in the prior study (Vos and Beems, 1971) using a lower dose (1 ml/day).



FIG. 10. Hepatocytes of an Aroclor* treated animal. Similar hydropic cells as seen in Fig. 9. Large confluent vacuoles with flocculent material within them (V). The hyalinized cytoplasm of hepatocytes shown in Fig. 5b consists of hypertrophied, densely packed SER. Note the arrangement of mitochondria around the vacuoles (arrow). Uranyl acetate and lead citrate. $\times 5500$.

When the lymphoid system was examined, no outstanding differences were found in the spleen, appendix and lymph nodes between control and treated groups. Some atrophy of the cortex, together with less pyroninophilic cells in the marrow, were found in the thymuses of rabbits in the PCB treated groups. The atrophy agrees with the absolute and relative (about 18%) thymus weights in these groups. No kidney damage was found, except for the presence of polyploid tubular epithelial cells in a liver of a biphenyl treated rabbit.

Pathologic changes in the livers of the treated animals were, in general, the same as described in a previous study (Vos and Beems, 1971). These findings included centrilobular degeneration and liver cell atrophy, focal hyalin degeneration of the cytoplasm.

NPC00026566

753903

of hepatocytes, enlarged nuclei and loss of glycogen. As shown in Table 2, liver damage by hexachlorobiphenyl was somewhat more severe when compared with the Aroclor[®] induced lesions. However, in the present study, areas of hydropic, degenerated and necrotic cells dominated, often in the subcapsular region (Figs. 1 and 2), forming bands



Fig. 11. Higher magnification of a part of Fig. 10 reveals the tubular structure of the densely hyperplastic smooth endoplasmic reticulum. Uranyl acetate and lead citrate. $\times 13,100$.

necrotic and degenerated cells, accompanied sometimes by proliferation of mesenchymal cells. Necrosis was usually accompanied by hemorrhage. Loss of glycogen was not limited to the degenerated, hydropic or necrotic areas, but was also seen in foci showing no alterations in hematoxylin-eosin sections. In the rest of the liver the glycogen content was also reduced when compared with the controls. The centrilobular damage was often zonal. This distribution can be explained by the concept of structural and functional hepatic unit (Rappaport *et al.*, 1954). The presence of Perl's positive material

NPC00026567

753904

and ceroid pigment in Kupffer and parenchymal cells (characterized by a brownish fluorescence) was evidently less than in the preceding study and was limited to some necrotic areas.

In both treated groups, numerous hepatocytes were seen with light staining cytoplasm. In these cells, the cell membrane and the nuclear membrane were empty by the presence of basophilic material (Fig. 3). When plastic embedded liver was examined by light microscopy, these alterations could be observed more clearly (Figs. 4 and 5a). This was the general trend in the PCB treated animals. In electron microscopy, the most striking feature of these cells was the proliferation of



FIG. 12. Severely damaged hepatocyte of the same 2,4,5,2',4',5'-hexachlorobiphenyl treated animal as shown in Fig. 6. Photograph shows a cluster of tubular smooth endoplasmic reticulum (microscopically visible as focal hyaline degeneration), forming a dense, closely packed aggregate (SER). Note also the lipid droplets (L), and hydropic swelling, as shown by a great number of small and large vesicles (V). Uranyl acetate and lead citrate. $\times 3000$.

surfaced membranes of the endoplasmic reticulum (SER). As a result of the proliferation of SER, a perinuclear and peripheral displacement of mitochondria and surfaced membranes of the endoplasmic reticulum (RER) was apparent (Figs. 7a). The proliferation of SER could well explain the accentuated linings of cell and nuclei in light microscopy and the increase in liver weight (Table 2) in the PCB treated group. The RER showed some dilatation and degranulation (Fig. 9).

In addition to the proliferative changes of the SER, degenerative changes were noted. Light microscopy showed the presence of hydropic cells (Fig. 2). Examination of plastic-embedded material under the light microscope showed some hydropic cells, which consisted of large vacuoles separated from a homogeneous cytoplasm,

NPC00026568

753905

as focal cytoplasmic degeneration. At the rim of the homogeneous cytoplasmic vacuoles were visible (Fig. 5b). The focal hyalinized cytoplasm (Fig. 5b) was shown by electron microscopy to consist of tightly packed tubules of proliferated SER (Figs. 10 and 11). The condensed SER was separated from large vacuoles, sometimes containing flocculent material in it. The mitochondria were arranged around the vacuoles. In the hyalinized cytoplasm, lipid droplets and degenerative changes of the nucleus were found in the more severely affected areas (Figs. 6 and 12). The nuclear changes included irregular outlines of the nuclei and karyopycnosis. Myelin figures were found rarely in the cytoplasm. In the present study, no differences were noted in the hepatotoxic action between the polychlorinated biphenyl mixture and 2,4,5,2',4',5'-hexachlorobiphenyl.

DISCUSSION

From the observed acnelike lesions, both from the PCB mixture and 2,4,5,2',4',5'-hexachlorobiphenyl, and assuming that the hexachlorobiphenyl is free from contamination with chlorinated dibenzofuran, it can be concluded that this particular compound in the mixture PCB has a slight acnegenic action of itself. Considering the already mentioned dermal toxicity study, it is evident that the major acnegenic action of crude PCB mixtures comes from chlorinated dibenzofurans. In Table 1 it is shown that 2,4,5,2',4',5'-hexachlorobiphenyl is even more porphyrogenic than the PCB mixture, so it is probable that hepatic porphyria comes only from PCB itself.

In contrast with the prior dermal toxicity study, only slight effects were noted from PCB on the lymphatic tissue. This difference is explained by the longer experimental period in the former study (38 days) which resulted in a less healthy condition of the experimental animals. Stress (e.g., release of glucocorticoids) must be considered responsible for the major effects on the lymphatic tissue. However, using the sensitive fluorescent antibody technique in tetanus toxoid stimulated guinea pigs, immunosuppression was found (Vos and de Ruij, 1972).

Liver damage was essentially the same after treatment with both the Aroclor[®] mixture and 2,4,5,2',4',5'-hexachlorobiphenyl. Liver damage caused by commercial PCB mixtures was due predominantly to the contaminants. This conclusion is based on the differences in liver toxicity between three PCB preparations (Vos and Koeman, 1970; Vos and Beenis, 1971). The probable contribution of chlorinated dibenzofurans and uncontaminated PCB in the toxicity of crude PCB mixtures is summarized in Table 3.

TABLE 3

PROBABLE CONTRIBUTION OF CHLORINATED DIBENZOFURAN AND PURE POLYCHLORINATED BIPHENYL IN THE TOXICITY OF TECHNICAL PCB PREPARATIONS

	Chloracne	Edema formation	Liver damage	Hepatic porphyria
Chlorinated dibenzofuran	++	+	+	+
Polychlorinated biphenyl	-	-	+	++

NPC00026569

753906

Focal cytoplasmic hyalin degeneration, as well as ceroid pigment deposits in hepatic cells, was more pronounced in the former dermal toxicity study (Vos and Beems, 1971), probably due to the longer experimental period. Hyalin inclusions were previously described after semichronic exposure to PCB in rat liver (Bennett *et al.*, 1938; Miller, 1944) and in mouse liver (Nishizumi, 1970). However, in the rabbit, most important liver lesions described varied from fatty degeneration to massive hepatocytic degeneration and necrosis (von Wedel *et al.*, 1943; Miller, 1944). In this study, the presence of cells with proliferated SER, resulting in a perinuclear and peripheral shift of RER and mitochondria, was more frequently seen (Fig. 8). This proliferation, also found by Nishizumi (1970) in mouse and monkey liver, must be considered the structural indication for enhanced metabolism of foreign lipophilic compounds. Thus, the activity of aniline hydroxylase and aminopyrine *N*-demethylase (determined *in vitro*) was increased in the rabbit after administration of Aroclor 1248 (Villeneuve *et al.*, 1971). A decrease in sleeping time after treatment with hexobarbital and enhanced *in vitro* rates of aniline hydroxylation and *p*-nitroanisole demethylation were demonstrated by Street *et al.*, (1969). They also found an increase of these effects with increasing chlorine content of different PCB preparations. Despite this increase in drug enzyme activity caused by the higher chlorinated mixtures, the lower chlorinated compounds are very probably metabolized to a greater extent (Grant *et al.*, 1971). On the other hand, the presence of focal cytoplasmic hyalin degeneration with glycogen (Figs. 5h and 6), in electron microscopy seen as densely packed agglomerations of SER, very probably represents hypertrophic, hypoactive SER. Hutterer *et al.* (1968) found reduced activity of microsomal oxidases (e.g. drug enzymes), while SER remained hypertrophic in livers of rats treated with high doses of dieldrin. As seen under the electron microscope, hypertrophic, hypoactive SER was recognized as tight clusters of tubular membranes, which could indicate a transition from adaptation to injury (Hutterer *et al.*, 1968, 1969). Liver damage was also suggested by elevated transaminase levels (Table 2). Myelin figures were seldom found in this study. Perhaps this depends on the animal species, since Nishizumi (1970) described myelin figures in mouse liver after PCB feeding, but not in monkey liver. The occurrence of SER proliferation and concentric membrane arrays is found in the livers of rats fed a chlorinated biphenyl (Norback and Allen, 1970) and toxic fat containing hexachlorodibenzo-*p*-dioxin, a compound related to chlorodibenzofuran (Norback and Allen, 1969). The results from the former study suggested that the concentric arrays of agranular membranes may be structural modifications of the endoplasmic reticulum having an enzymatic function similar to that associated with the SER.

ACKNOWLEDGMENTS

We are grateful to Dr. R. J. C. Kleipool and Mr. A. C. Tas of the Central Institute for Food and Nutrition Research, T.N.O., Zeist, for synthesis and supply of the 2,4,5,2',4',5'-hexachlorobiphenyl sample.

Special thanks are due to Drs. C. J. A. H. V. van Vorstenbosch of the Institute of Veterinary Histology (Head: Professor Dr. W. A. de Voogd van der Straaten) for use of the electron microscope, as well as for help and advice.

NPC00026570

753907

REFERENCES

- WATT, G. A., DRINKER, C. K. and WARREN, M. F. (1938). Morphological changes in the livers of rats resulting from exposure to certain chlorinated hydrocarbons. *J. Ind. Hyg. Toxicol.* 20, 97-123.
- WATKINS, K. D. (1970). Chloracne. A critical review including a comparison of two series of cases of acne from chloronaphthalene and pitch fumes. *Trans. St. John's Hosp. Dermatol. Soc.* 56, 79-99.
- WILKINSON, J. M. (1969). Effects of temperature, poststaining rinses and ethanol-butanol dehydrating pictures on methylogreen-pyronin staining. *Stain Technol.* 44, 201-204.
- GRANT, D. L., PHILLIPS, W. E. J. and VILLENEUVE, D. C. (1971). Metabolism of a polychlorinated biphenyl (Aroclor 1254) mixture in the rat. *Bull. Environ. Contam. Toxicol.* 6, 102-112.
- WITTEKAMP, F., SCHAFFNER, F., KLION, F. M. and POPPER, H. (1968). Hypertrophic, hypoactive smooth endoplasmic reticulum: A sensitive indicator of hepatotoxicity exemplified by dieldrin. *Science* 161, 1017-1019.
- WITTEKAMP, F., KLION, F. M., WENGRAF, A., SCHAFFNER, F. and POPPER, H. (1969). Hepato-cellular adaptation and injury. Structural and biochemical changes following dieldrin and methyl butyl yellow. *Lab. Invest.* 20, 455-464.
- WILKINSON, J. W. and ALDEN, H. S. (1936). An acneform dermatogesis. *Arch. Dermatol. Syphilol.* 33, 1022-1034.
- LEWIS, W. R. (1964). A reliable and easily sectioned epoxy embedding medium. *Anat. Rec.* 150, 129-140.
- WILKINSON, J. W., ALBOM, J. J. and KARTIN, B. L. (1954). Chloracne from an unusual exposure to Aroclor. *J. Amer. Med. Ass.* 154, 1417-1418.
- WILKINSON, J. W. (1944). Pathologic changes in animals exposed to a commercial chlorinated biphenyl. *U.S. Pub. Health Rep.* 59, 1085-1093.
- YONIZUMI, M. (1970). Light and electron microscope study of chlorobiphenyl poisoning. *Arch. Environ. Health* 21, 620-632.
- YONIZUMI, M. and ALLEN, J. R. (1969). Morphogenesis of toxic fat-induced concentric membrane arrays in rat hepatocytes. *Lab. Invest.* 20, 338-346.
- YONIZUMI, M. and ALLEN, J. R. (1970). Enzymatic and morphologic alterations of hepatic endoplasmic reticulum induced by a chlorinated aromatic hydrocarbon. *Fed. Proc. Fed. Amer. Soc. Exp. Biol.* 29, 816, 3220.
- LOPAPORT, A. M., BURROWS, Z. J., LOUGHEED, W. M. and LOTTO, W. N. (1954). Subdivision of hexagonal liver lobules into a structural and functional unit. Role in hepatic physiology and pathology. *Anat. Rec.* 119, 11-34.
- SCHWARTZ, L. (1936). Dermatitis from synthetic resins and waxes. *Amer. J. Pub. Health* 26, 586-592.
- LEWIS, J. C., URRY, F. M., WAGSTAFF, D. J. and BLAU, A. D. (1969). Comparative effects of polychlorinated biphenyls and organochlorine pesticides in induction of hepatic microsomal enzymes. Presented at ACS meeting Div. Pesticides, Paper No. 17. September 8-12, New York.
- LEWIS, A. C. and DE VOS, R. H. (1971). Characterization of four major components in a technical polychlorinated biphenyl mixture. *Environ. Sci. Technol.* 5, 1216-1218.
- VAN DER WAERDEN, B. L. (1957). *Mathematische Statistik*, pp. 269-284. Springer-Verlag, Berlin and New York.
- VILLENEUVE, D. C., GRANT, D. L., PHILLIPS, W. E. J., CLARK, M. L. and CLEGG, D. J. (1971). Effects of PCB administration on microsomal enzyme activity in pregnant rabbits. *Bull. Environ. Contam. Toxicol.* 6, 120-128.
- WEDEL, H., HOLLA, W. A. and DENTON, J. (1943). Observations on the toxic effects resulting from exposure to chlorinated naphthalene and chlorinated phenyls with suggestions for prevention. *Rubber Age (New York)* 53, 419-426.
- WATKINS, J. G. and BEEMS, R. B. (1971). Dermal toxicity studies of technical polychlorinated biphenyls and fractions thereof in rabbits. *Toxicol. Appl. Pharmacol.* 19, 617-633.

NPC00026571

- Vos, J. G. and DE ROU, T. (1972). Immunosuppressive activity of a polychlorinated biphenyl preparation on the humoral immune response in guinea pigs. *Toxicol. Appl. Pharmacol.* 21, 549-555.
- Vos, J. G. and KOEMAN, J. H. (1970). Comparative toxicologic study with polychlorinated biphenyls in chickens with special reference to porphyria, edema formation, liver damage and tissue residues. *Toxicol. Appl. Pharmacol.* 17, 656-668.
- Vos, J. G., KOEMAN, J. H., VAN DER HAAS, H. L., TEN NIEVER DE BRAUW, M. C., and DE VRIES, R. H. (1970). Identification and toxicological evaluation of chlorinated dibenzofuran and chlorinated naphthalene in two commercial polychlorinated biphenyls. *Food Cosmet. Toxicol.* 8, 625-633.
- Vos, J. G., STRIK, J. J. T. W. A., VAN HOLSTEIN, C. W. M. and PENNING, J. H. (1971). Polychlorinated biphenyls as inducers of hepatic porphyria in Japanese quail, with special reference to δ -aminolevulinic acid synthetase activity, fluorescence, and residues in the liver. *Toxicol. Appl. Pharmacol.* 20, 232-240.
- Zitko, V. and CHOI, P. M. K. (1971). PCB and other industrial halogenated hydrocarbons in the environment. *Fish. Res. Bd. Can. Techn. Rep.* 272.

NPC00026572

753909