

Study finds 33 brain tumor deaths at Port Arthur, Beaumont refineries

By ERNA SMITH
Post Reporter

Thirty-three brain tumor deaths have been found among union workers at three petroleum refineries in Port Arthur and Beaumont, The National Cancer Institute has reported.

The workers, members of the Oil, Chemical and Atomic Workers International Union, were employed at Texaco and Gulf refineries in Port Arthur, and the Mobil refinery in Beaumont.

The findings confirm a NCI "hypotheses-generating" study, released in May 1979, that suggested workers in the entire petrochemical industry are exposed to increased risks of several cancers, including those affecting digestive and respiratory systems. The greatest increase emerged in the number of brain cancer cases.

"Scientists at the National Cancer Institute and the National Institute for Occupational Safety and Health have found an apparent association between increased risk of brain cancer and employment in the oil refinery industry," reads an NCI fact sheet, obtained by the Post.

However, Gulf — using a more widely recognized statistical method and examining a larger number of deaths — reported the preliminary results of a company study "shows no increased risk of employees developing or dying from brain tumors" at its Port Arthur refinery.

Furthermore, the company contends, the "undue publicity" given the NCI findings has "caused needless anxiety and concern among our employees."

Three years ago, OCAW invited NCI to examine its membership records to evaluate for the first time cancer risks among a large group of workers exposed to petroleum and petroleum products. The union, the largest labor organization for petroleum workers, has nearly 20,000 members in Texas.

The NCI and NIOSH scientists in their study confirmed that 7 of the 33 OCAW members died of cancer. In the remaining six cases, the cause of death was listed only as brain tumor.

NCI's original study population consisted of 3,105 males whose deaths were reported between 1947 and 1977 to OCAW locals in Texas. For the brain cancer report, scientists examined 2,133 deaths of active and re-

tired men who worked in the refineries from 1943-78.

Researchers examined death certificates, obtained by the union and the state's vital records division, and used that data to compute the frequency of fatal diseases among the workers.

The number of deaths was statistically compared to deaths from all causes and deaths from all types of cancer among males in the U.S. population as a whole.

Researchers reviewed 1,008 deaths at Texaco, 729 at Gulf and 396 at Mobil. Out of the 2,133 deaths examined at all three refineries, researchers found 27 brain cancer cases, compared to the normally expected 15 fatalities from all forms of cancer.

At Texaco, scientists observed 16 fatal brain tumor cases, compared to an expected 7.5 fatalities from all causes of death. Ten brain tumor fatalities were observed among Gulf union members compared to an expected 7.5, and 7 cases were uncovered at Mobil compared to an expected 3.

Terry Thomas, NCI's chief project researcher, said the institute's findings are "preliminary" and more

Please see 33 deaths/page 19A

STG 2433932

RSV0034013

From page 1

study is needed to determine if a correlation can be drawn between brain cancer and exposure to petroleum products.

"A lot of other studies need to be done in other plants. It will take a great deal more research to draw correlations."

The cancer mortality rates among workers who are still employed at the time of their death is always higher than the expected rate from all causes of death in the population as a whole, she noted.

"Some chronic disabling diseases prevent some individuals from becoming employed in the first place, or force them to stop working many years prior to their death so these types of deaths are not as prominent within the worker population," she explained. "This is what we call the healthy worker effect."

She said researchers will be examining the work histories of the 33 union members. A case control study of stomach cancer fatalities at the three plants also is under way.

"The new results indicate there may be an increased frequency of stomach cancers and leukemia," she said.

Shortly after initial cancer study results were released last spring, NCI and NIOSH, with the cooperation of the companies, began in-depth epidemiological studies at the refineries.

The NCI's preliminary brain cancer findings were presented Oct. 28 at a New York Academy of Sciences meeting in New York. At that time, Gulf released the results of its in-house study into the deaths of 4,766 of its employees and retirees that occurred between June 15, 1935, and Dec. 31, 1979. The death certificates for 4,600 white males were examined.

Gulf's study, billed by the company as the "largest study of refinery employees ever reported in North America," included OCAW members as well as other union and non-union employees, said company spokesman Kirk Vogeley.

Using a different statistical methodology, the Gulf scientists observed 30 brain tumor deaths compared to an expected 31 brain tumor fatalities.

"In other words, the risk of these workers dying from brain tumors was essentially the same or slightly less than what would be expected for the total population of the United States," according to a Gulf study statement.

Despite company study results, Vogeley said, Gulf is continuing its investigation and is participating in four other studies, including NCI's.

"We have cooperated with NCI, OCAW, NIOSH in the NCI study by providing data and assistance. We will continue to work with them in investigating this important matter," the Gulf statement continued.

Texaco spokesman Mike McDermott said "studies of individual cases conducted to date have failed to confirm any correlation between work performed and cause of death."

Texaco and Gulf are participating in a study by the University of Texas, compiling and reviewing mortality records of workers at several sites in both Orange and Jefferson counties. Like Gulf, Texaco is conducting an in-house study.

In addition, both companies are involved in a comprehensive American Petroleum Institute study examining 100 petroleum refineries for adverse health effects in the workplace.

Mobil spokesman John Flint said the company has hired the Stanford Research Institute to "conduct an in-depth study of our entire mortality experience at the Beaumont refinery. We will share the results with government agencies as well as anyone else, including employees, when the study is complete in late 1981."

STG 2433933

RSV0034014

The Galveston Daily News Tuesday Morning, October 28, 1980

Xc: Kating
Bisher
Collins
D. J. K. to

Byano
Brasfield
F.I.
Stone

TC brain cancer deaths apparently work-related

By LEE HERMAN
News Staff Writer

TEXAS CITY—Eighteen brain cancer deaths among workers at the Union Carbide plant here are apparently work-related, a government occupational health investigator said in New York Monday.

Dr. Victor Alexander, an Occupational Safety and Health Administration (OSHA) medical officer, added in a telephone interview Monday night that the 20-month study of employment records at Union Carbide showed "very, very solid" evidence that the deaths were job-related.

Investigators said it was the "largest single series of presumably occupationally related brain cancers" in medical annals, United Press International reported.

Alexander said the research, compiled by a team of OSHA and National Institute for Occupational Health and Safety (NIOSH) scientists, was consistent with three other studies on the subject.

The other studies involve Dow Chemical Corp.'s Freeport plant, three refineries in Port Arthur and a survey of Oil, Chemical and Atomic Worker Union Local 4-449 (Texas City) records.

The brain cancer deaths—four or five times as many as those projected for Galveston County under normal conditions—occurred between 1956 and 1980, UPI reported.

"We can think of no other explanation and are left with the conclusion that they are occupationally related."

"Until somebody comes up with another conclusion, which would be hard or impossible to do, we think they are occupationally related," Alexander said.

Alexander added that he presented the information to the New York Academy of Sciences Workshop on Brain Tumors in the Chemical Industry to allow scientists to study the OSHA-NIOSH data and "have their say."

The workshop is a government-funded, three-day series of seminars.

He said that although the study is only 85 percent complete, the findings were "striking enough" to report.

manager for Union Carbide in Texas City, said that he read a copy of the presentation to be made at the conference.

He said that in the prepared text, there was no definite connection made between the deaths and the workplace.

Ritchie added that he would wait for a final report, which Alexander said is probably from two to six months away, before making additional comments.

UPI reported that another Union Carbide official said there was no reason to believe there is a connection between the brain cancer and occupational exposure.

"Our position is that Dr. Alexander's is an interim report and that more will be learned over the next two years as company toxicologists and OSHA-NIOSH scientists continue the investigation," the spokesman said.

Although Alexander said the evidence showed the cancers were probably job-related, he added that the investigation has yet to reveal if any particular chemical caused them.

However, he added that he hopes to be able to pinpoint a certain chemical before the study is finished, "but even if you don't know the specific chemical, there are still some things that can be done."

Alexander said that scientists will be looking for a common link among the cancer victims at the various plants in hope of determining what might be changed to lessen the risk.

"If workers are exposed to something that causes this cancer, we can almost be certain that that malignancy can be prevented," UPI reported NIOSH scientist Dr. Anthony Robbins said.

Although vinyl chloride has been suggested as a possible cause, Alexander said the study had not "made a strong tie" between the resin and these cancers.

"Even after we have looked over all the case histories, we can't link it to vinyl chloride," Alexander

said.

Six separate suits filed against Union Carbide by former employees or their widows between March, 1979 and Jan., 1980 have blamed vinyl chloride for illnesses and deaths.

The study of about 8,600 employees at Union Carbide—which has been "extremely cooperative," Alexander said—began after a worker complained to the government in November of 1978.

UPI reported that the median employment of those with fatal brain cancer was 21 years.

In February of 1979, the federal government announced the joint investigation by NIOSH and OSHA scientists.

Alexander added that although the study may result in new regulations, he is not trying to scare present workers.

"This isn't something where people should stop working. They should just be careful" and "stick to basics," he said.

Alexander said that data from related studies will be presented at the New York conference.

② f -
Brown
Yuma
Feb

STG 2433934

Science

Brain tumor risk in petrochemical workers

Epidemiologists struggle to sort out tenuous evidence that points to elevated incidence of brain tumors among petrochemical workers

Jeffrey L. Fox
C&EN, Washington

The current effort to show a link between exposures to petrochemicals and the incidence of brain tumors among workers in the petrochemical industry is stretching epidemiology to its limits.

Some critics even argue that epidemiologists risk a tumble by leaning somewhat beyond the reasonable limits of their science. This possibility was apparent at the recent workshop sponsored by the New York Academy of Sciences at which was assembled a blue-ribbon group of epidemiologists and other authorities familiar with brain tumor incidence in humans and in experimental animals (C&EN, Nov. 3, page 6). Some observers at the workshop could not help concluding that establishing a firm link between brain tumor incidence and employment in the petrochemical industry may prove very difficult. Moreover, even if established, that link may

represent the result of practices long since outmoded. Many naive and dangerously sloppy procedures of several decades ago have been eliminated.

But sloppy procedures still exist, and other procedures that are genuinely thought to be safe well might be otherwise. Thus, the petrochemical industry—and any other industry, for that matter—must rely on imperfect early warning systems, such as those mustered by epidemiologists, to ferret out potential health hazards in the workplace.

Epidemiologists often refer to their preliminary surveys as "hypothesis generating" devices. The current hypothesis, being voiced by epidemiologists working for the Occupational Safety & Health Administration, for the National Institute for Occupational Safety & Health, and for the National Cancer Institute, is that the incidence of brain tumors among petrochemical employees is anywhere from 1.5 to 2.5 times higher than the incidence in the general population. Several other epidemiologists, some of them employed by the petrochemical industry but others without such ties, do not find evidence for elevated incidence. These disagreements were by no means reconciled during the New York Academy of Sciences workshop.

"Polarizations came to mind during preparation of this meeting," says workshop organizer and epidemiologist Irving P. Selikoff. "We're conscious of this. But we were taught it was good to find causes of disease. Now, when causes in industry are found, there's no automatic pat on the back." There are also no automatic pats on the back when someone finds the obverse, no evidence for causes, he could have added.

Epidemiologic evidence regarding brain tumors is anything but straightforward. Many factors could serve to show the intense analytic efforts under way at the Texas petrochemical plants, including facilities run by Union Carbide in Texas City and Dow Chemical in Freeport but also at other installations owned by Texaco, Mobil, and Gulf Oil.

For instance, the incidence of brain tumors in the general population in the U.S. is on the rise, according to



Nation: dosage and route are important

Lawrence Garfinkel of the American Cancer Society. The rise is not a sharp one; it applies more to men than women; and it follows no immediately obvious pattern. Also, the apparent rise in incidence might be due, in part, to improved diagnostic procedures and to better reporting of cancers.

Quibbling about statistics and how to analyze them is just the sort of activity that preoccupies many of the epidemiologists now scrutinizing the petrochemical industry. There are plenty of good reasons for the quibbling—scarce data, reliance on old records of questionable validity, and sharp differences on how to approach an issue are all factors.

"It is clear there is a rise in the risk of brain cancer," asserted NIOSH's Anthony Robbins at the beginning of the workshop. "From the point of view of the workers, we've been too slow in recognizing that risk. We are attracting attention to this issue [by having a public workshop]—not to blame anyone. But until you bring the best minds together, you don't solve the problem."

What Robbins said was echoed often during the three-day meeting, particularly by his colleagues from NIOSH and OSHA. Their concerns, quite properly, revolve around the principle of ensuring the health of workers. But those concerns can act like a centripetal force, continually drawing one's attentions back to



Robbins: bring the best minds together

Animal studies provide troublesome hints

Ideally, the results of carefully controlled animal studies of carcinogens will complement epidemiology, confirming suspicions about humans by referring to systematic data about mice and rats.

But a good case may be made that animal studies are sometimes misleading and can be prejudicial to the epidemiologist. For example, humans might be more, less, or altogether insensitive to a particular chemical that induces cancer in rats and mice. How often such discrepancies crop up is not so important here. The point is that the existence of bad or even good animal studies can send epidemiologists off on an oriented search, but a wild goose chase nevertheless.

To anyone who is strict about the conduct of scientific investigations, the question of initial biases—of whatever nature—is bound to be troublesome. Thus, conscientious epidemiologists are likely to find themselves trapped in a paradox: wanting and needing helpful clues about enigmatic human health problems, but also wishing to avoid one of the worst pitfalls in science—namely, relying on false premises.

There is plentiful animal data for the epidemiologists searching for causes of human brain tumors to review. Scientists have identified several dozen chemical compounds that can cause brain tumors in various animals, including mice, rats, hamsters, rabbits, and nonhuman primates. When tested, however, many of these chemicals do not cause cancer consistently from one type of animal to another. Moreover, the anatomic site of tumors caused by a particular agent can vary from one species of animal to another. This lack of consistency is fairly commonplace in animal testing programs. It's a complicating factor that helps to muddy otherwise limpid clues now available.

Not all brain tumors are alike. The major kind of tumor being followed in the epidemiologic studies is called a glioma. The spontaneous incidence of gliomas in mice is "exceedingly low," according to Italian toxicologist, Cesare Maltoni of the University of Bologna. Thus, the incidence of gliomas arising after exposure to various chemicals is a relatively reliable test system, he says. Nonethe-

less, that incidence also varies according to how the chemicals are administered to the animals—by feeding, injection, implantation, or inhalation. "It is important to know the dosage and route by which a carcinogen gains entrance," Maltoni emphasizes.

Other observations of animals undoubtedly will complicate the assessment of the human health question. For example, there is the matter of varying susceptibility to particular cancer-causing agents during an animal's life span. "Embryos and newborn animals are far more sensitive to many agents than are adult animals," Maltoni notes, referring to the incidence of central nervous system tumors.

This age-dependent variation in susceptibility is pronounced in *Erythrocebus patas*, a species of reddish, African monkey being studied by Jerry M. Rice and colleagues at the National Cancer Institute in Bethesda, Md. "We cannot get central nervous system tumors by injection of carcinogens into adults," he says about such animals. "They need to have very early exposure during gestation to develop tumors later, two to three years after birth."

To Rice, these results suggest that there may be periods in life when humans are particularly or peculiarly susceptible to developing certain kinds of tumors. Thus, induction of brain tumors in adults "might have occurred very much longer in the past," he notes.

What all this means regarding workplace exposures to putative carcinogens and the incidence of brain tumors is difficult to say. And when these observations are tied into a widely held view of how cancer develops, the story looks even more complex.

Many scientists believe that cancer develops in two phases, called initiation and promotion. An early event, initiation, caused by one type of chemical, may not cause a tumor unless followed by exposure to (usually) other types of chemicals, called promoters. Thus, the search for a single type of workplace exposure to explain brain cancer incidence might be misguided.

Imagine the following situation. One group of adults, when children, either had a particular disease, took a certain

medicine, or was exposed to some other "initiating agent," whereas another group of adults avoided that agent. Then during adulthood, members from both groups were met with an onslaught of many "promoting agents." Some such promoting agents might be found in the workplace, particularly when that workplace is a petrochemical facility. But other kinds of workplaces and altogether different kinds of environments also might provide some of the adults with deleterious promoting agents. Sorting through all these possibilities to identify the culprit agents could prove a formidable challenge.

Studies of animals are introducing further complications. For example, some scientists contend that virus particles are associated with chemically induced brain tumors in mice. Other scientists dispute the importance and even the existence of such particles. Their importance to human brain tumors is unknown.

Other observations are puzzling. Why, for instance, does a particular chemical produce primarily brain tumors without affecting other organs? Access of the chemical to such organs partly accounts for their differing sensitivity. But other metabolic influences, including either different activation or detoxification in various organs, also could be important, according to Paul Kleihus of the University of Freiburg in West Germany. Thus, for instance, enzymes in the brain cells of rats are slow to rid DNA there of certain kinds of damage. Long-lived damage might lead to brain tumors during the later life of such animals.

By far, the most puzzling phenomenon is one noted by Maltoni and his colleagues. Varying the dose and the means by which it's administered for some carcinogens can cause systematic fluctuations in brain tumor incidence in animals. Thus, an elevated dose of vinyl chloride, for example, induces few (or no) brain tumors, whereas a lower dose may induce many such tumors. This "masking" phenomenon departs significantly from the typical observation that tumor incidence rises directly in proportion to the dose of carcinogen an animal receives. If such a phenomenon pertains to human brain cancer, the search for pertinent workplace hazards may be tortuous indeed. □

workers and their environment. Thus, well-intentioned beliefs can predispose one to look for clues more zealously in one place than in another. This is not meant to say that OSHA or NIOSH officials are tailoring their investigations to distort the truth. But these epidemiologists clearly are instilled with the belief that the workplace, particularly a modern

petrochemical facility, can and must be safeguarded.

That commitment, which was on clear display among many of the federal officials at the workshop, is naturally even stronger among officials of unions representing the petrochemical workforce. "We must identify the job places where cases of brain cancer have appeared to see if

there's a common denominator," says Raphael Moure of the Oil, Chemical & Atomic Workers International Union. "What are the locations in the plant?" Once that's known, he says, "conditions can be improved and exposures minimized." Like Moure, Sheldon Samuels of AFL-CIO now assumes that the risk for brain tumors is elevated for workers in pet-



Thomas: analyses suggest elevated risk

rochemical plants. "The failure to identify specific agents is a disappointment, but not an insurmountable difficulty," he says.

Although a fear about elevated brain cancer incidence certainly is justifiable, the statements of the two union officials indicate they have already gone from fear, to belief, to plan of action. Here again, their professional commitments more or less make that course inevitable. But it

comes when many of the epidemiologists are arguing over whether there is any elevated risk. If the risk is the same as that faced by the general population, it seems a rather striking challenge to the petrochemical industry to improve the workplace.

"I'm puzzled by the widespread nature of the problem [of brain tumor incidence]," says H. Michael Utidjian of Union Carbide. "It seems widespread over many populations and work classes, sometimes extending to white collar workers. I don't know of any other case of occupationally related cancer where it's appeared over such a wide range of workplaces." It is "oversimplistic," he adds, to say conclusively that brain tumors are due to exposures in the chemical industry.

Several epidemiologists have concluded that brain tumor incidence, at least among some petrochemical workers, is not elevated. For example, one such study was conducted by Chi-Pang Wen of Gulf Oil's medical and health resources division of the company's oil refinery in Texas—the same installation included in an NCI survey. Wen says his study "represents the largest retrospective cohort study of refinery workers to date in North America in terms of length of observation (44 years of employment experience) and person-years of observation." And he concludes that "the results indicate no increased risk of brain tumors among workers at this Texas refinery."

He and NCI's Terry L. Thomas (and her collaborators) differ sharply in their conclusions. "Our analyses suggest that refinery workers have an elevated risk of brain cancer," Thomas says. And it's possible that current studies "underestimate" that risk, she adds.

Eventually, their differing conclusions have to be reconciled: They are, after all, looking at overlapping (albeit not identical) populations. Hence, it seems that discrepancies in analytic methods and in deciding what information to include stand in the way of reconciliation.

Yet other epidemiologists, looking at different workplaces, come up with all kinds of puzzling clues. Michael R. Alderson of the Institute of Cancer Research in the U.K. has surveyed eight refineries there and concludes there is no increased risk for brain cancer, although preliminary evidence for increases in malignant melanoma (skin cancer) is being followed up, he says. "But even if you accept some of the U.S. data, the problem of brain cancer is in no way like other hazards."

Hints from other surveys provide a kaleidoscope of incongruent obser-



Wen: results indicate no increased risk

ventions. Women more than men employed in the Swedish chemical industry might be at greater than normal risk for brain cancer. Workers in the rubber tire building industry in the U.S. are also at greater than normal risk, says Thomas Mancuso of the University of Pittsburgh School of Public Health. But so are workers in the electrical machinery and also paper (and allied products) industries. And so also are workers in certain segments of the transportation industry, according to epidemiologists at the University of Southern California in Los Angeles.

Epidemiology is very much an outgrowth of clinical medicine. According to one wag, epidemiologic surveys are often really "hunches, shored up with statistics."

However, the fact is that many clinicians have hit upon important medical treatments by following up their hunches. The same phenomenon may be at work among the epidemiologists scrutinizing brain cancer incidence. Hence, the possibility that they are onto something means that their surveys, no matter how preliminary, cannot be disregarded.

All of this close scrutiny also may serve a broader purpose. For example, Selikoff and his colleague William J. Nicholson at Mount Sinai School of Medicine in New York City find evidence of higher incidence for other kinds of cancer besides brain tumors among several groups of industrial workers. "Brain tumors may be 'the canary' for locating other cancer risks," Nicholson says. Thus elevated risks of diseases, such as lung cancer and malignant melanoma, cited by several of the epidemiologists at the New York Academy of Sciences workshop well could prove to be a more significant public health problem than brain cancer. □

Many chemicals induce brain tumors in animals

2-Acetylaminofluorene
Acrylonitrile
1-Aryl-3,3-dialkyltriazenes
Azoethane*
Azoxyethane*
Azoxy methane
Butylnitrosourea
Cycasin
1,2-Diethylhydrazine
Diethylnitrosamine*
Diethyl sulfate*
Dimethylbenzanthracene
Dimethyl sulfate*
Elomycin
Ethyl methanesulphonate
Ethylnitrosobutyl*
Ethylnitrosourea*
1-Methyl-2-benzyl-hydrazine*
Methyl methanesulphonate
Methylnitrosourea*
Methylnitrosourethane
(intrapleural)
Procarbazine*
Propane sulfone
Propylene imine
Pyrrolizidine alkaloids
Vinyl chloride

* These chemicals can cross the placenta to induce brain tumors in offspring. Source: Victor Alexander et al., MOSH

ANOTHER TECHNOLOGICAL ADVANCE FROM SILARP.

MURPHY'S LAW DEFIED.

“If anything can go wrong, it will” and usually does, especially when you’re entering an equation into a calculator.

Until now, you may have been using the error-prone procedure of encoding and decoding the steps of an assembly code. The Sharp 5100 Scientific Calculator eliminates this needless procedure. It's the first programmable calculator that allows you to enter directly, using algebraic, logarithmic, exponential and other functions in their familiar form.

That's Murphy's Law defied.

The 5100 is also the first hand-held to give you a complete display of the equation: you see 24 characters at a time in a dot-matrix display that rolls right or left to accommodate one or more formulas of up to 80 steps. So that while other programmable calculators display only one *program step* at a time, the 5100 displays the actual equation in its entirety, and in the way you're accustomed to seeing it.


If, anywhere along the line, you've entered an incorrect character (Murphy's Law at work), the display makes it easy for you to spot. You can then move the 5100's computer-like cursor to any part of the display to correct it.

In addition to the fast solution of frequently used formulas, the 5100 is extremely convenient for deriving difference tables, performing data analysis, numerical integration, simulations of linear systems, statistical and probability calculations, and many other procedures. It has so many uses, it would take a book to fully describe them, and someone has even written one.²

Sharp offers you a complete line of highly advanced scientifics, and if you want the leading edge in scientific calculators, buy the 5100 (under \$90.00)*.

*Suggested Retail Price

*Suggested Retail Price



THE SHARP EDGE

SHARP

Sharp Electronics Corp. 10 Keystone Place Paramus, N.Y. 07652

STG 2433931

RSV0034019