REPORT TO

MONSANTO COMPANY

MUTAGENIC STUDY WITH
AROCLOR 1254
IN ALBINO MICE

JANUARY 13, 1972

IBT NO. E622

## I. Introduction

A mutation is a change in the character of a gene such that morphologic, physiologic, and/or biochemical alterations are produced. If this change in gene character occurs in the germinal cell, the alteration can be transmitted to succeeding generations. Changes of this nature can be artificially induced (irradiation, chemical exposure, etc.) or they may be spontaneous. A dominant lethal mutation, occurring in the male germinal cell, may lead to the inability of the affected cell to fertilize an egg or, once having fertilized, to the failure of development beyond the blastocyst stage (implantation). Male mice, treated with the test compound, are mated with untreated females. The numbers of pre-implantation losses and early resorptions in female mice, dissected at mid-gestation, are used to calculate the mutation rate based on the induction of dominant lethal mutations.



STLCOPCB4018801

## II. Summary

The potential mutagenicity of Aroclor 1254 was investigated by conducting a dominant lethal study using albino mice treated with either 500 (T-I) or 1,000 (T-II) mg of the material/kg of body weight. Methyl methanesulfonate was used as a positive control.

Three T-II males died within 2 days of treatment and 2 positive control males died during the course of the study.

At weeks 4 and 6 the mating index for T-II animals was somewhat low. However, mating indices of T-I animals were normal.

Autopsy findings and mutation rates for Aroclor 1254 treated animals were not unusual and compared favorably to the contemporary control findings.

The MMS treated animals showed a positive response for the first 2 weeks post-treatment.



STLCOPCB4018802

Therefore, treating male mice with a single intraperitoneal injection of either 500 or 1,000 mg Aroclor 1254/kg of body weight did not induce dominant lethal mutations.

Respectfully submitted,

INDUSTRIAL BIO-TEST LABORATORIES, INC.

Report prepared by: Dennis Arnold

Dennis Arnold, B.S.
Group Leader
Mutagenic Studies

Report approved by: Gerald Kennedy

Gerald Kennedy, B.S.
Senior Group Leader
Metabolic and Mutagenic Studies

M. L. Keplinger

M. L. Keplinger, Ph.D.
Manager, Toxicology

January 13, 1972

trm:psh

DSW 034976

STLCOPCB4018803

## III. Procedure

### A. Outline of Experiment

The test material was Aroclor 1254. Methyl methanesulfonate was used as a positive control. Charles River strain albino mice were received at this laboratory at 60 to 70 days of age for use in the study. The organization of groups is presented in Table I.

TABLE I

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Organization of Groups

| Group | Dose Level* (mg/kg) | Number of Males Treated |
|---|---|---|
| C | - | 12 |
| PC | 100 | 12 |
| T-I | 500 | 12 |
| T-II | 1,000 | 12 |

* Aroclor 1254 was administered as a 10 percent solution in corn oil. Control males received the vehicle in amounts equivalent to those received by the T-II males. Positive control males received a 1 percent corn oil solution of MMS.



STLCOPCB4018804

B. Dosage Levels

The treatment levels were selected by intraperitoneal administration of single graded doses of the test material to male mice. The maximum tolerated dose was determined and dose levels employed in the main study were based on these findings. The test material was administered intraperitoneally as a corn oil solution to male mice. Control animals received the vehicle in volumes equivalent to those given the high test group and positive control animals received methyl methanesulfonate (MMS).

C. Mating Schedule

Each group consisted of 12 male mice, each of which was placed in a cage with 3 untreated virgin females immediately after dose administration. At the end of 1 week, the females were removed from the cage and replaced by another group of 3 females. This procedure continued for 6 consecutive weeks, a period of time required for maturation of the male mouse germ cells from the spermatocyte to the mature spermatozoon. Experiments at this laboratory have shown that the reference mutagenic compound* affects germinal cells in the meiotic and post-meiotic stages prior to or during the spermatid stage. Recovery from the effects of MMS is generally seen 3 weeks post-treatment. Following the sixth week of mating, all males were sacrificed.

D. Female Sacrifice

The females were sacrificed approximately 1 week after removal from the breeding cage, at which time most females that had

* The reference mutagen is methyl methanesulfonate (MMS).



STLCOPCB4018805

mated were at mid-pregnancy. All animals were sacrificed by carbon dioxide asphyxiation. The numbers of implantation sites, resorption sites, and embryos were recorded. Females were judged to be pregnant if corpora lutea were present in the ovaries.

Resorption sites were divided into 2 groups, early deaths (deciduomata) and late deaths. Late deaths refer to embryos which develop to a relatively advanced stage prior to death, so that the placenta and fetal membranes or remnants thereof are visible. The frequency of late deaths is apparently unaffected by mutagens and, therefore, would appear to be non-genetic in this test system. Deciduomata occur at the sites of implanted blastocysts which fail to develop following implantation. The mutagenicity of the chemical can be measured by the proportions of all implantations which are deciduomata.

Mutagenicity can also be measured by comparing the mean number of viable embryos in the test group to the number obtained in the control group. Calculation of the mutation rate using this criterion assumes that pre-implantation losses among non-affected test animals will be essentially the same as losses observed among control animals.



STLCOPCB4018806

## IV. Results

### A. Treatment Levels

Male mice were each given a single intraperitoneal injection of either 30, 100, 300, or 1,000 mg of Aroclor 1254 per kg of body weight. No effects were noted at any of the levels tested. Treatment levels chosen for the study were 500 (T-I) and 1,000 (T-II) mg/kg of body weight.

### B. Mortality and Reactions

Two positive control males died, 1 on the day of treatment and 1 during week 3. Three T-II males died within 2 days of treatment.

### C. Mating Performance

Mating indices, defined as the number of animals pregnant divided by the number of females mated times 100, and the number of surviving males for each post-treatment week are shown in Table II. The mating indices of T-II animals were slightly low at weeks 4 and 6. However, all mating indices seen in the experiment are considered to be normal for the mouse strain employed.



STLCOPCB4018807

TABLE II

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Mating Performance

| Group | Test Week Number | Number of Surviving Males | Number of Animals Pregnant / Number of Females Mated | Mating Index (Percent) |
|---|---|---|---|---|
| C | 1 | 12 | 25/36 | 69.4 |
| | 2 | 12 | 28/36 | 77.8 |
| | 3 | 12 | 31/36 | 86.1 |
| | 4 | 12 | 29/36 | 80.6 |
| | 5 | 12 | 28/36 | 77.8 |
| | 6 | 12 | 29/36 | 80.6 |
| PC | 1 | 11 | 14/33 | 42.4 |
| | 2 | 11 | 19/33 | 57.6 |
| | 3 | 11 | 23/33 | 69.7 |
| | 4 | 10 | 26/30 | 86.7 |
| | 5 | 10 | 23/30 | 76.7 |
| | 6 | 10 | 24/30 | 80.0 |

DSW 034981

STLCOPCB4018808

TABLE II continued

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Mating Performance

| Group | Test Week Number | Number of Surviving Males | Number of Animals Pregnant / Number of Females Mated | Mating Index (Percent) |
|---|---|---|---|---|
| T-I | 1 | 12 | 26/36 | 72.2 |
| | 2 | 12 | 24/36 | 66.7 |
| | 3 | 12 | 24/36 | 66.7 |
| | 4 | 12 | 28/36 | 77.8 |
| | 5 | 12 | 31/36 | 86.1 |
| | 6 | 12 | 30/36 | 83.3 |
| T-II | 1 | 9 | 18/27 | 66.7 |
| | 2 | 9 | 16/27 | 59.2 |
| | 3 | 9 | 17/27 | 63.0 |
| | 4 | 9 | 13/27 | 48.1 |
| | 5 | 9 | 14/27 | 51.8 |
| | 6 | 9 | 12/27 | 44.4 |

DSW 034982

STLCOPCB4018809

D. Sacrifice Data

Data pertaining to the numbers of corpora lutea, implantation sites, resorption sites (early and late differentiated), and embryos are presented in Table III. Laboratory data have shown that the untreated female mouse has a mean of 13.5 corpora lutea. Therefore, the number of pregnant animals is multiplied by 13.5 to obtain the total number of corpora lutea.

The corpora lutea for the pseudo-pregnant female in the control group (at week 3) that did not have implantations are included in the results. These data show no significant differences between test and control animals with regard to the numbers of implantations, resorptions, and embryos.



STLCOPCB4018810

TABLE III

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Summary of Sacrifice Data

| Group | Test Week Number | Pregnant Animals Examined | Calculated Corpora Lutea | Implantation Sites | Resorption Sites Early | Resorption Sites Late | Embryos |
|---|---|---|---|---|---|---|---|
| C | 1 | 25 | 338 | 293 (11.7) | 5 (0.2) | 1 (0.1) | 287 (11.5) |
| | 2 | 28 | 378 | 355 (12.7) | 9 (0.3) | 0 (0.0) | 346 (12.4) |
| | 3 | 31* | 417 | 342 (11.0) | 14 (0.4) | 5 (0.2) | 323 (10.4) |
| | 4 | 29 | 392 | 357 (12.3) | 15 (0.5) | 5 (0.2) | 337 (11.6) |
| | 5 | 28 | 378 | 341 (12.2) | 13 (0.5) | 3 (0.1) | 325 (11.6) |
| | 6 | 29 | 392 | 349 (12.0) | 12 (0.4) | 3 (0.1) | 334 (11.5) |
| PC | 1 | 14 | 189 | 134 ( 9.6) | 41 (2.9) | 2 (0.1) | 91 ( 6.5) |
| | 2 | 19 | 256 | 178 ( 9.4) | 47 (2.5) | 0 (0.0) | 131 ( 6.9) |
| | 3 | 23 | 310 | 294 (12.8) | 13 (0.6) | 3 (0.1) | 278 (12.1) |
| | 4 | 26 | 351 | 334 (12.8) | 16 (0.6) | 4 (0.2) | 314 (12.1) |
| | 5 | 23 | 310 | 291 (12.6) | 14 (0.6) | 5 (0.2) | 272 (11.8) |
| | 6 | 24 | 324 | 292 (12.2) | 17 (0.7) | 2 (0.1) | 273 (11.4) |

Note: Numbers in parentheses are the means per female.

* Includes 1 pseudo-pregnant female (corpora lutea only).

DSW 034984



STLCOPCB4018811

TABLE III continued

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Summary of Sacrifice Data

| Group | Test Week Number | Pregnant Animals Examined | Calculated Corpora Lutea | Implantation Sites | Resorption Sites Early | Resorption Sites Late | Embryos |
|---|---|---|---|---|---|---|---|
| T-I | 1 | 26 | 351 | 279 (10.7) | 3 (0.1) | 3 (0.1) | 273 (10.5) |
| | 2 | 24 | 324 | 281 (11.7) | 5 (0.2) | 1 (0.1) | 275 (11.4) |
| | 3 | 24 | 324 | 315 (13.1) | 11 (0.4) | 0 (0.0) | 304 (12.7) |
| | 4 | 28 | 378 | 345 (12.3) | 19 (0.7) | 2 (0.1) | 324 (11.6) |
| | 5 | 31 | 418 | 403 (13.0) | 18 (0.6) | 4 (0.1) | 381 (12.3) |
| | 6 | 30 | 405 | 354 (11.8) | 17 (0.6) | 4 (0.1) | 333 (11.1) |
| T-II | 1 | 18 | 243 | 214 (11.9) | 9 (0.5) | 2 (0.1) | 203 (11.3) |
| | 2 | 16 | 216 | 197 (12.3) | 3 (0.2) | 3 (0.2) | 191 (11.9) |
| | 3 | 17 | 230 | 220 (12.9) | 3 (0.2) | 0 (0.0) | 217 (12.8) |
| | 4 | 13 | 176 | 150 (11.5) | 5 (0.4) | 1 (0.1) | 144 (11.1) |
| | 5 | 14 | 189 | 185 (13.2) | 9 (0.6) | 1 (0.1) | 175 (12.5) |
| | 6 | 12 | 162 | 138 (11.5) | 4 (0.3) | 2 (0.2) | 132 (11.0) |

Note: Numbers in parentheses are the means per female.

DSW 034985



STLCOPCB4018812

E. Mutagenic Data

A summary of results for the mutagenic study is presented in Table IV.

Pre-Implantation Loss is defined as:

$$\frac{\text{Number of Corpora Lutea - Number of Implantation Sites}}{\text{Number of Corpora Lutea}} \times 100$$

The mutation rate may be calculated by comparing the number of early resorptions (deciduomata) to the total number of implantations for each group.

$$\frac{\text{Number of Early Resorption Sites}}{\text{Number of Implantation Sites}} \times 100 \quad \text{(A in table)}$$

Another way of expressing the mutation rate, which takes into account pre-implantation losses, is by comparing the mean number of normal embryos in each test group to the mean number of embryos in the control group.

$$100 - \left( \frac{\text{Embryos Test Group/Female}}{\text{Embryos Control Group/Female}} \times 100 \right) \quad \text{(B in table)}$$

Values presented were obtained by comparing each group to the contemporary control group (a) and to cumulative control data (b). Values with a minus (-) sign indicate that the mean number of embryos for that group and week was greater than that of controls. The deviation from zero should not exceed 25 percent in the positive direction.



STLCOPCB4018813

Mutation rates among Aroclor 1254 treated animals are not unusual and give no indication of a dominant lethal response.



STLCOPCB4018814

TABLE IV

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Summary of Mutagenic Data

| Group | Test Week Number | Pre-Implantation Loss and Mutation Rates<br>Pre-Implantation Loss (Percent) | Mutation Rates<br>A | B<br>a | b |
|---|---|---|---|---|---|
| C | 1 | 13.5 | 1.7 | - | - 0.9 |
| | 2 | 6.1 | 2.5 | - | -10.7 |
| | 3 | 18.0 | 4.1 | - | 10.4 |
| | 4 | 8.9 | 4.2 | - | - 0.9 |
| | 5 | 9.8 | 3.8 | - | 0.9 |
| | 6 | 11.0 | 3.4 | - | 1.7 |
| PC | 1 | 29.1 | 30.6 | 43.5 | 43.0 |
| | 2 | 30.5 | 26.4 | 44.4 | 38.4 |
| | 3 | 5.2 | 4.4 | -16.3 | - 4.3 |
| | 4 | 4.8 | 4.8 | - 4.3 | - 5.2 |
| | 5 | 6.1 | 4.8 | - 1.7 | - 0.8 |
| | 6 | 9.9 | 5.8 | 0.9 | 2.6 |

DSW 034988

STLCOPCB4018815

TABLE IV continued

TEST MATERIAL: Aroclor 1254

Mutagenic Study - Albino Mice

Summary of Mutagenic Data

| Group | Test Week Number | Pre-Implantation Loss and Mutation Rates: Pre-Implantation Loss (Percent) | Mutation Rates A | Mutation Rates B a | Mutation Rates B b |
|---|---|---|---|---|---|
| T-I | 1 | 20.5 | 1.1 | 8.7 | 7.9 |
| | 2 | 13.3 | 1.8 | 8.1 | - 1.8 |
| | 3 | 2.8 | 3.5 | -22.1 | - 9.5 |
| | 4 | 8.7 | 5.5 | 0.0 | - 0.9 |
| | 5 | 3.6 | 4.5 | - 6.0 | - 5.1 |
| | 6 | 12.6 | 4.8 | 3.5 | 5.1 |
| T-II | 1 | 11.9 | 3.7 | 1.7 | 0.9 |
| | 2 | 8.8 | 1.5 | 4.0 | - 6.2 |
| | 3 | 4.3 | 1.4 | -23.1 | -10.3 |
| | 4 | 14.8 | 3.3 | 4.3 | 3.5 |
| | 5 | 2.1 | 4.9 | - 7.8 | - 6.8 |
| | 6 | 14.8 | 2.9 | 4.4 | 6.0 |

DSW 034989

STLCOPCB4018816