

57

59
76
18
40

74
73
21

13

7
0
9

0
2

3

7
1
1

1



Industrial Hygiene Digest

INDUSTRIAL HEALTH NEWS

LITERATURE ABSTRACTS

MEDICAL

ENGINEERING

CHEMICAL

TOXICOLOGICAL

LEGAL ... decisions and trends

MAY 2 - 1957

OF
MEDICINE

APRIL, 1957

(Vol. 21, No. 4)

INDUSTRIAL HYGIENE FOUNDATION

MELLON INSTITUTE

4400 FIFTH AVENUE • PITTSBURGH 13, PA.

**PLAINTIFF'S
EXHIBIT**

WV-00468

**PLAINTIFF'S
EXHIBIT**

13905

SC-ALL-06960

SCF-ALLF-01830

INDUSTRIAL HYGIENE FEATURES

Biological data which represents the correlated contributions of more than 4,000 scientists and the counsel of an additional 13,000.
Abstract No. 451.

What can be done about the shortage of physicians for industry?
Abstract No. 456.

Questions and answers on practical aspects of occupational dermatoses.
Abstract No. 465.

Circumstances in which soaps and detergents have been indicted as causing or aggravating dermatitis are much more complex than have been anticipated. Abstract No. 466.

An evaluation of the hazards of ozone and oxides of nitrogen, and factors modifying their toxicity.
Abstract Nos. 474 and 475.

Existing tendency to develop lung tumors may have some bearing in pulmonary carcinoma, experiment exposing mice to asbestos dust indicates. Abstract No. 506.

Hitherto unrecognized etiological factor in calcification of the pericardium revealed by clinicorontogenic study of workers employed in talc mills and mines.
Abstract No. 507.

How can the nuisance problem of noise in air-conditioning systems be eliminated? Abstract Nos. 513-517.

A proposed technique for measuring the noise of a ducted fan eliminates measuring errors caused by the existence of sounding waves.
Abstract No. 518.

What are the responsibilities of the medical profession in the use of x-rays and other ionizing radiation?
Abstract No. 528.

A study of a multiple Venturi wet collector. Abstract No. 541.

A study of carcinogenicity of atmospheric pollutants. Abstract No. 545.

In order to treat electric-arc steel furnace fume theoretically, accurate particle size distributions for each phase of the melting cycle must be known. Abstract No. 548.

How to go about hiring handicapped people and placing them in productive work. Abstract No. 553.

Safety statistics in their present form are only one indication of what a safety program is doing.
Abstract No. 554.





FOUNDATION FACTS

Foundation Facts is a monthly news-sheet issued
to members of the Industrial Hygiene Association. The Foundation
is a non-profit organization for the improvement of working conditions.

Volume 19

April, 1957

MAY 21 1957

OF
MEDICINE

No. 4

SCHRENK RECEIVES CUMMINGS MEMORIAL AWARD

Dr. H. H. Schrenk, Research Director of Industrial Hygiene Foundation, received the Donald E. Cummings Memorial Award from the American Industrial Hygiene Association for outstanding contributions to the knowledge and practice of the profession of industrial hygiene. The award was made at the Association's 18th Annual Meeting at the Industrial Health Conference in St. Louis on April 24.

Dr. Schrenk has directed environmental studies in industries throughout the United States and Canada and has been in charge of the programs of toxicological and biological research carried on in the Foundation's research laboratory.

Before joining the Foundation's staff Dr. Schrenk was associated with the Bureau of Mines from 1928 to 1948 where he was Chief of the Health Branch and directed laboratory and research investigations in industrial hygiene. The following year he joined the Commissioned Corps of the United States Public Health Service as Scientist Director. He served as Chief, Environmental Investigations Branch, Division of Industrial Hygiene. He directed the Donora smog investigation conducted by the United States Public Health Service.

In his Cummings Memorial Lecture on "The Growth and Progress of Industrial Hygiene," Dr. Schrenk stressed the need for research in industrial hygiene to keep pace with the new industrial developments and the hazards which may accompany them.

"The production of new materials carries with it the responsibility to make sure that the usefulness is not outweighed by risks to health," he said.

The benefits of many of the products available today, ranging from synthetic fibers, high energy fuels and new plastics, would not be available today unless it had not been for the efforts of industrial hygienists in protecting workers from trailing hazards. The role industrial hygiene plays in industrial progress is perhaps best illustrated by the development of nuclear energy and its application to chemical processes, Dr. Schrenk pointed out.

In order to be prepared for the industrial and social changes which are taking place at an ever-increasing pace, new tools, instruments and analytical procedures must be available to the industrial hygienist. An excellent example of progress in the analytical field, Dr. Schrenk stated, is the spectacular advance in gas chromatography. He pointed out also the rapid expansion in the fields of spectroscopy, photometry, electroanalytical chemistry, nuclear magnetic resonance, and polarographic instrumentation, and the many contributions in the more conventional analytical field.

"A knowledge of these new developments is essential in the development of the more practical instruments which are in day-to-day use in the field," he said.

"The true value of analytical tools depends upon our ability to interpret the results of such measurements in terms of a quantitative relationship with harmful or objectionable effects. This in turn requires a knowledge of the biological response to offending agents," Dr. Schrenk said.

Thus, the industrial hygienist cannot work alone in protecting the health of workers and consumers. "Laboratory studies of biological response serve the dual purpose of providing data for establishing standards and of providing better diagnostic

(Cont. on next page)

carborundum for sanding in the shoe industry; and slate or chalk for stone-dusting in c mines. Quartz sands have been prohibited since 1938 in grinding of glass, and exposure has been regulated. The higher cost of artificial abrasives in the glass and ceramic industries is offset by higher working speed.

-- Cond. from Bull. Hyg

- 503 Respiratory Symptoms and Pulmonary Disability in an Industrial Town. Survey of a Random Sample of the Population. I. T. T. Higgins, P. D. Oldham, A. L. Cochrane, and J. C. Gilson. *Brit. Med. J.* 2, 904-910 (Oct. 20, 1956).

In 1954 the Pneumoconiosis Research Unit of the Medical Research Council carried out a survey of the male population aged 55-64 of Leigh in Lancashire, in order to compare the prevalence of respiratory symptoms and disability in miners with that found in men who had worked only in other occupations. The random sample consisted of 135 miners and ex-miners and 85 non-miners. Pneumoconiosis was found in 23% of the miners and ex-miners. Chronic bronchitis was found in 10.7% of the non-miners, in 26.7% of the miners without pneumoconiosis, and in 12.9% of the miners with evidence of pneumoconiosis. Some of the conclusions are: (1) miners and ex-miners had significantly more chronic bronchitis, a lower ventilatory capacity, and more respiratory symptoms than men who had never worked in mines; (2) miners without pneumoconiosis were on the average more disabled than those with early simple pneumoconiosis; and (3) the differences between miners and non-miners could not be ascribed to differences in social class or smoking habits, and in the miners there was no obvious relationship between length of exposure to dust and bronchitis.

-- Cond. from Bull. Hyg.

- 504 The Influence of the Shape of Dust Particles on the Rate of Phagocytosis in Vitro. A. Lengerova, V. J. Lenger, M. Esslova, R. Tuscany, and M. Vollova. *Brit. J. Ind. Med.* 14, 43-46 (Jan. 1957).

The rate of phagocytosis of two shapes of dust particles—"long" and "round"—was studied in an investigation in vitro in cultures of macrophages from the spleen of chicken embryos. The round particles were phagocytosed preferentially. This result is discussed from the standpoint of existing theories of phagocytosis as well as from that of the relation of the rate of phagocytosis and the harmfulness of different kinds of dusts.

-- Authors' summary

- 505 The Prevalence of Tuberculosis at Necropsy in Progressive Massive Fibrosis of Coalworkers. D. Rivers, W. R. L. James, D. G. Davies, and S. Thomson. *Brit. J. Ind. Med.* 14, 39-42 (Jan. 1957).

One hundred fifty-three cases of coalworkers' pneumoconiosis were examined at necropsy for the presence of tubercle bacilli both by culture and by animal inoculation. The series contained 118 cases of progressive massive fibrosis which were the special object of this study. Tubercle bacilli were isolated from 41 cases (35%) of massive fibrosis, and in 31 (28%) of these naked-eye or histological evidence of tuberculosis was recognized pathologically. The pathological observations, therefore, correlated well with the bacteriological results. There is a marked difference between the prevalence of tuberculous infection found in a post-mortem series and that found in life in cases of progressive massive fibrosis. During life the observed prevalence is only 1.1%. The authors found tubercle bacilli in the bronchi at necropsy in 28%. Thus open tuberculosis is, in most cases, a late complication of massive fibrosis. Whether tuberculous infection occurs early or late cannot be deduced from this study.

-- Authors' summary

- 506 Pulmonary Tumors in Mice Exposed to Asbestos Dust. K. M. Lynch, F. A. McIver, and J. R. Cain. *Arch. Ind. Health* 15, 207-214 (Mar. 1957).

Mice of a selected strain (AC F₁) were exposed to a dusty atmosphere containing asbestos floats of a type used by others experimentally, at 150 to 500 million particles per cubic foot. The largest single experimental group consisted of 127 animals killed at 19 months after exposure for 17 months. This group was compared with 222 control animals. Primary lung tumors were demonstrated microscopically in 46% of the dusted mice and in

36% of the controls. Multiple pulmonary tumors were demonstrated in 18% of the dusted animals and in 8% of the control animals. No significant histologic or cytologic difference was observed between tumors in the two groups. No clear-cut proof of malignancy was found in any of the lung tumors studied. The authors have been unable to demonstrate any proof of the carcinogenicity of asbestos under the conditions of the experiment, but the increased incidence of multiple lung tumors in dusted animals must be regarded as a possible accentuation by the inhalation of asbestos dust of an existing tendency to develop lung tumors. Future experiments with asbestos should include the use of larger animals, probably monkeys. The equivocal nature of this report does not answer the questions raised by accumulating clinical evidence that asbestosis is strongly associated with, and therefore is a possible cause of, pulmonary carcinoma in man. -- Adapted from authors' summary

507 Calcification of the Pericardium Caused by a Hitherto Unrecognized Etiological Factor.
M. Kleinfeld and C. P. Giel. *Am. Heart J.* 52, 899-904 (Dec. 1956).

In a clinicoroentgenologic study of workers employed in talc mills and mines, calcification of the pericardium was observed in 3 of 37 workers. In one patient the condition had existed for 15 years. The advanced age of these patients (79, 68, and 64 years) is in contrast to that found in chronic constrictive pericarditis, where the age range reported has varied from 10 to 59 years. Cardiac enlargement was absent, and the electrocardiographic patterns were essentially normal. The clinical picture revealed a chronic, slowly progressive course, with the predominant symptoms of pulmonary insufficiency consistent with the average course of talc pneumoconiosis. The normal electrocardiograms, lack of evidence of failure of either side of the heart, lack of response to nitroglycerin, and the absence of any intensifying effect of effort would tend to exclude angina pectoris. The exact mechanism responsible for the calcification of the pericardium in exposure to talc is not known. It can best be explained by a mechanical effect of the talc fibers resulting in a chronic inflammatory response and increased capillary hemorrhage with subsequent calcium deposition. -- Cond. from *J. Am. Med. Assn.*

508 Dust Exposures During Ash Removal from Incinerators. W. H. Megonnell, J. H. Ludwig, and L. Silverman. *Arch. Ind. Health* 15, 215-222 (Mar. 1957).

Dust exposures to the operator during ash removal are not excessive for the average institutional-type incinerator, as the stack draft through the open furnace doors draws most of the dust produced by the operation back into the incinerator. Installations with small stacks, or when reverse flow is created by wind action, produce higher and more persistent dust concentrations. Incinerator operation is poorer during rainy weather. Only minor dust exposures are produced when water quenching of the ash is used, and wetting is advisable when it can be accomplished. Airborne dust in incinerator rooms is of the same general composition as the ash. It is composed of a high percentage of small particles (50% less than 0.5 micron), with larger particles which are predominantly aggregates. Exposure period for operators removing ash is a function of the degree of cleaning accomplished and the quantity of ash removed. Stack draft and room ventilation appear to be the most important factors in limiting the concentration of airborne dust in the incinerator room. A large proportion of the dust exposure incidental to ash removal could be eliminated by careful disposal cans to take advantage of the effect of stack draft and by care in depositing ashes from the shovel into containers. Obviously the furnace door and grate opening should be kept open throughout all cleaning in the incinerator room. -- Cond. from authors' summary

509 Changes in Lungs of Experimental Animals after Intratracheal Introduction of Mineral-Wool Dust. V. A. Grimaïlovskaya and others. *Gigiena i Sanit.* 21, No. 8, 24-29 (1956). Russian.

Mineral-wool dust with relatively low free silica and moderate contents of bound silica produces in rats after 2-3 months a moderate sclerosis of diffuse and nodular types; after 5 months a peribronchial sclerosis starts. This is caused by solubility of dust particles in the lymph. -- Chem. Absts.

AFFIDAVIT OF DAVID S. EGILMAN

STATE OF MASSACHUSETTS)
) ss.
COUNTY OF NORFOLK)

Before me, the undersigned authority, personally appeared Dr. David Egilman and swore as follows:

1. My name is Dr. David Egilman. I am over the age of twenty-one years old and am otherwise competent to make this affidavit. All of the statements contained in this affidavit are based upon my personal knowledge and are true and correct.
2. I am the custodian of records of the Industrial Hygiene Foundation.
3. I have personally reviewed the documents listed below, copies of which are attached to this affidavit, and can attest that the attached copies are true, complete and correct copies of Industrial Hygiene Foundation documents.

Participants in Seventh Saranac Symposium with attached Affidavit dated May 21, 1991 of Michael Rhode, employee of the Armed Forces Institute of Pathology, custodian of original archives of Dr. Arthur J. Vorwald, authenticating the listing. (SCF-ALLF-01290)

*Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal.
January, 1950 (Vol. 14, No. 1) (SCF-ALLF-01180)*

*Industrial Hygiene Foundation, Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal.
September, 1952 (Vol. 16, No. 9) (SCF-ALLF-01370)*

*Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal.
September, 1956 (Vol. 20, No. 9) (SCF-ALLF-01760)*

*Industrial Hygiene Foundation, Industrial Hygiene Digest.
April, 1957 (Vol. 21, No. 4) (SCF-ALLF-01830)*

*Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal.
August, 1960 (Vol. 24, No. 8) (SCF-ALLF-02040)*

Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal.
March, 1961 (Vol. 25, No. 3) (SCF-ALLF-02140)

Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal. (SCF-ALLF-02310)

Industrial Hygiene Foundation, 1967 Annual Report. (SCF-ALLF-02836)

Industrial Hygiene Foundation, 1969 Annual Report. (SCF-ALLF-03082)

Industrial Hygiene Foundation of America, Inc., Report of Preliminary Dust Investigation for Asbestos Textile Institute with attached Memorandums on Plant of (various companies) Supplementing the General Report "Preliminary Dust Investigation for Asbestos Textile Institute" of June, 1947. (SCF-ALLF-01000)

Members of Air Hygiene Foundation / Industrial Hygiene Foundation as of November 30, 1937 through 1983. (SCF-ALLF-00470)

Air Hygiene Foundation of America, Inc., References to Recent Literature of Industrial Diseases, January 24, 1939. (SCF-ALLF-00565)

Foundation Facts, Vol. 3, No. 6 with attached Digest of Industrial Hygiene, Literature - News - Developments, dated June, 1941. (SCF-ALLF-00685)

Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal, Foundation Facts, August, 1957.
(SCF-ALLF-01875)

An Epidemiological Study of Lung Cancer in Asbestos Miners, Industrial Hygiene Foundation Report by Bruan & Truan; study made possible through a grant from the Quebec Asbestos Mining Association (Note: date taken from published article), June, 1958. (SCF-ALLF-01939)

Industrial Hygiene Digest, Industrial Health News, Literature Abstracts, Medical, Engineering, Chemical, Toxicological, Legal, May, 1963, Vol. 27, No. 5.
(SCF-ALLF-02375)

Preliminary Program, 28th Annual Meeting, Industrial Hygiene Foundation, October 23-24, 1963, to Mr. Wagner. (SCF-ALLF-02441)

Industrial Hygiene Foundation, 1964 Annual Report with attached membership list.
(SCF-ALLF-02468)

Industrial Hygiene Foundation of America, Inc., Asbestos Bioeffects Research for Industry, Medical Series, Bulletin No. 11, 1966. (SCF-ALLF-02683)

Further, affiant sayeth not.

David S. Egilman, MD, MPH

David S. Egilman, MD, MPH

Commonwealth of Massachusetts
Norfolk, ss

SUBSCRIBED AND SWORN TO ME, on this 8th day of May, 2006, to certify which
witness my hand and official seal of office.

NOTARY PUBLIC IN AND FOR
THE STATE OF MASSACHUSETTS

David S. Egilman

My Commission Expires March 1/2011

