

REQUEST FOR APPROVAL TO RELEASE
PROPRIETARY TECHNICAL INFORMATION



B-101256

1

Risk of Angiosarcoma in Vinyl Chloride Workers as
Predicted from Studies in Rats

P. J. Gehring, C. N. Park, and P. G. Watanabe

☐ 1. AN ABSTRACT
OF THE
PAPER
19
1978

Toxicology and Applied Pharmacology

RELATED PATENTS OR DISCLOSURES

RELATED PRODUCTS OR PROCESSES

RELATED REPORTS OF PUBLICATIONS

SUGGESTIONS BY AUTHOR FOR POSSIBLE REVIEWERS OF THIS MANUSCRIPT

K. Groves

T. Torkelson

R&S 101256

QUESTIONS CONCERNING THIS MANUSCRIPT ARE TO BE DIRECTED TO

P. J. Gehring

Tox Res Lab H&ER

636-1089

1803

Midland

15 May 1978

Rush

2

DEPARTMENTAL APPROVAL FOR RELEASE

BASchwert

18 May 78

May 16, 1978

DEPARTMENTAL COLLECTION OF REVIEWERS

3

COPIED IN MIDLAND CRI

JUL 16 1978

4

W. Blair

RISK OF ANGIOSARCOMA IN VINYL CHLORIDE WORKERS
AS PREDICTED FROM STUDIES IN RATS

by

P. J. Gehring, P. G. Watanabe
Toxicology Research Laboratory
Health and Environmental Research
Dow Chemical U.S.A.
Midland, Michigan 48640

and

C. N. Park
Computations Research Laboratory
Dow Chemical U.S.A.
Midland, Michigan 48640

Abbreviated Title: Vinyl Chloride-Risk Assessment

Send Correspondence to: Dr. P. J. Gehring
Toxicology Research Laboratory
Health and Environmental Research
Dow Chemical U.S.A.
1803 Building
Midland, MI 48640

R&S 101257

ABSTRACT .

Risk of Angiosarcoma in Vinyl Chloride Workers as Predicted from Studies in Rats. Gehring, P. J., Park, C. N., and Watanabe, P. G. (1978). Toxicol. Appl. Pharmacol. 00,000-000. Dose-response data for the induction of angiosarcoma in rats exposed to various levels of vinyl chloride (VC) together with attendant biotransformation data were used to estimate the risk of developing angiosarcoma in persons exposed to VC. Since a biotransformation product of VC, not VC per se, is responsible for the induction of angiosarcoma, the body surface area of people relative to rats was used to estimate the dose of the carcinogen biotransformed from VC by the former. Four models were used to extrapolate the data. Using a probit model, 10 hepatic angiosarcomas were predicted to occur in a recently reported epidemiological cohort of 9677 workers whereas 5 have occurred. Linear models and that based on the equation, $\text{Risk} = 1 - e^{-2x}$, where x = dose, do not appear as reliable. For an 8-hr day, 5 days/week, 35-yr time-weighted-average exposure of 1 ppm, the predicted incidence of hepatic angiosarcoma using the probit model is 1.5×10^{-8} .

R&S 101258

RISK OF ANGIOSARCOMA IN VINYL CHLORIDE WORKERS AS PREDICTED
FROM STUDIES IN RATS

INTRODUCTION

Exposure of rats (Maltoni and Lefemine, 1975) and man (Creech and Johnson, 1974; Tabershaw and Gaffey, 1974; Makk, et al, 1976; and Fox and Collier, 1977) to vinyl chloride (VC) has been associated with the development of hepatic angiosarcoma. Numerous studies in rats indicate that it is not VC per se which is responsible for production of angiosarcoma but rather a reactive metabolite formed from it in the body (Bartsch, et al, 1975; Barbin, et al, 1975; Kappus et al, 1976; Malavielle, et al, 1976; Watanabe, 1977; Bolt et al 1975; and Bolt, et al, 1976). Biotransformation of VC in rats is a nonlinear process occurring in accordance with Michaelis-Menten kinetics:

$$1) \quad v = V_m S / K_m + S$$

(Watanabe, et al, 1976a, 1976b, and 1976c and Bolt et al 1976). In this equation, v and V_m are velocity and maximum velocity, respectively, for the biotransformation of VC expressed as μg equivalents VC metabolized daily. S and K_m are the concentration of VC being inhaled and the Michaelis constant expressed as $\mu\text{g VC/l air}$, respectively.

Utilizing the foregoing information, Gehring, et al (1978) revealed a good correlation between the probit percent incidence of angiosarcoma in rats exposed to varying concentrations of VC, 4 hr daily, and the amount of vinyl chloride

biotransformed, v, μ g equivalents VC metabolized per day.

The probit equation is:

$$2) \quad \text{Probit Percent Incidence} = -1.625 + 1.543 \log v.$$

This correlation substantiates further the hypothesis that the induction of angiosarcoma is attributable to a biotransformation product of VC, not VC per se.

Hypothesizing that the biotransformation of VC is related directly to body surface area, Gehring, et al 1975 determined that the amount of VC transformed to a reactive form by man on a mass equivalent basis to rat is:

$$3) \quad V \text{ } \mu\text{g/8 hr} = \frac{1675 \text{ } \mu\text{g/hr} \cdot S_{\text{H}/\text{L}}}{850 \text{ } \mu\text{g/r} + S_{\text{H}/\text{L}}}.$$

Utilizing this relationship together with the foregoing probit equation, it was determined that incidence of hepatic angiosarcoma in workers exposed to 200 ppm should be approximately 13. This incidence is 50-fold higher than that experienced (Fox and Collier, 1977). Various plausible explanations were put forth to explain this difference including greater sensitivity of rats and increasing latency for development as v decreases.

One plausible explanation not mentioned was that in worker populations studied to-date, many were exposed for only fractions of their working life. Furthermore, for some, the

duration since initial exposure was likely insufficient for development of clinical detectable disease.

Recently available is a comprehensive epidemiological study¹ entitled, "Epidemiological Study of Vinyl Chloride Workers, Final Report". In this study, reasonably detailed information on 10,173 workers has been obtained for duration of exposure and lapsed time since first exposure.

Thus, much better information is available now to assess the reliability of the model of Gehring et al, 1978, for predicting the incidence of hepatic angiosarcoma in man utilizing data collected in rats. This assessment is reported here. In addition, three other models for estimating the incidence have been used for comparison of their reliability for predicting the incidence being experienced by workers exposed to vinyl chloride.

¹ Prepared for Manufacturing Chemists Association, 1825 Connecticut Avenue, N.W., Washington, D.C. 20009, Prepared by Equitable Environmental Health, Inc., 6000 Executive Blvd, Suite 308, Rockville, Maryland 20852, January 1978.

-4-

METHODS

Four models were selected to analyze their reliability for calculating the incidence of hepatic angiosarcoma in rats as function of μgVC equivalents metabolized daily by rats exposed 4 hr/day, 5 days/week for one year. These models were as follow:

A) Probit %

$$\begin{aligned}\text{Probit } \% &= a + b(\text{Log } v) \\ &= -1.625 + 1.543(\text{Log } v)\end{aligned}$$

B) Linear

$$\begin{aligned}\% &= a + b(v) \\ &= -1.48 + 0.339 \times 10^{-2}(v)\end{aligned}$$

C) Linear Forced through Origin

$$\begin{aligned}\% &= bv \\ &= 0.3565 \times 10^{-2}(v)\end{aligned}$$

D) One-Hit Model (Cancer Assessment Group, National Cancer Institute)

$$\begin{aligned}\% &= 1 - e^{-\beta(v)} \\ &= \{ 1 - \exp(-0.38 \times 10^{-4} \times v) \} 100\end{aligned}$$

The constants given for the preceding models were determined by linear regression analysis and represent the best fit of the data for the incidence of hepatic angiosarcoma versus the amount of VC metabolized daily in rats exposed to different concentrations of VC (see Gehring, et al, 1978).

The data used to derive the constants are given in Table 1. Using the equations derived for the models, the predicted incidence of angiosarcoma for the groups of rats exposed to different concentrations of VC were calculated. These results are shown also in Table 1.

All four models overpredict the incidence of hepatic angiosarcoma in rats exposed to 10,000 ppm VC. This is as expected because this exposure caused excessive mortality unrelated to development of angiosarcoma.

With respect to the predicted versus experimental incidence of hepatic angiosarcoma experienced by groups of rats exposed to the remaining concentrations of VC, none of the four models is unequivocally more reliable than any other. Thus, any one of the four models represents adequately the experimental data. Since no clear preference is discernible, all four models will be used to predict the incidence of hepatic angiosarcoma experienced or to be experienced by workers exposed to VC.

R&S 101263

RESULTS AND DISCUSSION

The incidences of hepatic angiosarcoma in subgroups of 9677 workers with respect to their duration of exposure are given in Table 2. These workers include 95.1% of the 10,173 in the cohort study; the remainder could not be traced.

The atmospheric concentrations of vinyl chloride to which these workers had been exposed were not quantitated, except subjectively into high, medium, and low categories. Since the time-weighted-average (TWA) exposure recommended by the American Conference of Governmental Industrial Hygienists (ACGIH) prior to 1972 was 500 ppm and subsequently 200 ppm until adoption of the Occupational Safety and Health Act standard of less than 1 ppm in 1974, it is assumed that even those exposures subjectively deemed low were high. Indeed, it is reasonable to expect the TWA exposures of 200 ppm and greater for 8 hours were common rather than the exceptions.

Table 3 depicts the average theoretical rat equivalent daily doses of biotransformation products of VC received by workers exposed to 200 or 500 ppm for 8 hr daily, 5 days/week adjusted for the fraction of their working life during which exposure occurred. The doses, v , are expressed as $\mu\text{g}/8 \text{ hr}$ and were calculated from equation 3 given in the introduction multiplied by the fraction of an assumed 35-year working life during which exposure occurred.

R&S 101264

Having estimated the rat equivalent daily doses of biotransformation products of VC received by the various subgroups of 9677 workmen exposed to a TWA of 200 or 500 ppm VC, the predicted incidence of hepatic angiosarcoma for each subgroup was calculated, Table 4. Examination of the data in this table indicates that Models C and D overestimate the incidence of hepatic angiosarcoma experienced by these workers while Model B underestimates the incidences. Models C and D may overpredict because they do not address either the enzymatic biotransformation of VC or repair of the lesion leading to development of angiosarcoma both of which are likely to be distributed normally in human or animal populations or because the time-weighted-average exposures for an 8 hr day were less than 200 ppm.

When a TWA of 200 ppm is utilized to represent the atmospheric concentration to which these workers were exposed, the predicted incidence of angiosarcoma of 10 cases using Model A compares favorably with that experienced, 5 cases. For a significant number of workers, the time elapsed since initial exposure has yet to reach 15 years, the shortest time needed for development of angiosarcoma in 1 of 5 of the afflicted workers, Table 2. Thus, it may be anticipated that a few additional cases of hepatic angiosarcoma will occur in this population. It is comforting that an epidemic of cases, as has been envisioned by some, is not to be anticipated.

As indicated in the introduction, the previous attempt to extrapolate data from experiments in rats to predict the occurrence of hepatic angiosarcoma in vinyl chloride workers appeared to overpredict the observed incidence even when differences in the biotransformation of VC to active products by man and rats were incorporated in the extrapolation.

The foregoing analysis using a probit percent incidence model 1 (Model A) has revealed that data collected in experiments on rats may be used to predict the incidence of hepatic angiosarcoma in man with reasonable accuracy. Further verification of this model will be assisted greatly by determination of the rate of biotransformation of VC by men exposed to sufficient concentrations of VC to allow such measurement. Currently, this must be estimated by the rate of biotransformation in rats and its assumed relationship to body surface area (Gehring, et al, 1973). Nonetheless, the correlation between predicted and observed incidence of hepatic angiosarcoma in vinyl chloride workers seems to justify both the assumptions and the model (Model A, Probit 3) used to make the predictions.

Since the predicted incidence of hepatic angiosarcoma in man using results from studies in rats appears justifiable, it is of interest to use the same procedures to estimate the incidence in workers exposed to 1 ppm 8 hr/day for a 35-year

working life, the upper limit of the current Occupational Safety and Health Act standard. These predictions are shown in Table 5 for all four Models.

Models C and E predict an incidence comparable to that experienced by workers exposed to much higher levels, again revealing their inadequacy because of overprediction.

Model B, although a linear model, predicts zero response at 93.7 ppm VC. Thus, exposures to levels below this are not predicted to induce hepatic angiosarcoma.

Using what appears to be the most reliable model (Model A, Probit 5), the predicted incidence of hepatic angiosarcoma is 1.5×10^{-6} , or 1.5 cases in 100,000,000 workers.

It is emphasized that use of Models A, C, and D to predict the incidence of hepatic angiosarcoma in workers exposed to 1 ppm VC assumes no threshold or even an inflection point in the dose-response curve.

In summary, the use of animal data for the assessment of risk in humans exposed to a chemical is not as mystical as frequently assumed. Dose-response data for the induction of angiosarcoma in rats exposed to various levels of VC together with attendant biotransformation data have been

used to estimate, reliably, the risk of developing angio-sarcoma in persons exposed to VC.

There is every reason to believe that similar data can and should be used in the assessment of risk of exposure to specified levels of other substances.

R&S 101268

REFERENCES

- 1) Barbin, A., Brévil, H., Créisy, A., Jacquignon, P., Malaveille, C., Montesano, R., and Bartoch, H. (1975). Liver-microsome-mediated formation of alkylating agents from vinyl bromide and vinyl chloride. Biochem. and Biophys. Res. Comm., 67, 596-603.
- 2) Bolt, H. M., Kappus, H., Kaufmann, R., Appel, K. E., Buchter, A., and Bolt, W. (1975). Metabolism of ^{14}C -vinyl chloride in vitro and in vivo. Inserm, Vol 52, pp. 151-164.
- 3) Bolt, H. M., Kappus, H., Buchter, A., and Bolt, W. (1976). Disposition of (1,2- ^{14}C) vinyl chloride in the rats. Arch. Toxicol. 35, 153-162.
- 4) Cresch, J. L. and Johnson, M. N. (1974). Angiosarcoma of liver in the manufacture of polyvinyl chloride. J. Occup. Med., 16, 150-151.
- 5) Fox, A. L. and Collier, P. F. (1977). Mortality experience of workers exposed to vinyl chloride monomer in the manufacture of polyvinyl chloride in Great Britain. Brit. J. Ind. Med., 34, 1-10.

- 6) Gehring, P. J., Watanabe, P. G., and Park, C. H. (1978). Resolution of dose-response toxicity data for chemicals requiring metabolic activation: Example--vinyl chloride. Toxicol. Appl. Pharmacol. (In press).
- 7) Kappas, H., Bolt, H. M., Buchter, A., and Bolt, W. (1976). Liver microsomal uptake of (14 C) vinyl chloride and transformation to protein alkylating metabolites in vitro. Toxicol. Appl. Pharmacol., 37, 461-471.
- 8) Makk, L., DeImore, F., Creech, J. L., Ogden, L. L., Fadell, E. H., Songster, C. L., Clanton, J., Johnson, M. N., and Christopherson, W. H. (1976). Clinical and morphologic effects of hepatic angiosarcoma in vinyl chloride workers. Cancer, 37, 149-163.
- 9) Malavielle, C., Bartsch, H., Barbin, A., Camus, A. M., and Montesano, R. (1975). Mutagenicity of vinyl chloride chloroethyleneoxide, chloroacetaldehyde and chloroethanol. Biochem Biophys. Res. Comm., 63, 363-370.
- 10) Maltoni, C. (1975). The value of predictive experimental environmental carcinogenesis. An example: vinyl chloride. Ambio, 4, 18-23.

- 11) Maltoni, C. and Lefemine, G. (1975). Carcinogenicity assays of vinyl chloride: Current results. Ann. N.Y. Acad. Sci., 2446, 1 5-224.
- 12) Rannar, U., Johansson, A., Ramel, C., and Wachtmeister, C. A. (1974). The mutagenicity of vinyl chloride after metabolic activation. Ambio, 3, 194-197.
- 13) Tabershaw, I. R. and Gaffey, W. R. (1974). Mortality study of workers in the manufacture of vinyl chloride and its polymers. J. Occup. Med., 16, 509-513.
- 14) Watanabe, P. G., McGowan, G. R., Madrid, E. O., and Gehring, P. J. (1976a). Fate of ^{14}C -vinyl chloride following inhalation exposure in rats. Toxicol. Appl. Pharmacol., 37, 49-59.
- 15) Watanabe, P. G., McGowan, G. R., and Gehring, P. J. (1976b). Fate of ^{14}C -vinyl chloride after single oral administration in rats. Toxicol. Appl. Pharmacol., 36, 339-352.
- 16) Watanabe, P. G., Hefner, R. E., Jr., and Gehring, P. J. (1976c). Vinyl chloride induced depression of hepatic nonprotein sulphydryl content and effects on bromo-sulphthalein (BSP) clearance in rats. Toxicology, 6, 1-8.

- 17) Watanabe, P. G., Zempel, J. H., Pegg, D. G., and Gehring, P. J. (1973) Hepatic macromolecular binding following exposure to vinyl chloride. Toxicol. Appl. Pharmacol. (In press).

TABLE 1 - PREDICTED INCIDENCE OF ANGIOSARCOMA IN RATS EXPOSED TO VINYL CHLORIDE USING VARIOUS MODELS VERSUS DOSE (v, μ g VC METABOLIZED PER DAY) COMPARED TO EXPERIMENTAL RESULTS

Exposure ppm ^a	Dose v, μ g/day ^b	Experimental ^a %	Predicted % ^c			
			Model A	Model B	Model C	Model D
10,000	5521	14.8 (9/61)	19.8	20.0	19.7	18.9
6,000	5403	21.7 (13/60)	19.3	20.5	19.3	18.6
2,500	5030	22.0 (13/59)	17.9	18.1	17.9	17.4
500	3413	11.9 (7/59)	12.1	11.8	12.2	12.2
250	2435	6.8 (4/59)	8.1	8.0	8.7	8.8
50	739	1.7 (1/59)	.4	1.4	2.6	2.8

a) From Maltoni and Lefemine (1975)

b) Calculated from equation $v = \frac{5706(\mu\text{gVC}/4 \text{ hr}) \cdot S(\mu\text{g}/\text{l})}{866(\mu\text{g}/\text{l}) + S(\mu\text{g}/\text{l})}$, where

$S = 2.56(\mu\text{g}/\text{ppm}/\text{l}) \times \text{Exposure, ppm}$; see Gehring, et al, 1978.

c) Model A, Probit $z = -1.625 + 1.543 \log (\text{Dose})$; Model B, $z = -1.48 + 0.389 \times 10^{-2} (\text{Dose})$;
Model C, $z = 0.3565 \times 10^{-2} (\text{Dose})$; Model D, $z = \{1 - e^{-z(\text{dose})}\} 100$ where $z = 0.38 \times 10^{-4}$.

TABLE 2 - INCIDENCE OF ANGIOSARCOMA IN 9677 WORKMEN EXPOSED TO VINYL CHLORIDE^a

<u>Population</u>	<u>Duration of Exposure, Yrs</u>	<u>Mean Exposure Duration, Yrs^b</u>	<u>Fraction of Working Life^c</u>	<u>Observed Angiosarcomas</u>
4384 (0.31) ^d	<4	2	0.06	1 (18 Yrs) ^e
2339 (0.27)	5-9	7	0.20	0
946 (0.58)	10-14	12	0.34	2 (15 & 23 Yrs)
1007 (1.0)	15-19	17	0.49	2 (18 & 19 Yrs)
677 (1.0)	20-24	22	0.63	0
324 (1.0)	25+	27	0.77	0
9677				

- a) From "Epidemiological Study of Vinyl Chloride Workers", Final Report January 1978, prepared by Equitable Environmental Health, Inc., Rockville, MD, for Manufacturing Chemists Association, Washington, DC.
- b) Assumed mean duration of exposure.
- c) Mean duration of exposure divided by 35 years.
- d) Fraction of population incurring first exposure 15 or more years prior to study.
- e) Duration between first exposure and diagnosis.

R&S 101275

TABLE 3 - AVERAGE THEORETICAL RAT EQUIVALENT DAILY DOSES OF BIOTRANSFORMATION PRODUCTS RECEIVED BY A 70 KG MAN EXPOSED TO 200 OR 500 PPM VINYL CHLORIDE FOR VARIOUS FRACTIONS OF AN ASSUMED 35-YEAR WORKING LIFE

Mean Exposure Duration, Yrs.	Fraction of Working life	v, $\mu\text{g}/8 \text{ hr}$ Averaged over 35 hrs ^a (ppm)	
		500	200
2	0.06	60	38
7	0.20	200	125
12	0.34	341	213
17	0.49	491	306
22	0.63	631	394
27	0.77	771	481

a)
$$v = \frac{1675 \mu\text{g}/8\text{hr} \cdot S_{\text{mg}/\%}}{360 \mu\text{g}/\% + S_{\text{mg}/\%}} \times (\text{Fraction of working life})$$

TABLE 4 - PREDICTED NUMBER OF ANGIOSARCOMAS IN WORKMEN EXPOSED TO 500 OR 200 PPM VINYL CHLORIDE 8 HOURS/DAY, 5 DAYS/WEEK FOR VARIOUS FRACTIONS OF A 35-YEAR WORKING LIFE VERSUS THE NUMBERS OBSERVED USING BIOTRANSFORMATION AND ANGIOSARCOMA INCIDENCE DATA FROM RATS AND FOUR MODELS FOR EXTRAPOLATION^a

Population	Angiosarcomas Observed	Angiosarcomas Predicted ^b							
		Model A		Model B		Model C		Model D	
		500	200	500	200	500	200	500	200
4384	1	0.2	0.0	0	0	9.4	5.9	10.0	6.3
2339	0	2.4	0.3	0	0	16.7	10.4	17.7	11.1
946	2	3.0	1.1	0	0	11.5	7.2	12.1	7.6
1007	2	6.8	2.7	4.0	0	17.6	11.0	18.6	11.7
677	0	7.3	3.0	6.6	0.3	15.2	9.5	16.0	10.1
324	0	4.9	2.1	4.9	1.3	8.9	5.6	9.4	5.9
9677	5	25	10	16	2	79	50	84	53

a) Observed number of angiosarcomas from "Epidemiological Study of Vinyl Chloride Workers", Final Report January 1978, prepared by Equitable Environmental Health, Inc., Rockville, MD, for Manufacturing Chemists Association, Washington, DC.

b) Model A, Probit $\lambda = -1.025 + 1.543 \log(\text{Dose})$; Model B, $\lambda = -1.48 + 0.388 \times 10^{-2}(\text{Dose})$; Model C, $\lambda = 0.3565 \times 10^{-2}(\text{Dose})$; Model D, $\lambda = \{1 - e^{-\beta(\text{dose})}\} 100$ where $\beta = 0.38 \times 10^{-4}$.

The constants in these Models were obtained using the biotransformation and angiosarcoma incidence data for rats exposed to vinyl chloride, Gehring, et al (1978).

TABLE 5 - PREDICTED INCIDENCE OF ANGIOSARCOMA IN WORKMEN EXPOSED TO 1 PPM VINYL CHLORIDE 8 HOURS/DAY, 5 DAYS/WEEK FOR 35 YEARS USING BIOTRANSFORMATION AND ANGIOSARCOMA INCIDENCE DATA FROM STUDIES IN RATS^a.

A) Probit 2 or Logarithm Probability

$$1.5 \times 10^{-6}\% \text{ or } 1.5/100,000,000$$

B) Linear Not Forced Through Origin

None predicted below exposures of 98.7 ppm.

C) Linear Forced Through Origin

$$1.7 \times 10^{-2}\% \text{ or } 1.7/10,000 \text{ or } 177/1,000,000$$

D) One Hit (CAG)

$$1.89 \times 10^{-2}\% \text{ or } 1.9/10,000 \text{ or } 189/1,000,000$$

^a Rat equivalent daily dose of biotransformation products received by workmen exposed for 8 hours to 1 ppm vinyl chloride is:

$$V = \frac{1675 \mu\text{g/hr} \cdot 2.56 \mu\text{g/g}}{866 \mu\text{g/g} + 2.56 \mu\text{g/g}} = 4.97 \mu\text{g/8 hr}$$

INDEX TERMS

Risk Assessment

Vinyl Chloride

Extrapolation

Angiosarcoma

Rats

Humans

R&S 101278